

**Nonstandard Analysis Applied to
Advanced Undergraduate Mathematics**
◇ Infinitesimal Modeling and Very Elementary Physics ◇

Robert A. Herrmann

A (July - August) 1989, 1990, 1991 Instructional Development Project from the
Mathematics Department
United States Naval Academy
572C Holloway Road
Annapolis, Maryland 21402-5002

IMPORTANT NOTICE

Since the writing of this book was financed entirely by a designated grant from the Federal Government that was specifically obtained for this sole purpose then a copyright for this specific book cannot be obtained by its author. Any portion of its contents can be copied and used without seeking permissions from the author. However, when such copying or use is made of this material, it is necessary that the author and the U. S. Naval Academy be indicated as the source of the material being used. All typographical error have NOT be corrected. Further note that certain new results that appear in this book will be published under the author's name in scholarly journals.*

*This publication process is now being instituted with certain results published as of January 1992, accepted for publication or in preparation. Further, some typographical errors have been corrected as of this 17 June 1997 version.

CONTENTS

Chapter 1

Introduction

1.1 A Brief History	4
1.2 Manual construction	8

Chapter 2

Infinitesimals, Limited And Infinite Numbers

2.1 Some Notation and Definitions	10
2.2 Basic Algebra	12
2.3 Euclidean N-spaces	13
2.4 The Standard Part Operator	13
2.5 A Slight Quandary	15

Chapter 3

Some Set Theory, Convergence And Leibniz's Principle

3.1 Some Set Theory	16
3.2 Convergence	18
3.3 Constants and More Constants	19
3.4 The Leibniz Principle of *-transfer	21
3.5 A Few Simple Applications	23

Chapter 4

Some Modeling With The Infinite Numbers

4.1 Historical Confusion	26
4.2 The Internal Definition Principle	27
4.3 Hyperfinite Summation	30
4.4 Continuity and a Few Examples	32

Chapter 5

Standard Rules For Integral Modeling

5.1 The Riemann Styled Integral	35
5.2 The Infinite Sum Theorems	37
5.3 Extensions	39
5.4 Applications and The Standard Modeling Rules	54
5.5 Extensions of the Standard Rules	39

Chapter 6

Nonstandard Rules For Integral Modeling

6.1 Historical Examples	44
6.2 The Monadic Environment	46
6.3 Simple Applications	49
6.4 The Method of Constants	50
6.5 The Hyperfinite Method	54

6.6	Instruction	55
6.7	Realism	56

Chapter 7

Pure Infinitesimal Integral Modeling

7.1	Brief Discussion	58
7.2	Geometric Elements	59

Chapter 8

Refinements For Integral Modeling

8.1	A Very General Approach	65
8.2	The Line Integral	65
8.3	Order Ideals and Approximations	68
8.4	n th Order Increments	70
8.5	Microgeometry - Tangents to Curves	71
8.6	Microgeometry - Surface Elements	72
8.7	Microgeometry - Other Stuff	74
8.9	Gauge Integrals	75

Chapter 9

The Beginnings Of Differential Equation Modeling

9.1	Brief Discussion	76
9.2	The Limit	77
9.3	Fluxions and Dynamic Geometry	77
9.4	Fluxions and Higher Order Infinitesimals	81
9.5	What is a Tangent?	82
9.6	What is an (k-surface) Osculating Plane?	84
9.7	Curvature	86

Chapter 10

The Differential And Physical Modeling

10.1	Basic Properties	88
10.2	Some General Observations	91
10.3	Vibrating Membrane	92
10.4	Internal Heat Transfer	98
10.5	Concluding Remarks	101

Appendices

For Chapter 2	103
For Chapter 3	106
For Chapter 4	115
For Chapter 5	120
For Chapter 6	127
For Chapter 7	136
For Chapter 8	137
For Chapter 10	133

References	149
Special Symbols	151
Very Elementary Physics	154
Index for pages 1-153	183
 Applications	
Length Of A Continuous Curve (Ex. 4.4.4.A)	33
Fractals (Ex. 4.4.2)	33
The 2-dimensional Area Between Two Curves	40
Volume Obtained By A 2-dimensional Integral	41
Mass Obtained By 3-dimensional Integral	41
Jordan-Volume Obtained By A 2-dimensional Integral	42
Incompressible Fluid	49
Moment Of Inertia	50
Incompressible Fluid Second Derivation	52
Moment Of Inertia Second Derivation	53
The 2-dimensional Area Between Two Curves (Single Integral)	59
Volume Of Revolution (Single Integral)	61
Surface Area of Revolution (Single Integral)	61
Volume Obtained By 2-dimensional Integral	63
Value Of An Electric Field Vector	64
Energy Expended Within a Force Field	67
Tangents To Curves	83
The Osculating Plane	84
Curvature	86
Vibrating Membrane	93
Internal Heat Transfer	98

Chapter 1.

INTRODUCTION

1.1 A Brief History.

Scientists who use mathematical analysis as a tool have traditionally relied upon a vague process called “infinitesimal reasoning” - a process that from the time of Archimedes until 1961 had no fixed rules nor consistent language. However, application of this intuitive process is the exact cause that has led to our great analytical successes both in scientific and engineering endeavors. Unfortunately, it also led to great controversy.

Beginning in about 1600 a schism developed between some mathematicians and the foremost appliers of this analytical tool. Leibniz approved entirely of the concept of the infinitely small or infinitely large numbers but stated that they should be treated as “ideal” elements rather than real numbers. He also believed that they should be governed by the same laws that then controlled the behavior of the ordinary numbers. He claimed, but could not justify the assertion, that all arguments involving such ideal numbers could be replaced by arguing in terms of objects that are large enough or small enough to make error as small as one wished. De l’Hospital [1715] when he wrote the first Calculus textbook used the terminology exclusively and utilized a formal “definition - axiom” process supposedly delineating the notion of the infinitesimal. Unfortunately, his first axiom is logically contradictory. D’Alembert insisted that the Leibniz concepts were without merit and only a process using a modified “limit” idea was appropriate. Euler contended in opposition to D’Alembert that the Leibniz approach was the best that could be achieved and fought diligently for the acceptance of these ideal numbers.

Due to what appeared to be logical inconsistencies within the methods, those mathematicians trained in classical logic began to demand that applied mathematicians produce “proofs” of their derivations. In answer to this criticism Kepler wrote, *“We could obtain absolute and in all respects perfect demonstrations from the books of Archimedes themselves, were we not repelled by the thorny reading thereof.”* The successes of these vague methods and those scientists and mathematicians such as Leibniz, Euler and Gauss who championed their continued use quieted the “unbelievers.” It should be noted that the concern of the critics was based upon the fact that they used the same vague processes and terminology in their assumed rigorous demonstrations.

The major difficulty was the fact that mathematicians had not as yet developed a precise language for general mathematical discourse, nor had they even decided upon accepted definitions for such things as the real numbers. Within their discussions they conjoined terms such as “infinitely small” with the term limit in the hopes of bringing some logical consistency to their discipline.

The situation changed abruptly in 1821. Cauchy, the foremost mathematician of this period, is believed by many to be the founder of the modern limit concept that was eventually formalized by Weierstrass in the 1870’s. A reading of Cauchy’s *Cours d’Analyse (Analyse Algébrique)*[1821] yields the fact, even to the causal observer, that he relied heavily upon this amalgamation of terms and in numerous cases utilized infinitesimal reasoning entirely for his “rigorous” demonstrations. He claimed to establish an important theorem using his methods - a theorem that Abel [1826] showed by a counterexample to be in error. No matter how mathematicians of that time period described their vague infinitesimal methods they failed to produce the appropriately altered theorem - a modified theorem that is essential to Fourier and Generalize Fourier Analysis.

Beginning in about 1870, all of the language and methods of infinitesimal reasoning were replaced in the mathematical textbooks by the somewhat nonintuitive approximation methods we

term the “ $\delta - \epsilon$ ” approach. These previous difficulties are the direct causes that have led to the modern use of axiom techniques and the great linguistic precision exhibited throughout modern mathematical literature.

However, scientists and engineers continued to use the old incorrect infinitesimal terminology. As an example, Max Planck wrote in his books on theoretical mechanics that *“a finite change in Nature always occurs in a finite time, and hence resolves into a series of infinitely small changes which occur in successive infinitely small intervals of time.”* He then attempts to instruct the student in how to obtain mathematical models from this general description. Unfortunately, at that time, such terms as “infinitely small” had no mathematical counterpart.

In many textbooks that claim to bridge the gape between abstract analysis and applications, students often receive the impression that there is no consistent and fixed method to obtain applied mathematical expressions and indeed it takes some very special type of “intuition” that they do not possess. In fact, Spiegel in his present day textbook “Applied Differential Equations” writes the following when he discusses how certain partial differential equations should be “derived.” He states that rigorous methods should not be attempted by the student, but *“it makes much more sense, however, to use plausible reasoning, intuition, ingenuity, etc., to obtain such equations and then simply postulate the equations.”*

In 1961, Abraham Robinson of Yale solved the infinitesimal problem of Leibniz and discovered how to correct the concept of the infinitesimal. This has enabled us to return to the more intuitive analytical approach of the originators of the Calculus. Keisler writes that this achievement *“will probably rank as one of the major mathematical advances of the twentieth century.”* Robinson, who from 1944 - 1954 developed much of the present supersonic aerofoil theory, suggested that his discovery would be highly significant to the applied areas. Such applied applications began in 1966, but until 1981 were confined to such areas as Brownian motion, stochastic analysis, ultralogic cosmogonies, quantum field theory and numerous other areas beyond the traditional experience of the student.

In 1980, while teaching basic Differential Equations, this author was disturbed by the false impression given by Spiegel in the above quotation relative to the one dimensional wave equation. It was suggested that I apply my background in these new infinitesimal methods and find a more acceptable approach. The approach discovered not only gives the correct derivation for the n-dimensional general wave equation but actually solves the d’Alembert - Euler problem and gives a fixed derivation method to obtain the partial differential equations for mechanics, hydrodynamics and the like. These rigorous derivation methods will bridge the gape between a student’s laboratory, and basic textbook descriptions for natural system behavior, and the formal analytical expressions that mirror such behavior. Indeed, slightly more refined procedures can even produce the relativistic alteration taught in modern physics. Moreover, pure nonstandard models are now being used to investigate the properties of a substratum world that is believed to directly or indirectly effort our standard universe. These include pure nonstandard models for the fractal behavior of a natural system, nonstandard quantum fields, a necessary and purely nonstandard model for a cosmogony (or pregeometry) that generates many different standard cosmologies as well as automatically yielding a theory of ultimate entities termed subparticles.

The major goal for writing this and subsequent manuals is to present to the faculty, and through them to the student, these rigorous alterations to the old infinitesimal terminology so that the student can once again benefit from the highly intuitive processes of infinitesimal reasoning - so that they can better grasp and understand exactly why infinitesimal models are or are not appropriate and

when appropriate why they predict natural system behavior. Except for the basic calculus and the more advanced areas, there are no textbooks nor any properly structured documentation available which presents this material at the undergraduate level. In my opinion it will be 15 to 20 years, if not much longer, before such material is available in the commercial market and instructors properly trained. An immediate solution to this problem will give your students a substantial advantage over their contemporaries at other institutions and place your institution in the forefront of what will become a major worldwide trend in mathematical modeling.

1.2 Manual Construction.

The basic construction of these manuals will be considerably different from the usual mathematical textbook. No proofs of any of the fundamental propositions will appear within the main body of these manuals. However, all propositions that do not require certain special models to establish are proved within the various appendices. A large amount of attention is paid to the original intuitive approaches as envisioned by the creators of the Calculus and how these are modified in order to achieve a rigorous mathematical theory.

Another difference lies in the statements of the basic analytical definitions. Many definitions are formulated in terms of the original infinitesimal concepts and not in terms of those classical approximations developed after 1870. Each of these definitions is shown, again in an appropriate appendix, to be equivalent to some well-known “ $\delta - \epsilon$ ” expression. Moreover, since these manuals are intended for individuals who have a good grasp of either undergraduate analysis or its application to models of natural system behavior then, when appropriate, each concept is extended immediately to Euclidean n -spaces.

Nonstandard analysis is NOT a substitute for standard analysis, it is a necessary rigorous extension. Correct and efficient infinitesimal modeling requires knowledge of both standard and non-standard concepts and procedures. Indeed, the nonstandard methods that are the most proficient utilize all of known theories within standard mathematics in order to obtain the basic properties of these nonstandard extensions. It is the inner play between such notions as the standard, internal and external objects that leads to a truly significant comprehension of how mathematical structures correlate to patterns of natural system behavior. Our basic approach employs simple techniques relative to abstract model theory in order to take full advantage of all aspects of standard mathematics. The introduction of these techniques is in accordance with this author’s intent to present the simplest and direct approach to this subject.

Since it is assumed that all readers of these manuals are well-versed in undergraduate Calculus, then your author believes that it is unnecessary to follow the accepted ordering of a basic Calculus course; but, rather, he will, now and then, rearrange and add to the standard content. This will tend to bring the most noteworthy aspects of infinitesimal modeling to your attention at the earliest possible moment. I have this special remark for the mathematician. *These manuals are mostly intended for those who apply mathematics to other disciplines. For this reason, many definitions, proofs and discussions are presented in extended form. Many would not normally appear in a mathematician’s book since they are common knowledge to his discipline. Some would even be considered as “trivial.” Please be patient with my exposition.*

It has taken 300 years to solve what has been termed “The problem of Leibniz” and it should not be assumed that the solution is easily grasped or readily obtained. You will experience some startling new ideas and encounter procedures that may be foreign to you. Hopefully, experience, intuition and knowledge are not immutable. It is my firm belief that, though proper training, these

three all important aspects of scientific progress can be expanded in order to reveal the true, albeit considerably different, mathematical world that underlies all aspects of rigorous scientific modeling. It has been hoped for many years that individuals who have a vast and intuitive understanding of their respective disciplines would learn these concepts and correctly apply them to enhance their mathematical models. It is through your willingness to discard the older less productive, and even incorrect, modeling language that this goal will eventually be met.

Chapter 2.

**INFINITESIMALS, LIMITED
AND INFINITE NUMBERS**

2.1 Some Notation and Definitions.

When Robinson [1961] first introduced his new concepts he used both abstract algebraic and logic notions. A few years later, for simplicity in exposition, the basic algebraic ideas were expanded and, indeed, many of these algebraic results appear here for the first time. Utilizing some fundamental facts about algebraic structures the general properties of the infinitesimals can be readily obtained. Your author has successfully used this approach while instructing a series of 400 level courses. This algebraic approach is based upon but two assumed requirements, requirements that are established from fundamental set theory in a later section. First, however, we need to recall certain elementary definitions and identify notation.

Throughout these manuals the symbol $\mathbb{R}$ denotes the real numbers which we assume is a complete ordered field.

(I) Assume that there exists another ordered field ${}^*\mathbb{R}$ with the properties that $\mathbb{R}$ is a ordered subfield of ${}^*\mathbb{R}$ (i.e. $\mathbb{R}$'s basic order and field properties are those of ${}^*\mathbb{R}$ but restricted to members of $\mathbb{R}$) **AND** $\mathbb{R} \neq {}^*\mathbb{R}$.

The set ${}^*\mathbb{R}$ is called by various names, **hyperreals**, **star-reals** or **extended reals**. Also let the symbol $\mathbb{N}$ denote the natural numbers (including zero).

The ordered field $\mathcal{F}$ is assumed to contain a copy of non-negative integers (natural numbers) $\mathbb{N}$ formed by adjoining to the additive identity (the zero of $\mathcal{F}$) finite sums of the multiplicative identity (the 1 of $\mathcal{F}$). You can then simply consider $\mathbb{N} \subset \mathcal{F}$, in general. The concept **complete** is discussed in most elementary analysis books where it is almost always shown that *if $\mathcal{F}$ is a complete ordered field and $r \in \mathcal{F}$, then there exists some $n \in \mathbb{N}$ such that $|r| < n$* . This property for a field is called the **Archimedean** property for a field. As will be shown, an important and general algebraic result states that **whatever the hyperreals may be they cannot be Archimedean**.

(II) The second assumption is the following: **assume that there exists some nonzero $\epsilon \in {}^*\mathbb{R}$ such that for each positive $r \in \mathbb{R}$; it follows that**

$$0 < \epsilon < r.$$

Historically, it is unfortunate that abstract algebra was not investigated at a much earlier date since much of the difficulty Leibniz and Euler faced in having the infinitesimals accepted as genuine entities would have been erased.

Theorem 2.1.1. *Assumption (II) holds for ${}^*\mathbb{R}$ if and only if ${}^*\mathbb{R}$ is not Archimedean.*

What Theorem 2.1.1 means is that the more we know about ordered fields that are not Archimedean the better. Of course, such fields also are not complete. It is now possible to define explicitly the “infinitesimals.” Note that we interpret the logical symbol $\wedge$ as the word “and,” the symbol $\forall$ by any one of the expressions “for each, for all, for every” and the symbol $\exists$ by one of the expressions “there exists some, there exists one, there exists an.” Since such logical notation will become significant later in this manual we might as well practice its use as soon as possible. Let $\mathbb{R}^+$ denote the positive real numbers.

Definition 2.1.1 (Infinitesimals). Let $\mu(0) = \{x | (x \in {}^*\mathbb{R}) \wedge \forall r(r \in \mathbb{R}^+ \rightarrow |x| < r)\}$. Or in words, $\mu(0)$ is the set of all hyperreal numbers x such that if r is a positive real number, then the absolute value of x is less than r . The set $\mu(0)$ is called the set of **INFINITESIMALS**. In the literature the set $\mu(0)$ is also denoted by the symbols M_1 and o .

Definition 2.1.2. (Limited). Let $\mathcal{O} = \{x | (x \in {}^*\mathbb{R}) \wedge \exists r(r \in \mathbb{R}^+ \wedge |x| < r)\}$. Or in words, $\mathcal{O}$ is the set of all hyperreal numbers x such that there exists some positive real number r such that $|x| < r$. The set $\mathcal{O}$ is called the set of **LIMITED** numbers. Robinson first called $\mathcal{O}$ the “finite” numbers and denoted them by the symbol M_0 .

The term “limited” is relatively new in the literature and many nonstandard analysts still employ the term “finite.” When this is done, there is some confusion when the term finite is used in the ordinary sense of set theory. The concept of limited can also be interpreted as meaning that each of these hyperreal numbers is “limited by” or “bounded by” a real number. There are many reasons why the term “bounded” would be confused with the same term as employed in standard analysis and for this reason is not used for the concept of limited.

Definition 2.1.3. (Infinite). The set ${}^*\mathbb{R} - \mathcal{O} = {}^*\mathbb{R}_\infty$ is the set of **INFINITE** hyperreal numbers. Or in words, those hyperreal numbers that are not limited are the infinite. The infinite numbers are also called the unlimited numbers.

I point out that the set of infinite numbers can also be characterized analytically as follows: $x \in {}^*\mathbb{R}$ is infinite if and only if for every $r \in \mathbb{R}^+$, $|x| > r$. This characterization is often very useful. The next definition relates the infinitesimals to the original concept of when two numbers are “infinitely close.” This is probably the most significant concept for infinitesimal modeling and deserves attention.

Definition 2.1.4. (Infinitely Close). Two hyperreal numbers x, y are **INFINITELY CLOSE** if $x - y \in \mu(0)$. Or in words, if their difference is an infinitesimal. The symbol used for infinitely close is $\approx$.

How do we extend these definitions for the case of the Euclidean n -spaces, $\mathbb{R}^n$ where we are using the Euclidean norm $\|(x_1, \dots, x_n)\| = \sqrt{x_1^2 + \dots + x_n^2}$?

Definition 2.1.5. (Euclidean Extensions). In Definitions 2.1.1 and 2.1.2 substitute for the absolute value symbol $|\cdot|$ the norm symbol $\|\cdot\|$, as it would be defined relative to ${}^*\mathbb{R}^n$. Of course, we should also substitute the term “vector” or the symbol $\mathbb{R}^n$ when appropriate.

In the next section, the basic algebraic properties for the infinitesimals, limited numbers etc. are explored along with the relationships between the above hyperreal concepts and those of the hyperreal n -spaces.

2.2 Basic Algebra.

You should expect that the basic properties for the above defined entities will be presented rapidly since the proofs only appear in the appendix. This has both advantages and disadvantages. These properties are couched in terms of some very well-known algebraic structures and we need a very brief refresher course relative to field theory.

The ordered fields $\mathbb{R}$, ${}^*\mathbb{R}$ have the weaker structural property of being a **ring**. As far as nonempty subsets of a field $\mathcal{F}$ are concerned *rings may be characterized as any nonempty $A \subset \mathcal{F}$ that are closed under the operations of subtraction and multiplication*. A ring is also closed under addition, has a zero and additive inverses. Rings share with the field itself the right and left distributive laws, and the usual associative and commutative properties used in parentheses manipulation. Rings can differ greatly from a field in that they need not contain an element that has a multiplicative inverse. In the case that $\mathcal{F}$ is ordered, then rings, in general, only share the basic simple order property for the field $\mathcal{F}$ when it is considered to be restricted to the ring. Our last general ring concept is that of the “ideal.” Let the ring $\mathfrak{R} \subset \mathcal{F}$. Then nonempty $\mathfrak{S} \subset \mathfrak{R}$ is an **ideal** of (or in) $\mathfrak{R}$ if $\mathfrak{S}$ is a subring (i.e. a ring with respect to the ring structure of $\mathfrak{R}$ and a subset of $\mathfrak{R}$) and for each $r \in \mathfrak{R}$ and each $x \in \mathfrak{S}$ it follows that $rx \in \mathfrak{S}$. Thus an ideal “absorbs” the members of $\mathfrak{R}$ by multiplication. Before proceeding with our first list of properties notice that the infinitesimals $\mu(0) = \{x | x \in {}^*\mathbb{R} \wedge x \approx 0\}$.

Theorem 2.2.1. *The set of limited numbers, $\mathcal{O}$, is a subring of ${}^*\mathbb{R}$, $\mathbb{R} \subset \mathcal{O}$ and $\mathcal{O}$ is not a field, but if $x \in \mathcal{O} - \mu(0)$, then $x^{-1} \in \mathcal{O}$.*

Theorem 2.2.2. *The infinitesimals, $\mu(0)$, form a subring of $\mathcal{O}$ and Γ is infinite if and only if there is some nonzero $\epsilon \in \mu(0)$ such that $\Gamma = 1/\epsilon$.*

Theorem 2.2.3. *The set of infinitesimals, $\mu(0)$, is an ideal of $\mathcal{O}$.*

The fact that $\mu(0)$ forms an ideal of $\mathcal{O}$ is very significant for the correct theory of infinitesimals. Not only are the infinitesimals closed under finite addition and finite product, but absorb, under product, all of the real numbers. It’s unfortunate that Leibniz and others could not establish such results rigorously since, if they could have, undoubtedly much of the criticism of their concepts would not have developed.

We now consider what happens when we take any real number and add to it the infinitesimals.

Definition 2.2.1. (Monad). Let $r \in \mathbb{R}$. Then a **MONAD OF (ABOUT) r** is the set $\mu(r) = \{x | x \in {}^*\mathbb{R} \wedge (x - r) \in \mu(0)\} = \{x | x \in {}^*\mathbb{R} \wedge x \approx r\}$.

Theorem 2.2.4. *The binary relation $\approx$ is an equivalence relation on ${}^*\mathbb{R}$.*

Theorem 2.2.5. *For each $x, y \in \mathbb{R}$,*

- (i) $\mu(x) \cap \mu(y) = \emptyset$ if and only if $x \neq y$,
- (ii) $\mathcal{O} = \bigcup \{\mu(r) | r \in \mathbb{R}\}$.

Corollary 2.2.5.1. *If $x, y \in \mathbb{R}$, $x < y$, $z \in \mu(x)$, $w \in \mu(y)$, then $z < w$.*

Corollary 2.2.5.2. *If $x, y \in \mu(r)$, $z \in {}^*\mathbb{R}$, $x < z < y$, then $z \in \mu(r)$.*

The important Theorem 2.2.5 can be expressed in words by stating that the set of monads forms an ordered partition of the limited numbers. It is beginning to appear as if the infinitesimals are indeed behaving in the manner first envisioned by the founders of the infinitesimal calculus. HOWEVER, in at least one respect the monads as well as the set of limited numbers do not share an important real number property. From the previous discussion, ${}^*\mathbb{R}$ is known not to be complete. But, are there significant sets that have upper bounds in ${}^*\mathbb{R}$ and do not have a least upper bound?

Theorem 2.2.6. *Each monad and the set of limited numbers are bounded above [resp. below], but do not possess a least upper bound [resp. greatest lower bound].*

2.3 Euclidean n-spaces.

The following are additional propositions that extend some of the above results to Euclidean n-spaces. Whenever possible the same notation is used and should be understood from the context. It should be obvious how the previous definitions extend to Euclidean n-spaces.

Theorem 2.3.1. *For any $n \in \mathbb{N}$, $n \geq 1$, the set of limited vectors in ${}^*\mathbb{R}^n$ is equal to*

$$\mathcal{O}^n = \overbrace{\mathcal{O} \times \cdots \times \mathcal{O}}^{n \text{ factors}}.$$

Theorem 2.3.2. *For any $n \in \mathbb{N}$, $n \geq 1$, then monad of $\vec{v} = (x_1, \dots, x_n) \in \mathbb{R}^n$ is equal to*

$$\mu(\vec{v}) = \overbrace{\mu(x_1) \times \cdots \times \mu(x_n)}^{n \text{ factors}}.$$

Theorem 2.3.3. *For any $n \in \mathbb{N}$, $n \geq 1$, and for each $\vec{v}, \vec{w} \in \mathbb{R}^n$,*

- (i) $\mu(\vec{v}) \cap \mu(\vec{w}) = \emptyset$ if and only if $\vec{v} \neq \vec{w}$,
- (ii) $\mathcal{O}^n = \bigcup \{\mu(\vec{v}) | \vec{v} \in \mathbb{R}^n\}$.

2.4 The Standard Part Operator

In infinitesimal modeling, various methods exist that allow us to investigate what might be termed as the *micro-effects* that occur within a world called the *Nonstandard Physical World*, (i.e. *NSP-world*) or *Deductive World* (i.e. *D-world*). When these micro-effects are modeled by means of the infinitesimals, certain mathematical operators applied to such infinitesimals yield standard mathematical objects that are often perceived to measure the corresponding *natural world* (i.e. *N-world*) effects that govern phenomenological behavior. The next operator is probably the most significant of these standardizing processes.

Definition 2.4.1. (Standard Part). Let $x \in \mathcal{O}$. Define the function $\text{st}: \mathcal{O} \rightarrow \mathbb{R}$ as follows:

- (i) let $\text{st}(x) = r$, where $r \in \mathbb{R}$ is the unique real number such that $x \in \mu(r)$.
 - (ii) The function st is often extended to all of ${}^*\mathbb{R}$ by letting $\text{st}(x) = +\infty$, when $x \in {}^*\mathbb{R} - \mathcal{O}$ and $x > 0$ or $\text{st}(x) = -\infty$, when $x \in {}^*\mathbb{R} - \mathcal{O}$ and $x < 0$. The map st is called the **standard part operator**.
-

Since $\mathbb{R} \subset \mathcal{O}$ then, as will be seen, the range of st is $\mathbb{R}$ (i.e. st is a surjection.) Please note the basic properties of the standard part operator are highly important in elementary infinitesimal analysis.

Theorem 2.4.1. Let $x, y \in \mathcal{O}$. Then

- (i) $x \approx y$ if and only if $\text{st}(x) = \text{st}(y)$,
- (ii) $x \approx \text{st}(x)$,
- (iii) if $x \in \mathbb{R}$, then $\text{st}(x) = x$,
- (iv) if $x \leq y$, then $\text{st}(x) \leq \text{st}(y)$,
- (v) if $\text{st}(x) \leq \text{st}(y)$, then either $x \leq y$ or $x - y \in \mu(0)$ with unknown order.

Part (iii) of Theorem 2.4.1 is what implies that the range of $\text{st} = \mathbb{R}$. The next theorem is established from the basic definitions and the fact that $\mu(0)$ is an ideal of $\mathcal{O}$. The map st is purely algebraic in character and its application is a remarkable indication of how pure abstract algebra can be utilized to obtain classical results. For algebraists, these pure algebraic characterizations are discussed following the next theorem.

Theorem 2.4.2. Let $x, y \in \mathcal{O}$. Then

- (i) $\text{st}(x \pm y) = \text{st}(x) \pm \text{st}(y)$,
- (ii) $\text{st}(xy) = (\text{st}(x))(\text{st}(y))$.

Corollary 2.4.2.1 Let $x, y \in \mathcal{O}$. Then

- (i) if $\text{st}(y) \neq 0$, then $\text{st}(x/y) = \text{st}(x)/\text{st}(y)$.
- (ii) if $y = \sqrt[n]{x}$, then $\text{st}(y) = (\text{st}(x))^{(1/n)}$, where it is always the case that if $x \in \mathcal{O}$, then $\sqrt[n]{x} \in \mathcal{O}$.

Theorem 2.4.3. For $\vec{v} \in \mathcal{O}^n$, let $\text{st}(\vec{v}) = (\text{st}(x_1), \dots, \text{st}(x_n))$. The set $\mathcal{O}^n$ forms a vector space with respect to the ring $\mathcal{O}$ (i.e. a module) and as such the map st distributes over the vector space algebra as well as the component defined dot and for $n = 3$ the cross product operators.

Notice that even though $\mu(0)$ is an ideal of $\mathcal{O}$ it is most definitely only a subring of ${}^*\mathbb{R}$. Indeed, products are somewhat indefinite. For example, let $0 \neq \epsilon \in \mu(0)$. Then $\epsilon^2 \in \mu(0)$. But $\epsilon(\epsilon^{-2}) \in {}^*\mathbb{R} - \mathcal{O}$, while $\epsilon(1/\epsilon) \in \mathcal{O}$ and $\epsilon^2(1/\epsilon) \in \mu(0)$.

[This is for the algebraist.] The above theorems show that the mapping st is a ring epimorphism with $\ker(\text{st}) = \mu(0)$.

Theorem 2.4.4. *The set $\mu(0)$ is a maximal ideal in $\mathcal{O}$ and the quotient ring $\mathcal{O}/\mu(0)$ is isomorphic to $\mathbb{R}$.]*

2.5 A Slight Quandary.

In 1961 when Robinson first published his new theory, it was restricted to subsets of ${}^*\mathbb{R}$ and functions defined on ${}^*\mathbb{R}$. Shortly after this Luxemburg [1962] expanded upon Robinson's paper but still retained the same general restrictions. It became obvious that any extensive applications of infinitesimal analysis to functional analysis would require an extension to other set-theoretic objects. Robinson [1966] did just this but at a great expense to elementary exposition. He used a purely logical approach and the simplified theory of types. Indeed, this author entered this field in 1968 and first used the type-theoretic approach. There is no doubt that the type-theoretic approach is beyond almost all mathematicians and appliers of mathematics that have not had specialized training. Machover and Hirschfeld [1969] introduced a simplification to the Robinson type-theoretic approach by restricting their nonstandard analysis to set theory itself. Unfortunately, their approach requires that individuals utilize two "set theories" - a pseudoset theory and the standard set theory. Your author wrote his doctoral dissertation in pseudoset theory. At approximately the same time Robinson and Zakon [1969] published a paper that further simplified Robinson's approach. They were able to show how set theory itself could be used for all the necessary formations of nonstandard analysis and that pseudoset theory was no longer necessary. Since 1969 there have been other attempts at simplifying the foundations of nonstandard analysis but it is this author's belief that they are not appropriate to those mathematicians who are not accustomed to special logical procedures.

The quandary I face is that there are no other appropriate simplifications of Robinson's theory except for the 1969 Robinson - Zakon approach. This approach definitely needs to be presented within this basic manual so that you can encounter the full power of infinitesimal analysis. The difficulty is my wish not to present numerous definitions and constructions from mathematical logic in the next chapter; but, rather, actually to do some infinitesimal analysis immediately. There is a procedure that does allow this if we postpone until a later chapter the reasons why this procedure functions.

Chapter 3.

**SOME SET THEORY, CONVERGENCE
AND LEIBNIZ'S PRINCIPLE**

3.1 Some Set Theory.

Historically we have over 5000 years of standard mathematics available. It is certainly reasonable to utilize all of this standard mathematics in the development of what has become known as “nonstandard analysis”. The term “nonstandard” should not be taken to mean that the properties to be discussed are not relative to the known properties for standard mathematics. Indeed, the opposite is the case. In the development of nonstandard mathematics, it has always been required that nonstandard structures have all the same ordinary properties as the standard mathematical structures. They are nonstandard in the sense that they have additional useful properties not possessed by the standard structures. The simplification procedures mentioned at the conclusion of the last chapter have, for the most part, eliminated certain model theoretic difficulties associated with Robinson’s application of the simplified theory of types. Depending upon the type of set-theoretic entity used no special considerations need to be considered. Even though what is about to be presented seems very straightforward and is easy to grasp, it should not be assumed that these set-theoretic procedures hold in the exact form given for more extensive set-theoretic processes that are not specifically mentioned. [Note: The formal set-theory used is found in **Suppes [1960]**.]

Why do we need to consider basic set theory at all? Are not the properties of the infinitesimals, limited and infinite numbers sufficient for analysis? Didn’t Leibniz stress the calculus of the infinitesimal and infinite numbers as the basis for analysis? Since mathematicians investigate the *relations between individual objects* and our simplification is set-theoretical then basic set theory is required. Indeed, even an order pair (a, b) is set-theoretically defined as $(a, b) = \{\{a, b\}, \{a\}\}$.

Theorem 3.1.1. *It is permissible to assume that*

- (i) if $A \subset \mathbb{R}$, then $A \subset {}^*\mathbb{R}$,
- (ii) if $A \subset \mathbb{R}^n$, then $A \subset {}^*\mathbb{R}^n$,
- (iii) if $A \subset (\mathbb{R}^n) \times (\mathbb{R}^m)$, then $A \subset ({}^*\mathbb{R}^n) \times ({}^*\mathbb{R}^m)$.

Notice that the subsets considered in Theorem 3.1.1 exhaust almost all of the mathematical objects studied in an ordinary course in undergraduate analysis and much more. However, at present, the notion of a “constant” in a formal mathematical language needs to be discussed. From the viewpoint of a formal language, it is assumed that every subset of $\mathbb{R}$, $\mathbb{R}^n$ or $(\mathbb{R}^n) \times (\mathbb{R}^m)$ is associated with at least one constant symbol that “names” that subset. These constants are employed in two contexts. First, as language constants and nothing more, then as names for mathematical objects within our special set theory. Technically these are two different usages which are not usually mentioned in introductory texts. In infinitesimal analysis these technicalities are significant, however. In the next theorem, the notation ${}^*A$ also has two meanings. ${}^*A$ is a constant symbol in a mathematical language and, in most cases, represents a mathematical entity different from A .

Theorem 3.1.2. *If $A \subset \mathbb{R}$ [resp. $\mathbb{R}^n$, $(\mathbb{R}^n) \times (\mathbb{R}^m)$], then there exists ${}^*A \subset {}^*\mathbb{R}$ [resp. ${}^*\mathbb{R}^n$, $({}^*\mathbb{R}^n) \times ({}^*\mathbb{R}^m)$] such that*

- (i) $A \subset {}^*A$ and
- (ii) $A = {}^*A$ if and only if A is finite.

Thus Theorem 3.1.2. tells us that a collection of new sets exist - the “star-sets” - and obviously we need a certain amount of new terminology in order to discuss their properties effectively. The basic operators that define the ordered field $\mathbb{R}$ may also be considered as subsets of $\mathbb{R} \times \mathbb{R}$ or $\mathbb{R}^3$. Do we then consider these operators extended to $^*\mathbb{R}$ and thus “star” the operator such as writing $a \text{ } ^*+ \text{ } b$, where $a, b \in ^*\mathbb{R}$? If one wants to be technical about the matter, then the answer would be yes. However, in practice these specific operators are not so denoted. On the contrary, one may consider the * -field operators as basic and that $\mathbb{R}$ is defined by restricting them to the set $\mathbb{R}$. In all cases, the particular defining field operator is determinable from the context.

In all that follows let $\mathcal{U} = \mathbb{R} \cup \mathbb{R}^n \cup (\mathbb{R}^n) \times (\mathbb{R}^m)$. Now a major consideration in the writing of these manuals relates to mathematical rigor. Even though all stated propositions are established in the appendixes, how explicit should the main text be when confronted with the basic construction of a nonstandard model for analysis? Analysis of the fundamental objects in $\mathcal{U}$ often requires collections of members from $\mathcal{U}$ that are themselves not members of $\mathcal{U}$. In order to satisfy the modern trend that rejects pronouncements that are not at least superficially established, a slight acquaintance with the actual construction of the standard model for this analysis seems appropriate. It is definitely not essential that you have any in depth knowledge of the fine details of the construction of this standard structure. But, be assured that every standard set-theoretic object that will ever be consider in this analysis is a member of the standard structure.

For any set W in our set theory, let $\mathcal{P}(W)$ denote the set of all subsets of W , where the operator $\mathcal{P}$ is often called the **power set operator**. Hence $A \in \mathcal{P}(W)$ if and only if $A \subset W$. Starting with $\mathbb{R} = X_0$ an object called a **superstructure** is constructed by induction as follows: assume that X_n has been defined for $n \in \mathbb{N}$. Then define $X_{n+1} = \mathcal{P}(\bigcup\{X_i \mid i \leq n\})$. Then the standard structure - **the superstructure** - is the set $\mathcal{H} = \bigcup\{X_n \mid n \in \mathbb{N}\}$. Various properties relative to $\mathcal{H}$ are discussed in Appendix 3.

Definition 3.1.1. (Individual, Entity, Star-Sets). Each member of $\mathbb{R}$ is called an **INDIVIDUAL**, while each set in $\mathcal{H}$ is called an **ENTITY** and for each entity A , the set $^*A$ is called an **EXTENDED STANDARD** set.

It turns out that for any $W \in \mathcal{H}$ the set $\mathcal{P}(W) \in \mathcal{H}$ and there exists the set $^*\mathcal{P}(W)$. {What appears between these two braces is a refinement for what appeared in this place originally. It clears up a certain notational confusion. The star operator “ * ” behaves like a mapping from $\mathcal{P}(W)$ into $^*\mathcal{P}(W)$. As such, there is a slight confusion in symbols that “name” the images. The usual notation for the range would be $^*(\mathcal{P}(W))$. The same notation holds for $A \in \mathcal{P}(W)$, where $^*(A)$ is the image notation. In order to relate this notation to our original when “ * ” is considered as an mapping, we let $^*A$ be a name for the image $^*(A)$. That is $^*(A) = ^*A$.}

The next few propositions reveal the fundamental behavior of * and the first shows that * at least preserves all finite set-theoretic operations.

Theorem 3.1.3. *Unless otherwise stated all constants represent individuals or entities.*

- (i) $a \in A$ if and only if $^*a \in ^*A$; $A \neq B$ if and only if $^*A \neq ^*B$.
- (ii) $A \subset B$ if and only if $^*A \subset ^*B$.
- (iii) $^*\{A_1, \dots, A_k\} = \{^*A_1, \dots, ^*A_k\}$.
- (iv) $^*(A_1, \dots, A_k) = (^*A_1, \dots, ^*A_k)$.
- (v) $(A_1, \dots, A_k) \in A$ if and only if $(^*A_1, \dots, ^*A_k) \in ^*A$.

(vi) Let $A, B \subset \mathcal{U}$. Then $*(A \cup B) = *A \cup *B$, $*(A \cap B) = *A \cap *B$, $*(A - B) = *A - *B$, $*(A \times B) = *A \times *B$, $*\emptyset = \emptyset$.

(vii) If $A \in \mathbb{R}$, then $*A = A$.

Recall that if R is any n -ary relation ($n > 1$), then the i **th projection** is the set $P_i(R) = \{x_i | (x_1, \dots, x_i, \dots, x_n) \in R\}$, where $1 \leq i \leq n$. Also since $R \subset A^n$ if and only if $*R \subset (*A)^n$ then $*R$ is any n -ary relation in $(*A)^n$ and the set-theoretic projections are defined for $*R$. If R is an binary relation, then P_1 is called the **domain** and P_2 the **range** of R .

Theorem 3.1.4. Let $\emptyset \neq R \subset A^n$, ($n > 1$),

(i) $*P_i(R) = P_i(*R)$.

(ii) If R is a binary relation (i.e. $n = 2$), then $*R^{-1} = (*R)^{-1}$.

The next theorem presents two technical results and is stated here for completeness.

Theorem 3.1.5. Let A be an entity. Then

(i) $*\{(x, x) | x \in A\} = \{(x, x) | x \in *A\}$.

(ii) $*\{(x, y) | x \in y \in A\} = \{(z, w) | z \in w \in *A\}$.

Theorems 3.1.3, 3.1.4 can be paraphrased by simply considering the $*$ map to be a natural map which appears to distribute over finite “everything.”

[This is for the algebraist. The map $*$ when restricted to $\mathcal{P}(W)$ is a homomorphism on the Boolean algebra $\mathcal{P}(W)$. The map $*$ is a monomorphism and it is a significant fact that it is not an epimorphism.]

3.2 Convergence.

Since the time of Zeno [350 BC] the concept of the non-finite has been somewhat controversial. In modern axiomatic set theory, except for considerations of consistency, such controversies have little meaning. However, in physical modeling, accepting the reality of the non-finite has numerous philosophical ramifications. Unfortunately, these manuals are not the appropriate form to discuss these side issues. We have mentioned the concept from an intuitive set-theoretic point of view and this will suffice since infinitesimal analysis eliminates, to a great degree, much of the more esoteric and philosophic concerns that plagued early mathematical discourse.

Leibniz [1701] discussed what he claimed to be a relation between his calculus of infinitesimal and infinite numbers and the “method of Archimedes” (the so-called method of exhaustion). He, of course, never established this claim and did not seem to use any definable approximation process such as the “ $\delta - \epsilon$.” He mentions such things as fractions with infinite numbers as denominators, infinite sequences and the like. In particular, Leibniz, de l’Hospital and even Cauchy specifically required, as axiomized by de l’Hospital, that “a curved line be considered an infinite assemblage of straight line segments each infinitely small.” This concept should be distinguished from the completely different idea of a curve as composed of an infinite number of points, with no magnitude, - the *indivisibles* - as put forth by Cavalieri [1635]. Leibniz also instituted a notion he called the “sovereign principle.” This principle served as his justification that the infinitesimals and his infinite numbers obey the

same rules as the ordinary real numbers. As will be seen, many of these early ideas were not correct from the rigorous view point and, moreover, Robinson's methods are very broad in context since they may be applied to all standard mathematical structures not merely to $\mathbb{R}^n$.

Even though such statements as *a sequence converges to a real number "as n goes to infinity or as n grows without bound or as n gets infinitely large"* and other such intuitive expressions may be eliminated entirely from our language of convergence and replaced with statements involving the behavior of extended standard functions that are property defined on $^*\mathbb{N}$, it is a misconception that the formal limit definition with its approximating techniques is no longer needed. From the view point of mathematical modeling, nonstandard and standard analysis complement each other. The effects that appear in the natural world are modeled by standard analysis. The limit approach, with all of its approximation techniques, is often considered to be a natural world approximation for events modeled within the NSP-world by nonstandard objects. Our basic definition for sequential convergence is stated in a language that parallels some of Leibniz thoughts; but, as established in appendix 3 it is equivalent to the Bolzano-Weierstrass concept. **In all that follows, let $\mathbb{N}_\infty$ denote the infinite natural numbers.**

Definition 3.2.1. (Sequential Convergence). Let the sequence $S: \mathbb{N} \rightarrow \mathbb{R}^n$, ($n \geq 1$). Then S converges to $\vec{r} \in \mathbb{R}^n$ (notation $S \rightarrow \vec{r}$), if for each $\Gamma \in \mathbb{N}_\infty$ it follows that $^*S(\Gamma) \in \mu(\vec{r})$. Thus S converges to $\vec{r} \in \mathbb{R}^n$ if $^*S$ maps all of the infinite natural numbers into a single monad (i.e. $^*S[\mathbb{N}_\infty] \subset \mu(\vec{r})$). Recall that the value $S(n)$ of a sequence is often written as S_n .

Theorem 3.2.1. *Definition 3.2.1 (i.e. $S \rightarrow \vec{r}$) is equivalent to the limit definition for convergence of a sequence (i.e. $\lim_{n \rightarrow \infty} S(n) = \vec{r}$).*

The next result simply recasts definition 3.2.1 in terms of the algebraic standard part operator. However, this seemingly "trivial" fact yields immediately all of the basic "limit theorems" that tend to be a first stumbling block for the first year calculus student.

Theorem 3.2.2. *Let the sequence $S: \mathbb{N} \rightarrow \mathbb{R}^n$, ($n \geq 1$). Then $S \rightarrow \vec{r} \in \mathbb{R}^n$ if and only if for each $\Gamma \in \mathbb{N}_\infty$, $\text{st}(^*S_\Gamma) = \vec{r}$.*

Notice that when definition 3.2.1 is stated without the symbolism then it closely parallels the conceptional process outlined by Leibniz. **A sequence converges to a real number r if its value for each infinite number is infinitely close to r .** Thus Leibniz and others explained the result that the sequence $S(n) = n^{-2}$ converges to 0 by asserting that if Γ is an infinite number, then Γ^2 is an infinite number. This implies that Γ^{-2} is an infinitesimal and thus $^*S(\Gamma)$ is infinitely close to 0. (They did not use the * idea, however. And, when no confusion results, the * is often omitted from the extended standard function notation.) Unfortunately, in order to apply such intuitive procedures one must know how the * -function behaves. In the next section, such an investigation is initiated.

3.3 Constants and More Constants.

In section 3.2, it was pointed out that Leibniz believed that the infinitesimal and infinite numbers "obey the same rules as the ordinary real numbers." It was necessary to develop abstract model

theory before Leibniz's vague comment could be rigorously justified. As previously discussed, every object in $\mathfrak{R} = \mathcal{U} \cup \mathcal{P}(\mathcal{U})$ is denoted by a “constant” symbol as is every object in $\mathcal{H}$. When theorems about the set $\mathcal{U} \cup \mathcal{P}(\mathcal{U})$ are written in terms of these constants, variables and the symbols ϵ and $=$, an intuitive “first-order language” is often used. Indeed, every theorem and definition that appears in this manual has been stated or can be reformulated in such a “language.” Here are a few examples of how these intuitive expressions appear, where all the constants are assumed to represent members of $\mathcal{H}$.

Example 3.3.1. In our set theory you have the basic definition for the union of two sets $A, B \subset \mathfrak{R}$. This might be expressed as follows: for each $x \in \mathfrak{R}$, $x \in A \cup B = C$ if and only if $x \in A$ or $x \in B$. This can also be expressed in a formalized first-order statement as $\forall x(x \in \mathfrak{R} \rightarrow (x \in A \cup B = C \leftrightarrow (x \in A) \vee (x \in B)))$.

In the above first-order statement, it is required that the variable symbol x be restricted to members of $\mathfrak{R}$. *The absolute requirement is that all quantified language variables must be restricted to specific sets that are entities (i.e. to sets that are elements of various X_p) and that are represented by constants within our language.* Mostly, the variables may simply be restricted to the set $\mathfrak{R}$, with certain exceptions mention in future chapters. However, better comprehension is often produced when the language variables are restricted to sets of immediate interest.

Example 3.3.2. Suppose that you have a sequence that is strictly increasing. How might this be stated? For each x and y if $x, y \in \mathbb{N}$ and $x < y$, then $S(x) < S(y)$.

Example 3.3.3. Since the usual operator and relation symbols have been retained as constant symbols, it is a simple matter to give explicit functional statements. For each $n \in \mathbb{N}$, $S(n) = \sin(n^2)$. Notice that the symbol n has now been used as a variable and that the symbol $\sin$ is a constant that represents the sine function.

Obviously, in order to formulate properly these first-order expressions it is essential that individuals gain facility with the basic language of mathematics. In particular, great precision must be maintained in both written and oral mathematical exposition. This is also one of the goals of the core mathematics program at the Academy.

Are there significant mathematical concepts that might not be expressible by such first-order statements? Suppose you wanted to express the general concept of mathematical induction as put forth by Peano - a concept considered to be one of the most basic in all of mathematics. *If Q is any property which may or may not hold for the natural numbers, and if (1) the natural number 0 satisfies property Q and (2) whenever a natural number n satisfies property Q , then $n + 1$ has property Q , then all natural numbers satisfy property Q .* The difficulty in expressing the induction axiom lies in the requirement that we express the phrase “ Q is any property.”

There are two ways of expressing a property within a first-order language. First, a property is expressed by other first-order statements or secondly, it may be claimed that a property can be represented by a member of $\mathbb{N}^m$. In the first case, we need a variable that, at least, represents

predicates in our language and this is not allowed in a first-order language. A third approach would be to replace the single induction axiom by infinitely many axioms where each axiom refers to one specific property expressed by one specific first-order expression. Unfortunately, it can be shown that you would not obtain all of the possible properties by this process.

For the second set-theoretic case, it turns out that all of the possible properties for the natural numbers still cannot be obtained since we do not have a set in our slightly restricted set theory that contains all of the objects that represent all of the properties that can be expressed by our first-order set-theoretic language. Further, using the basic language of set theory one can conceive of a property determined by the predicate $P(x) = “\emptyset \subset x,”$ where $\emptyset$ (the empty set) is the object that satisfies the set-theoretic expression: there exists a set x such that for all sets y , $y \notin x$. However, for set theory to be consistent it must be assumed usually that if there does exist something that satisfies this property, it is not to be called a set. Consequently, one must be very sure that an informally stated notion is indeed expressible by means of our simplistic first-order language, even though it may not be done within an informal discussion or argument.

Every object in $\mathcal{H}$ is denoted by a logical constant. Some of these constants are in the customary form such as $\mathbb{N}$, $\mathbb{R}$, 2 , $\sqrt{2}$, $+$, $\sin$ and all the very well-known symbols used in standard analysis. As an abbreviation, let $C(\mathcal{H})$ denote the set all such constants. As previously indicated, in nonstandard analysis the set ${}^*\mathcal{H} = \bigcup \{ {}^*X_n | n \in \mathbb{N} \}$ is constructed and it is assumed that each object in ${}^*\mathcal{H}$ is denoted by a constant. The set of all such constants is denoted by $C({}^*\mathcal{H})$. In particular, it follows that ${}^*\mathbb{R} = {}^*\mathcal{U} \cup {}^*\mathcal{P}(\mathcal{U})$ and each object in ${}^*\mathbb{R}$ is also denotable by a constant. (Note: ${}^*(\mathcal{P}(\mathcal{U}))$ is denoted by ${}^*\mathcal{P}(\mathcal{U})$ since the symbol $\mathcal{P}(\mathcal{U})$, in this form, is considered as one constant.) Now if $A \in C(\mathcal{H})$, it denotes an object in $\mathcal{H}$ and the notation ${}^*A$ is the constant that denotes the object in ${}^*\mathcal{H}$ obtained by means of the * map. **Also note that Theorems 3.1.2 and 3.1.3 indicate that some members of ${}^*\mathcal{H}$ are denotable by both a starred and unstarred constant and by a previous convention the basic field operators defined on ${}^*\mathbb{R}$ are not starred. In these cases, we tend to use only the unstarred notation.** Theorems 3.1.2 and 3.1.3 imply that there are infinitely many members of ${}^*\mathcal{H}$ that are not named by starred members of $C(\mathcal{H})$. But, nevertheless, they do have constant names in an extended language.

Example 3.3.4. We know that $\mathbb{N}_\infty \neq \emptyset$. Moreover, ${}^*\mathbb{N} \subset {}^*\mathcal{U}$ implies that each member of ${}^*\mathbb{N}$ is a member of ${}^*\mathbb{R}$ and $\mathbb{R}$ is a member of some X_p . We have used capital Greek letters as the names for some of the members of ${}^*\mathbb{N}_\infty$. By the * process we only obtain the stars of the natural numbers, where by convention we drop the * . Since the * process is one-to-one into ${}^*\mathcal{H}$ by Theorem 3.1.3 then no member of ${}^*\mathbb{N}_\infty$ is the * of any of the constants in $C(\mathcal{H})$.

3.4 The Leibniz' Principle of * -transfer.

Thus far it may not appear that the previous rules and conventions associated with infinitesimal analysis yield a simplification. However, I invite you to compare the second chapter of Robinson's 1966 book with above procedures and judge for yourself. But, why do we need to consider these specialized first-order languages in the first place? The reason lies within certain very powerful results from the subject area of mathematical logic. These fundamental propositions coupled with our conventions lead to a completely correct formulation of the Leibniz Principle, which now becomes a theorem. But, first, one final procedure needs to be discussed prior to formalizing this highly useful principle.

Definition 3.4.1. (*-transfer). Let Φ be an intuitive first-order sentence written with respect to the above rules and only containing constants from $C(\mathcal{H})$ and the basic set-theoretic binary operators $\in, =, \cup, \cap, \subset, \times, -$ etc. Then $^*\Phi$ is the same sentence as Φ except every constant that appears in Φ is now proceeded by a $*$. In this process, the conventions as to when the $*$ may be dropped are also followed.

The sentence $^*\Phi$ is called the ***-transform** of the sentence Φ . The mathematical object $^*\mathcal{H}$ is actually obtained by application of the compactness theorem for a formal first-order language or by an algebraic construction called the ultraproduct construction [Stroyan and Luxemburg [1976]] and a process known as the Mostowski collapse [Barwise [1977], Herrmann [1986]]. The Leibniz Principle is an immediate consequence of the fundamental definition of what it means to say that $\mathcal{H}$ and $^*\mathcal{H}$ are the universes for first-order models. You can learn about first-order models by referring to any good book in mathematical logic.

Theorem 3.4.1 (The Leibniz' Principle). *A sentence Φ holds true for members of $\mathcal{H}$ if and only if the sentence $^*\Phi$ holds true for members of $^*\mathcal{H}$.*

What Theorem 3.4.1 says is that if we let K be the set of all of those specially written sentences that hold true for members of $\mathcal{H}$, then the members of $^*\mathcal{H}$ along with the basic set-theoretic operators form a model for the set of all sentences obtained from K by $*$ -transfer. *Probably much more significant is the fact that infinitely many other statements not obtainable by $*$ -transfer hold true for members of $^*\mathcal{H}$.* Each member of $^*\mathcal{H}$ is called an **internal** entity (or internal individual if it is a member of $^*\mathbb{R}$). Observe that an object A is internal if and only if there exists some $p \in \mathbb{N}$ such that $A \in ^*X_p$. Consequently, the extended standard sets are internal entities; but, there are many internal entities that are not extended standard sets and internal individuals that are not individuals. Shortly the additional terms *extended standard* and *internal* are more fully exploited. But, first a few examples.

Example 3.4.1. Let S_1, S_2 be two sequences. The definition of the addition of two such functions can be stated as follows: for every x if $x \in \mathbb{N}$, then the function $C = S_1 + S_2$ if and only if $C(x) = S_1(x) + S_2(x)$. The $*$ -transform becomes: for every x if $x \in ^*\mathbb{N}$, then the function $^*C = ^*S_1 + ^*S_2$ if and only if $^*C(x) = ^*S_1(x) + ^*S_2(x)$. Notice that such forms as $^*C(x)$ mean that the $*$ is applied to the C only. Thus $*$ -transfer extends to all of the ordinary definitions of this type.

Example 3.4.2. Let's look at a few of the parts of Theorem 3.1.3. (i) Using just constants themselves is legal. Thus if $a, A \in C(\mathcal{H})$, and the expression $a \in A$ holds for $\mathcal{H}$, then the $*$ -transform is simply the expression $^*a \in ^*A$. (iii) Notice that if $\{A_1, \dots, A_n\}$ is a finite set of members of $\mathcal{H}$, then there is a constant C in $C(\mathcal{H})$ that represents this set. Clearly, it is unnecessary to mention continually that a symbol is a constant on one hand and a name for an object in $\mathcal{H}$ on the other. The context of a statement will usually serve to indicate a constants usage. This first-order statement is: for every x , $x \in C$ if and only if $x = A_1 \vee x = A_2 \vee \dots \vee x = A_k$. This is a collection of finitely many symbols and is thus allowed. The $*$ -transfer becomes: for every x , $x \in ^*C$ if and only if $x = ^*A_1 \vee x = ^*A_2 \vee \dots \vee x = ^*A_k$. [Note: translate $\vee$ by the word "or."]

[Note: Our defined operators $+$, $\cdot$, $|\cdot|$, $\|\cdot\|$, etc. are usually considered but the $*$ -transfer of the operators (the nonstandard extension to ${}^*\mathbb{R}$, ${}^*\mathbb{R}^n$) as they are defined on $\mathbb{R}$ and $\mathbb{R}^n$.]

Example 3.4.3. An argument. $*$ -transfer is certainly important, but it is our ability to argue by using statements that contain symbols that are not obtainable by $*$ -transfer which is the key to nonstandard analysis. Indeed, as will be established, there are three different categories of constant mathematical symbols used in the following argument. Let $\Gamma \in {}^*\mathbb{N} - \mathbb{N}$. [$\Gamma \in C({}^*\mathcal{H})$ and there is no symbol for ${}^*\mathbb{N} - \mathbb{N}$ in $C(\mathcal{H})$.] Since $\Gamma \notin \mathbb{N}$ [it turns out that $\mathbb{N} \notin {}^*\mathcal{H}$] then for each $x \in \mathbb{N}$, $x < \Gamma$ for if not, then there exists some $n \in \mathbb{N}$ such that $0 < \Gamma < n$ and for each $x \in \mathbb{N}$, $x \neq \Gamma$. However, we also know that the set $\{x | 0 < x < n\}$ is a finite set and thus each member is an element of $\mathbb{N}$ (even under $*$ -transfer by our conventions). This would yield a contradiction. Now since $\mathbb{R}$ [$\mathbb{R} \in C(\mathcal{H})$] is Archimedean then for each $r \in \mathbb{R}$ there exists some $n \in \mathbb{N}$ such that $|r| < n$. Hence, $|r| < \Gamma$ implies that $\Gamma \in \mathbb{N}_\infty$. Therefore, ${}^*\mathbb{N} - \mathbb{N} \subset \mathbb{N}_\infty$. Since it is obvious that $\mathbb{N}_\infty \subset {}^*\mathbb{N} - \mathbb{N}$ then this implies that $\mathbb{N}_\infty = {}^*\mathbb{N} - \mathbb{N}$.

Theorem 3.4.2. *The set of infinite natural numbers $\mathbb{N}_\infty = {}^*\mathbb{N} - \mathbb{N}$.*

Mathematicians are so practiced in arguments such as illustrated in example 3.4.3 that it is often not realized that there are at least five categories of constants as symbols being used. (1) There are the **standard (unstarred) constants** that appear in $C(\mathcal{H})$. (2) The **extended standard constants** that appear in $C({}^*\mathcal{H})$. (3) The **internal constants** that appear in $C({}^*\mathcal{H})$ but are not obtained by $*$ -transfer. (4). Then **external constants**, such as $\mathbb{N}_\infty$ that represent mathematical objects but do not fit categories (1) (2) (3). Last, there are the constants the comprise that word forms of the “metalanguage” that discusses and establishes things about the mathematical objects, such as the phrase “there is not a symbol for.” Theorem 3.1.2 and part (vii) of Theorem 3.1.3 indicate the only instances where two categories - (1) and (2) - overlap in the sense that there are definitely two distinct constants representing the same mathematical object. With respect to the next definition, these basic facts about the usage of the constants - especially their correspondence to various mathematical objects - should always be kept in mind.

Definition 3.4.2. (Internal, External.) Any member of ${}^*\mathcal{H}$ will be termed, in general, an **INTERNAL ENTITY** or **INTERNAL INDIVIDUAL**. Any $A \subset {}^*X_p$ that IS NOT INTERNAL is termed an **EXTERNAL ENTITY**.

Definition 3.4.2 now allows for a somewhat better but equivalent statement of the Leibniz Principle for objects such as $\mathbb{R}$.

Theorem 3.4.3. (Leibniz Principle Restated). *A sentence Φ holds true for members of $\mathbb{R} = \mathcal{U} \cup \mathcal{P}(\mathcal{U})$ if and only if the sentence ${}^*\Phi$ holds true for the members of ${}^*\mathcal{U}$ or internal subsets of ${}^*\mathcal{U}$.*

3.5 A Few Simple Applications.

Using infinitesimal procedures, there are simple and specifically describable methods for determining the limit of a sequence, where Theorem 3.2.1 allows us to use the expression “limit of a

sequence” as an abbreviation for the phrase “real number to which the a sequence converges.” The next applications amply illustrate the most significant of these procedures.

Application 3.5.1. *This is an example of the direct application of the algebraic properties of the infinite and infinitesimal numbers.*

Let $p \in \mathbb{N}$, $p > 0$ and assume that for each $0 < n \in \mathbb{N}$, $S_n = (1/n)^p$. Then $S \rightarrow 0$.

Proof. Let $\Gamma \in {}^*\mathbb{N} - \mathbb{N}$. Then $(1/\Gamma) \in \mu(0)$ implies that $(1/\Gamma)^p \in \mu(0)$. Thus result follows from Definition 3.2.1.

Application 3.5.2. *A bounding method.*

Let $a \in \mathbb{R}$, $0 < |a| < 1$. For each $0 < n \in \mathbb{N}$, let $S_n = a^n$. Then $S \rightarrow 0$.

Proof. By induction it is not difficult to show that if $0 \leq b \in \mathbb{R}$, then for each $n \in \mathbb{N}$, $(1+b)^n \geq (1+nb)$. Let $b = (1/|a|) - 1$. Then $b > 0$ and $|a| = 1/(1+b)$. Thus for $n \in \mathbb{N}$,

$$0 < (|a|)^n = \frac{1}{(1+b)^n} \leq \frac{1}{1+nb} < \frac{1}{bn}, \quad (n > 0).$$

Thus by *-transfer for each $n \in {}^*\mathbb{N}$

$$0 < (|a|)^n < (1/b)(n)^{-1}, \quad (n > 0).$$

In particular, for each $\Gamma \in {}^*\mathbb{N} - \mathbb{N}$,

$$0 < (|a|)^\Gamma < (1/b)(\Gamma)^{-1}.$$

The result now follows since $a^\Gamma \in \mu(0)$.

Application 3.5.3 *A bounding method, redefinition and the standard part operator.*

Let $1 < a \in \mathbb{R}$. For each $0 < n \in \mathbb{N}$, let $S_n = (a)^{1/n}$. Then $S \rightarrow 1$.

Proof. Define the sequence $Q_n = a^{1/n} - 1$, $0 < n \in \mathbb{N}$. Then

$$a = (1 + Q_n)^n = 1 + nQ_n + \text{other positive terms},$$

where $0 < n \in \mathbb{N}$. Hence $a > nQ_n$, $\forall n \in \mathbb{N}$, $(n > 0)$. By *-transfer $a > nQ_n$, $\forall n \in {}^*\mathbb{N}$ $(n > 0)$. In particular, $a > \Gamma Q_\Gamma$, $\forall \Gamma \in {}^*\mathbb{N} - \mathbb{N}$. Hence

$$\forall \Gamma \in {}^*\mathbb{N} - \mathbb{N}, \quad 0 < Q_\Gamma < a(1/\Gamma) \in \mu(0).$$

Thus $Q \rightarrow 0$ implies that $\text{st}({}^*Q_\Gamma) = 0 = \text{st}({}^*S_\Gamma) - \text{st}(1) = \text{st}({}^*S_\Gamma) - 1$, $\forall \Gamma \in {}^*\mathbb{N} - \mathbb{N}$. The result follows from Theorem 3.2.2.

Application 3.5.4. *A bounding method, redefinition and the standard part operator.*

Let $S_n = \sqrt[n]{n}$, $\forall n \in \mathbb{N}$, $(n > 0)$. Then $S \rightarrow 1$.

Proof. Let $Q_n = S_n - 1 = \sqrt[n]{n} - 1$, $\forall n \in \mathbb{N}$, $(n > 0)$. Then

$$n = (1 + Q_n)^n \geq (n(n-1)/2)Q_n^2, \quad \forall n \geq 2,$$

by the binomial expansion. Hence

$$0 \leq Q_n \leq \sqrt{\frac{2}{n-1}}, \quad \forall n \geq 2.$$

By *-transfer,

$$0 \leq Q_\Gamma \leq \sqrt{\frac{2}{\Gamma-1}}, \quad \forall \Gamma \in {}^*\mathbb{N} - \mathbb{N}.$$

But, $\sqrt{\frac{2}{\Gamma-1}} \in \mu(0)$, $\forall \Gamma \in {}^*\mathbb{N} - \mathbb{N}$ implies that $Q \rightarrow 0$; which implies that $S \rightarrow 1$.

The bounding techniques and algebraic manipulations illustrated by above examples are actually of the same type that were used originally with old style infinitesimal analysis to argue for these results. As previously mentioned the standard part operator establishes the classical limit theorems which, of course, can be applied always. However, the standard part operator can be applied directly without referring to the limit theorem at all.

Application 3.5.5 *Using the standard part operator.*

Find the limit of the sequence $S_n = ((1/n)^{10}) \sqrt[n]{n}$, $n > 0$. Let arbitrary $\Gamma \in {}^*\mathbb{N} - \mathbb{N}$. Then $\text{st}(1/\Gamma) = 0$ implies that $\text{st}((1/\Gamma)^{10}) = 0$. From application 3.5.4, $\text{st}((\Gamma)^{1/\Gamma}) = 1$. Thus $\text{st}({}^*S_\Gamma) = 0 \cdot 1 = 0$. Hence, $S \rightarrow 0$.

One of the most significance statements made by a researcher is “What if...?” The basic propositions of infinitesimal analysis tend to lend themselves to many such “What if...?” type questions. Consider, for example, Theorem 3.2.2. One is prone to ask; what if $\forall \Gamma \in \mathbb{N}_\infty$, $\text{st}({}^*S_\Gamma) \in \mathcal{O}^n$? Or, what if there exists some $\Gamma \in \mathbb{N}_\infty$ and $\text{st}({}^*S_\Gamma) \in \mathcal{O}^n$? We state a theorem that represents an interesting result relative to the last “What if ...?” - a result established in appendix 3 solely by means of the standard part operator.

Theorem 3.5.1. *Let $S: \mathbb{N} \rightarrow \mathbb{R}$. If S is an increasing [resp. decreasing] sequence and there exists some $\Gamma \in \mathbb{N}_\infty$ such that ${}^*S_\Gamma \in \mathcal{O}$, then $S \rightarrow \text{st}({}^*S_\Gamma)$.*

Chapter 4.

**SOME MODELING WITH
THE INFINITE NUMBERS**

4.1 Historical Confusion.

First, I point out that the infinite numbers are completely different from the idea of extending the real numbers by adjoining the new objects $\pm\infty$ to $\mathbb{R}$ and impressing upon these objects certain topological and algebraic properties. This can be readily shown by considering the concept of when $\lim_{n \rightarrow \infty} = +\infty$. Let $\mathbb{R}_{\infty}^+$ denote the positive infinite hyperreal numbers.

Definition 4.1.1. ($S \rightarrow +\infty$). Let $S: \mathbb{N} \rightarrow \mathbb{R}$. Then $S \rightarrow +\infty$ if for each $\Gamma \in \mathbb{N}_{\infty}$, ${}^*S(\Gamma) \in \mathbb{R}_{\infty}^+$.

Theorem 4.1.1. *Definition 4.1.1 is equivalent to the limit definition for a sequence $S \rightarrow +\infty$.*

Theorem 4.1.1 indicates that the values of ${}^*S$ are scattered throughout the set $\mathbb{R}_{\infty}^+$.

Theorem 4.1.2. *Let $r \in {}^*\mathbb{R}^+$, $S: \mathbb{N} \rightarrow \mathbb{R}$ and $S \rightarrow +\infty$. Then for each $\Omega \in \mathbb{N}_{\infty}$ there exists some $\Delta \in \mathbb{N}_{\infty}$ such that ${}^*S(\Omega) + r \leq {}^*S(\Delta)$.*

Thus the sequential property that $S \rightarrow +\infty$ does not correspond to the idea that a sequence might “converge” to an infinite number. Indeed, we have not even mentioned the idea of a monad about an infinite number even though it is possible to extend monad theory to cover such cases. This is particular significant for applied modeling when the infinite series is discussed relative to the sequence of partial sums each infinitely small.

Recall that except for modern times most mathematics was almost used exclusively for applied physical or geometric modeling and was not consider as a study of abstract entities. De l’Hospital apparently believed in the objective reality of both the infinitesimal and infinite quantities as well as the existence of sets that contained infinitely many members. To him, they existed in the natural world. When de l’Hospital wrote that it was a requirement that one must regard a curve as a totality of an infinity of straight line segments, each infinitely small: or “(which is the same) as a polygon with an infinite number of sides, each infinitely small, which determine by the angle at which they meet, the curvature of the curve...” he apparently meant that all of these intuitively expressed objects exist in reality. Leibniz did not approve of this interpretation. Over and over again, Leibniz proclaimed that such objects as the infinitesimal or infinite numbers were “ideal” or imaginary. He claimed that they are theoretically useful but that they did not correspond to real natural things. Moreover, Leibniz apparently accepted the concept of a potentially infinite set as well, rather than the possibility of an objectively real infinite set of objects. Almost all of the ancient controversy as to the reality of such notions apparently came about as a direct result of the basic philosophical and theological predilections of the investigators. Their general philosophic belief systems were reflected in both their scientific and mathematical views.

With respect to the physical possibility of there being objectively real objects that are characterized by infinitesimals or even infinite numbers, Robinson has replace these abstruse philosophical

considerations with the following observation. In the first fundamental paper delineating his theory, he wrote: “*For phenomena on a different scale, such as are considered in Modern Physics, the dimensions of a particle may not be observable directly. Accordingly, the question whether or not a scale of non-standard analysis is appropriate to the physical world really amounts to asking whether or not such a system provides a better explanation of certain observable phenomena than the standard system of real numbers. The possibility that this is the case should be borne in mind.*” [Fine Hall, Princeton University] Robinson [1961]

As previously mentioned Zeno proposed his famous physical paradox of Achilles’ and the Tortoise in their never ending (potentially infinite time) foot race. It is claimed, that this paradox is resolved by modern mathematics through application of the infinite series. Such a series can supposedly model this physical foot race - a statement that is obviously false. Such a series would require the non-mythological object, the Tortoise, and the mythological Achilles to have variable “sandal” sizes - sizes that decreased to the point of being unmeasurably small.

De l’Hospital accepted a curve as being identifiably the same as a polygon with a fixed infinite number of line segments comprising its sides. On the other hand, Eudoxus [370 BC] devised the method of exhaustion which assumes the true existence of a finite sequence of inscribed and circumscribed polygons. In general, for a closed non-polygonal curve none of these Eudoxus polygons were considered to be the curve under investigation; but, rather, by the “continuity process” they would continually squeeze the curve between these two types of polygons and “exhaust” the space in between. By this process the length of a curved segment was conceived of as an intuitive sequence composed of portions of the polygon’s perimeters. Thus developed the idea of a partial sum that represented the sum of the lengths of the sides of an n -gon - a finite sum that remained finite but acquired more and more terms. Those that employed this method often guessed at a specific formula then justified their guess by indirect and not direct argument. In modern times, Planck described the accepted procedure for modeling the behavior of a natural system when he wrote that: “*a finite change in Nature always occurs in a finite time, and hence resolves into a series of infinitely small changes which occur in successive infinitely small intervals of time.*” What might the term “series” mean in Planck’s statement?

Leaving aside the ontological question associated with the notion of the infinite, it will be demonstrated that Robinson’s theory of the infinitesimal and infinite hyperreal numbers brings a concrete and rigorous language to the above vague methods of physical and geometric modeling. Unfortunately, in doing so, many of these previous ideas will require modification.

4.2. The Internal Definition Principle.

Clearly there is a need to acquire a better understanding of the relationship between the concepts of the infinite series, the sequence of partial sums, the internal process of partial summing with its arithmetic and the notion expounded by Planck. But, prior to examining these concepts, yet another technical procedure needs to be discussed.

In definition 3.4.1, our first-order language was extended to include the basic set-theoretic binary operators. As explained in the Appendix to Chapter 3 immediately after the proof of Theorem 3.1.3, our first-order language may also include the symbol $(\cdot, \dots, \cdot)$ for n -tuple formation where the coordinates are either variables or constants. Under $*$ -transfer these operators and the n -tuple formation symbol are not starred in any Φ that has been properly formulated with the variables restricted to entities in $\mathcal{H}$.

Example 4.2.1. Assume that you are given some relation R , a nonempty set A and the

next statement. For each x if $x \in \mathcal{P}(\mathcal{U})$, then $(x, A) \in R$ and $x \subset A$. Or, as a formal statement $\forall x(x \in \mathcal{P}(\mathcal{U}) \rightarrow (x, A) \in R \wedge x \subset A)$. Then the *-transfer would read: For each x if $x \in {}^*\mathcal{P}(\mathcal{U})$, then $(x, {}^*A) \in {}^*R$ and $x \subset {}^*A$.

Please note that we needed to star the symbol $\mathcal{P}(\mathcal{U})$. This would also be the case if a variable appeared where the $\mathcal{U}$ appears.

When a collection C of mathematical expressions are written they include “mathematical” variables and constants. Within our standard set theory the constants represent individuals or entities. From the view point of the mathematical structure, the entities represent such things as basic sets, operators, relations, specific n-tuples and other definable objects. In general, the variables either vary over every member of some set or represent some one unknown element. It is common practice for mathematicians to use different variable symbols to represent elements of distinct sets or distinct positions in n-tuples and the like. Within C certain of these variables may also be “quantified.” What this means is that if v is one of the variables in C , than there also appears in C in the customary location the phrase “for all v ” (i.e. $\forall v$) or “there exists some v ” (i.e. $\exists v$). Any variable in C that is not associated with some quantifier is termed a **free variable**. Now in formal logic the concept of the free and quantifier bounded variable must be more carefully described since formally the same variable can appear both in a free and not free position. Since we are trying to be as non-technical as possible and since it is customary to use many different variable symbols in mathematical prose, this somewhat vague free variable definition should suffice. If it still is confusing, then most elementary logic books have a more formal presentation.

Example 4.2.2. When the definition of continuity is expressed one usually states the following: The function $f: A \rightarrow \mathbb{R}$ is continuous at $p \in A$ if for each $\epsilon \in \mathbb{R}^+$ there exists some $\delta \in \mathbb{R}^+$ such that whenever $0 \leq |x-p| < \delta$ and $x \in A$, then $0 \leq |f(x)-f(p)| < \epsilon$. The symbols $f, A, \mathbb{R}, \mathbb{R}^+, p, -, |, \leq, <, 0$ are all considered as constants. The symbols ϵ, δ, x are variables. This entire definition may be formally expressed by $\Phi = \forall x(x \in \mathbb{R}^+ \rightarrow \exists y(y \in \mathbb{R}^+ \wedge \forall z(z \in A \wedge 0 \leq |z-p| < \delta \rightarrow |f(z)-f(p)| < y))$.

Suppose that you are interested in the subset of A in example 4.2.2 that comprises all of the points of continuity. In this case, the constant p is considered a variable and one writes such a set as $\{p|p \in A \wedge \Phi(p)\}$. Built into our set builder notation is the quantifier $\forall$. “The set of all $p \in A$ such that $\Phi(p)$ holds true.” The expression $\Phi(p)$ is now considered a formula in one variable p . Note that p is a free variable. Of course, these set builder formulas should present no difficulties since these language constructions are the ordinary and customary ones used by the mathematical community. Two more examples should sufficiently illustrate this easily grasped relation between our first-order language and common mathematical usage.

Example 4.2.3. Most of the time when we write mathematical formula in variables they are considered as free. Such an expression as $y = 3x$ has been defined for many different structures. Even though to obtain the graph of such an expression one needs to know the domain and codomain, this is not necessary until the structure itself is considered. Thus the graph may be the $\{(x, y)|x \in [0, 3] \subset \mathbb{R} \wedge y \in \mathbb{R} \wedge y = 3x\}$. On the other hand, we might also have $\{(x, y)|x \in \mathbb{R}^7 \wedge y \in \mathbb{R}^7 \wedge y = 3x\}$. Thus $y = 3x$ is a mathematical formula, $\Phi(x, y)$, in two variables with a constant 3 and a constant operator (multiplication) that can be used to generate many different sets in our set theory.

Formulas that appear in set builder notation can also contain quantifiers as the next example indicates.

Example 4.2.4. Let the sets $A, B \in X_p$ and let B^A denote the set of all functions with domain A and codomain B . Then $B^A \in X_{p+3}$. One might want to consider a special subset of B^A defined by $\{x|x \in B^A \text{ and there exists } y \in A \text{ such that } x(y) > 3\}$ (i.e. $\{x|x \in B^A \wedge \exists y(y \in A \wedge x(y) > 3)\}$) or $\{x|x \in B^A \wedge \exists y \exists z(y \in A \wedge z \in B \wedge (y, z) \in x \wedge z > 3)\}$. This set exists by the axiom of comprehension and every though the defining expression contains more than one variable, all but the x are bounded by a quantifier that appears immediately to the left.

The basic reason for discussing and presenting these simple examples lies in two most interesting results that allow for the set-theoretic generation of all extended standard or internal sets and n -ary relations by means of such set builder formulas. Of course, n -ary relations are sets but are singled out specifically due to their obvious usefulness. Recall that $C(\mathcal{H})$ denotes the set of all constants that signify members of the set $\mathcal{H}$ while $C(*\mathcal{H})$ the names for members of $*\mathcal{H}$. A formula in our first-order language is called standard [resp. internal] **bound** if each quantified variable is restricted to an object represented by a constant in $C(\mathcal{H})$ [resp. $C(*\mathcal{H})$]. What this means is that, for a $\mathcal{H}$ [resp. $*\mathcal{H}$] interpretation of the formula, each bounded variable must be interpretation as varying over a set contained in $\mathcal{H}$ [resp. $*\mathcal{H}$]. The formula in Example 4.2.2 is bound. A formula like $\forall x \exists y (x \in y)$ is not bound.

Theorem 4.2.1. (The Extended Standard Definition Principle)

(i) A set A in our set theory is an extended standard set (i.e. there exists some $B \in C(\mathcal{H})$ such that $A = *B$) if and only if there exists some standard set D and a standard bound formula $\Phi(x)$ in one free variable where each constant in $\Phi(x)$ is a member of $C(\mathcal{H})$ and

$$A = \{x|x \in *D \wedge *\Phi(x)\}.$$

(ii) A set A is an extended standard n -ary relation ($n > 1$) if and only if there exist n standard sets $D_1, \dots, D_n$ and a standard bound formula $\Phi(x_1, \dots, x_n)$ in n free variables where each constant in $\Phi(x_1, \dots, x_n)$ is a member of $C(\mathcal{H})$ and

$$A = \{(x_1, \dots, x_n)|x_1 \in *D_1 \wedge \dots \wedge x_n \in *D_n \wedge *\Phi(x_1, \dots, x_n)\}.$$

Theorem 4.2.2. (The Internal Definition Principle)

(i) A set A in our set theory is an internal set if and only if there exists some internal set D and an internal bound formula $\Phi(x)$ in one free variable where each constant in $\Phi(x)$ is a member of $C(*\mathcal{H})$ and

$$A = \{x|x \in D \wedge \Phi(x)\}.$$

(ii) A set A is an internal n -ary relation ($n > 1$) if and only if there exist n internal sets $D_1, \dots, D_n$ and an internal bound formula $\Phi(x_1, \dots, x_n)$ in n free variables where each constant in $\Phi(x_1, \dots, x_n)$ is a member of $C(*\mathcal{H})$ and

$$A = \{(x_1, \dots, x_n)|x_1 \in D_1 \wedge \dots \wedge x_n \in D_n \wedge \Phi(x_1, \dots, x_n)\}.$$

Internal objects are basic to nonstandard analysis since it is only internal properties that hold for the model ${}^*\mathcal{H}$. As far as physical modeling is concerned **internal objects represent NSP-world effects that directly or indirectly yield the observed natural world behavior being modeled approximately by a standard mathematical structure.**

4.3. Hyperfinite Summation.

In example 4.2.4 the set-theoretic notation B^A is utilized to denote the set of all functions with domain A and codomain B . As previously mentioned there are procedures that require objects in $\mathcal{H}$ that may not be members of $\mathfrak{R}$. Assume that $\mathcal{A}, \mathcal{B}$ are sets of subsets and $\mathcal{A}, \mathcal{B} \in X_p$. Now let the sets $A \in \mathcal{A}, B \in \mathcal{B}$. Then $A, B \in X_{p-1}$, ($p \geq 2$), $A \cup B \subset X_0 \cup X_{p-2}$ implies that $A \cup B \in X_{p-1}$ and if $a \in A, b \in B$, then $\{a\}, \{a, b\} \in X_{p-1}$. Hence $(a, b) \in X_p$. Thus if $f \in B^A$, then $f \in X_{p+1}$. Indeed, $A \times B \in X_{p+1}$. Therefore, $B^A \in X_{p+2}$. It is possible to consider the formation of each set B^A as an operator $F(x, y)$ where $x \in \mathcal{A}$ and $y \in \mathcal{B}$. The $*$ -transfer process can be extended to this operator in the same manner as is done with the $\cup, \cap, \times, -, (\cdot, \dots, \cdot)$ operators in the sense that it is not starred when written entirely in variable form. For specific members of $C({}^*\mathcal{H})$, if $D \in {}^*\mathcal{A}$ and $E \in {}^*\mathcal{B}$, then ${}^*F(D, E) \in {}^*X_{p+2}$ and ${}^*F(D, E)$ is the internal set of all internal functions with domain D and codomain E . This last fact comes from the $*$ -transfer of a general characterization for the set-theoretic concept of x^y .

Example 4.3.1. Let $p \in {}^*\mathbb{N}$. Then the set $\{x | x \in {}^*\mathbb{N} \wedge 0 \leq x \leq p\} = [0, p]$ is an internal subset of ${}^*\mathbb{N}$. Notice that the symbol $[0, p]$ can be used as an abbreviation for this set's defining property (i.e. $x \in [0, p]$ if and only if $x \in {}^*\mathbb{N} \wedge 0 \leq x \leq p$) which can be restated in an appropriate first-order expression and substituted for the notation $[0, p]$. With this in mind it is clearly possible to now consider p as a variable.

Hence it follows that $BHF = \{y | y \in {}^*\mathcal{P}(\mathbb{N}) \wedge \exists p(p \in {}^*\mathbb{N} \wedge y = [0, p])\}$ is an internal (indeed, an extended standard set) of subsets of ${}^*\mathbb{N}$. The set BHF is called the set of all **basic hyperfinite subsets of ${}^*\mathbb{N}$** .

A nonempty $A \subset B$ is **finite** if there exists some $n \in \mathbb{N}$ and a function $f: [0, n] \rightarrow B$ such that the range of f (i.e. $P_2(f)$) = A . The intuitive idea of a finite set appears to be equivalent to this functional definition if you are willing to accept such things as the “finite” axiom of choice and elementary procedures of recognition. The intuitive idea of the finite is based upon the human recognition of a distinction between symbols written on paper as they are considered geometric forms and consequently it is related to the most basic aspects of concrete geometry. It is this accepted recognition of the differences between geometric forms that than allows one to give a concrete meaning to a correspondence between these forms and $\mathbb{N}$. One does not really establish that such a correspondence exists but its existence is accepted as part of the metamathematical methods.

Theorem 3.1.4 states that for any standard function f , ${}^*(P_i(f)) = P_i({}^*f)$, $i = 1, 2$. The projections P_i can be considered as maps from the set of all nonempty subsets of $A_1 \times \dots \times A_n$ into the sets A_i , where $1 \leq i \leq n$. A modification of proof of Theorem 3.1.4 yields

Theorem 4.3.1. Let $n > 1$. Then for each internal $R \subset {}^*A_1 \times \dots \times {}^*A_n$ and for each i , $1 \leq i \leq n$ it follows that ${}^*P_i(R) = P_i(R)$, where $P_i(R)$ is an internal subset of ${}^*A_i$.

Definition 4.3.1 (Hyperfinite). An internal subset A of a set ${}^*B$ is **hyperfinite** if it is empty or there exists some $[0, p] \in BHF$ and $f \in {}^*F([0, p], {}^*B)$ such that $P_2(f) = A$.

Theorem 4.3.2. Let $F(B)$ be the set of all finite subsets of B . Then nonempty $A \subset {}^*B$ is hyperfinite if and only if $A \in ({}^*F)(B)$, where F is considered as an operator that generates all of the finite subsets of a set.

Theorem 4.3.3. Any nonempty finite set of internal individuals or entities is internal and hyperfinite.

With respect to Definition 4.3.1 the maps in each ${}^*F([0, p], {}^*B)$ are internal and behave like internal (partial) sequences. The ordinary finite manipulation we do with finite sets of real numbers can be extended to the hyperreals by means of * -transfer and by describing these processes by means of (partial) sequences defined on various $[0, p]$. As far as the NSP-world is concerned hyperfinite sets have all the same set-theoretic first-order properties as the finite sets and we do not usually establish these basic hyperfinite properties each time they are first employed. Thus, not only is the union of finitely many hyperfinite subsets of a set ${}^*B$ a hyperfinite subset of ${}^*B$ but the union of a hyperfinite collection of hyperfinite subsets of ${}^*B$ is a hyperfinite subset of ${}^*B$. *However, from the external or metamathematical point of view most hyperfinite sets are not finite as the next result indicates.*

Theorem 4.3.4. Let $A \in \mathfrak{R}$ and assume that A is infinite. Then there exists a hyperfinite set F such that $F \neq A$, $F \neq {}^*A$ and $A \subset F \subset {}^*A$.

It is precisely the concept of the hyperfinite that leads not only to a clear understanding of the processes that underlie a convergent infinite series, but also leads to the basic notion of the integral and Planck's meaning of the term "series."

Finite summation of elements of $\mathfrak{R}$ can be consider a function Σ defined on $\{\mathfrak{R}^{[0, n]} | n \in \mathbb{N}\}$ with values in $\mathfrak{R}$ and a corresponding function for the finite summation of members of $\mathfrak{R}^n$. It is customary to express the values as $\sum_{i=0}^n a_i$. By * -transfer if $\Gamma \in \mathbb{N}_\infty$, then the **hyperfinite sum** ${}^*\sum_{i=0}^\Gamma a_i \in \mathfrak{R}$. By convention, the symbol ${}^*\sum_{i=0}^\Gamma$ is written as $\sum_{i=0}^\Gamma$. It is a simple matter to translate Definition 3.2.1 for sequential convergence into the following theorem for convergence of a infinite series.

Theorem 4.3.5. An infinite series, $\sum_{i=0}^\infty a_i$, converges to $r \in \mathfrak{R}$ if and only if for each $\Gamma \in \mathbb{N}_\infty$ it follows that $\sum_{i=0}^\Gamma a_i \in \mu(r)$.

My experience indicates that Theorem 4.3.5 is not the most paramount application of the concept of hyperfinite summation for either physical or geometric modeling and, indeed, does not correspond to Planck's description for a series of infinitely small changes. In the next section, examples are discussed that substantially indicate the true character of this concept when applied to geometry or natural system behavior.

4.4. Continuity and a Few Examples.

De l'Hospital's concept of what constitutes a **curve** is not satisfactory for Robinson's theory. Recall that one acceptable analytical definition for the notion of a curve in $\mathbb{R}^n$ is the following: a curve is a continuous map $c: [0, 1] \rightarrow \mathbb{R}^n$. This is equivalent to considering c as determined by n continuous coordinate functions $x_i = f_i(t)$, $1 \leq i \leq n$ each defined on $[0, 1] \subset \mathbb{R}$. Of course, the geometric curve C determined by these functions is usually considered as the set $\{(x_1, \dots, x_n) | t \in [0, 1]\}$. The $*$ -transform of these defining functions leads to the functions $x_i = {}^*f_i(t)$, $1 \leq i \leq n$ each defined on ${}^*[0, 1] \subset {}^*\mathbb{R}$ and they generate the "hypercurve" ${}^*C \subset {}^*\mathbb{R}^n$. The analytic geometry of ${}^*\mathbb{R}^n$ is similar to the customary geometry except that it must be considered non-Archimedean in character. From the viewpoint of the geometry of the NSP-world, if the C is not linear, then ${}^*C$ is not linear and this would entail a necessary rejection de l'Hospital's infinitesimal description. As to the definition of continuity the following captures the envisioned belief that continuous functions preserve the infinitely close.

Definition 4.4.1. (Continuity). For any nonzero $n, m \in \mathbb{N}$ and any nonempty $A \subset \mathbb{R}^n$ a function $f: A \rightarrow \mathbb{R}^m$ is **CONTINUOUS** at $p \in A$ if ${}^*f[\mu(p) \cap {}^*A] \subset \mu(f(p))$. Also, f would be **UNIFORMLY CONTINUOUS** on A if for each $p, q \in {}^*A$ such that $p \approx q$, then ${}^*f(p) \approx {}^*f(q)$.

Observe that uniform continuity seems to preserve the infinitely close in the most satisfactory manner. On the other hand, pointwise continuity is a monad preserving property.

Theorem 4.4.1. *Definition 4.4.1 for continuity and uniform continuity is equivalent to the classical $\delta - \epsilon$ definition.*

One of the most powerful ideas in elementary analysis is that of the compact subset [or subspace if you wish] of the space $\mathbb{R}^n$. Rather than dwell upon the many equivalent standard definitions for this notion a direct nonstandard assault is very enlightening since it reveals immediately the relationship between compactness and continuity as well as an intuitive comprehension of what is being compressed or compacted.

Definition 4.4.2. (Compactness). For any nonzero $n \in \mathbb{N}$ a nonempty set $A \subset \mathbb{R}^n$ is **COMPACT** if ${}^*A \subset \bigcup \{\mu(r) | r \in A\}$.

Theorem 4.4.2. *Definition 4.4.2 for compactness is equivalent to the standard definition utilizing open covers.*

The reason that some 17'th century geometers considered non-linear curves to be collections of infinitesimal line segments was in their desire to use infinitesimal analysis to measure a curves length by corresponding this measure to the polygons of Eudoxus. Even though de l'Hospital's original description is inadequate, a modification does secure the accepted analytical results.

Example 4.4.1.A. Let $\Gamma \in \mathbb{N}_\infty$. Then $F = \{t_i | t_i = i/\Gamma \wedge 0 \leq i \leq \Gamma\}$ is an internal and hyperfinite subset of ${}^*[0, 1]$. By $*$ -transfer, F behaves like an ordered partition of the interval $[0, 1]$ as

defined in the standard sense. Such a set is termed a **fine partition** (i.e. hyperfinitely many members of $[0, 1]$ generating subintervals that are infinitesimal in length). The internal set F generates the internal set of “points” $P = \{(*f_1(t_i), \dots, *f_n(t_i)) | t_i \in F\}$ that are members of the hypercurve $*C$. Now for each $i = 0, \dots, \Gamma - 1$, and each j , $0 \leq j \leq n$ let $*f_j(t_{i+1}) - *f_j(t_i) = d(j, i)$. (If c is continuous, then each $d(j, i) \in \mu(0)$.) For each $i \in *\mathbb{N}$ such that $0 \leq i \leq \Gamma - 1$, the internal set $\ell_i = \{(x_1, \dots, x_n) | \forall j \in *\mathbb{N}, 0 \leq j \leq n, x_j = *f_j(t_i) + t(d(j, i)) \wedge t \in *[0, 1]\}$ is a hyperline segment connecting the two points $(*f_1(t_i), \dots, *f_n(t_i))$, $(*f_1(t_{i+1}), \dots, *f_n(t_{i+1}))$ on the curve $*C$. From this one obtains the internal hyperpolygonal curve $\mathcal{P}_\Gamma = \bigcup \{\ell_i | 0 \leq i \leq \Gamma - 1\}$. As to the length of $\mathcal{P}_\Gamma$ simply extend the concept of length in the classical sense by defining for each $i = 0, \dots, \Gamma - 1$ the vector $\vec{v}_i = (d(1, i), \dots, d(n, i)) \in *\mathbb{R}^n$. Then let the hyperfinite sum $\sum_{i=0}^{\Gamma-1} \|\vec{v}_i\| = |\mathcal{P}_\Gamma| \in *\mathbb{R}$. Even though, in general, you would have a different hyperpolygon with a different hyperreal length for $\forall \Gamma \in \mathbb{N}_\infty$ Robinson [1966, 84–86] showed that if c is continuously differentiable, then for all $\Gamma \in \mathbb{N}_\infty$, $|\mathcal{P}_\Gamma| \in \mu(r)$ and the real number r was the length of the curve obtained in the classical sense by means of the integral.

Example 4.4.1.B. Under the same criterion as stated in example 4.4.1.A that c is continuously differentiable, the length of a curve is actually closer to the limit concept than it is to the de l’Hospital description. This is seen by simply following the same process but replacing Γ with an arbitrary nonzero $n \in \mathbb{N}$. As n increases this yields an increasing sequence $|\mathcal{P}_n|$. Application of Theorem 3.5.1 implies that if there exists but one $\Gamma \in \mathbb{N}_\infty$ such that $|\mathcal{P}_\Gamma| \in \mathcal{O}$, then from example 4.4.1.A this sequence converges to $\text{st}(|\mathcal{P}_\Gamma|)$ and has the same standard part for all $\Gamma \in \mathbb{N}_\infty$.

The process used to obtain the length of a curve in example 4.4.1.A shows that for most ordinary curves there are infinity many hyperpolygons that have the same standard part generated length. Thus each can be used as a NSP-world representative for the curve itself, at least as far as length is concerned. The fact that there does not, in general, exist a unique hyperpolygon is a disadvantage from the viewpoint of the founders of the infinitesimal method and forces a rejection of the de l’Hospital description. However, if in the natural world a curve is envisioned to be a path of motion produced by physical processes, then the lack of uniqueness could be an advantage. Under this interpretation, it would indicate that different and possibly interesting NSP-world ultranatural processes yield the same effect when they are restricted to the natural world.

There has arisen recently a significant application of the seeming esoteric idea that there may exist a multitude of distinct NSP-effects that yield the same natural world effects. Next is an example of how this might occur.

Example 4.4.2. (Fractals) Science has become interested in order and design as reflected in what has become known as “fractal” behavior. For this example, let $c: [0, 1] \rightarrow \mathbb{R}^n$ be a curve; but, assume that the sequence of polygon approximations, $|\mathcal{P}_i|$, discussed in example 4.4.1.B has the property that $|\mathcal{P}_i| \rightarrow +\infty$. This is apparently one of the salient features of a fractal curve. In Herrmann [1989] it is shown that for any nonempty compact $K \subset \mathbb{R}$ and for any continuous $c: K \rightarrow \mathbb{R}^n$ there exists an internal $G: *K \rightarrow *\mathbb{R}^n$ such that G is $*$ -differentiable of any order $m \in *\mathbb{N}$, G has a well-defined hyperreal length and $\text{st}(G) = c$. From the viewpoint of the NSP-world of processes and paths of motion, this G represents the same effects as does the function c except that G has an associated length concept and is ultrasmooth. Moreover, the internal object

G is somewhat less arbitrary in character than are those in example 4.4.1 since it is selectable from a specific algebra of functions. This may be significant since when it has been rigorously shown that certain physical attributes are representable by fractal curves then the theory that models such physical attributes is usually associated with some specific algebra of functions. Thus a standard fractal curve may be replaced by the standard part of an ultrasmooth curve with a well-defined length.

The idea of hyperfinite summation as representing the geometric length of a curve is, of course, closely associated with the elementary integral. Indeed, when we quoted Planck's fundamental description for physical modeling a question was asked, "What might the term 'series' mean in Planck's statement?" Apparently, what Planck meant by this term is the type of hyperfinite summation that, as seen in the next chapter, produces the integral.

Chapter 5.

**STANDARD RULES
FOR INTEGRAL MODELING**

5.1 The Riemann Styled Integral.

The mathematical concept variously termed “integration” (i.e. to bring together the parts or to make whole) was, until after the time of Cauchy, always considered to be a specifically defined summation process. In 1823 Cauchy wrote the following description, where $f: [a, b] \rightarrow \mathbb{R}$ is assumed to be continuous. “...if one divides $X - x_0$ into infinitesimally small elements $x_1 - x_0, x_2 - x_1, \dots, X - x_{n-1}$ the sum

$$S = (x_1 - x_0)f(x_0) + (x_2 - x_1)f(x_1) + \dots + (X - x_{n-1})f(x_{n-1})$$

converges to a limit represented by the definite integral $\int_{x_0}^X f(x) dx$.” From our new nonstandard point of view such a sum is produced by a hyperfinite partition [see Example 4.4.1.A] generated by some $\Gamma \in \mathbb{N}_\infty$ and the internal hyperfinite sequence of values $f(x_i)$, $0 \leq i \leq \Gamma - 1$. Obviously, the Riemann sum, where f may be evaluated at any member of each subinterval $[x_i, x_{i+1}]$, is styled after this Cauchy definition. However, does the standard part of such a hyperfinite sum exist and is it independent of the partition? Moreover, can the concept be extended to bounded not necessarily continuous functions?

In the following very brief discussion, neither the Stieltjes nor Lebesgue generalization is considered, even though these have been extensively investigated by nonstandard means. Indeed, research indicates that the use of arbitrary partitions along with the Darboux concept of the upper and lower sums and the upper and lower integrals which he proved to be equivalent to the Riemann integral are note worthy in that they more easily yield the rigorous proofs that establish the properties of the highly applicable converging Riemann sum notion. Since our paramount concern is modeling with the Riemann styled integral, our stated results are in terms of such easily conceived hyperfinite sums. A simple nonstandard definition for an integral of bounded functions - the H-integral - is given in Herrmann [1985]. In this paper, it is shown that the H-integral is equivalent to the Darboux integral. [Note: in Herrmann [1985] the Darboux integral is called the Riemann integral. Further, Theorem 3.3 and Corollaries 3.3.1, 3.3.2, 3.3.3, 3.3.4 in this paper are incorrect as stated. However, our use of the results from this paper are not related to these few erroneous conclusions.] As is well-known the Darboux integral is equivalent to the Riemann integral conceived of as approximated by the Riemann Sums. It seems expedient, however, to consider all such integral concepts extended to reasonable subsets of $\mathbb{R}^n$ using the simplest possible procedures [Apostol [1957], Spivak [1965]].

For an n -dimensional space $n \geq 1$, the closed set $R = [a_1, b_1] \times \dots \times [a_n, b_n]$, $a_i < b_i$, $1 \leq i \leq n$ is called a **rectangle**. Of course, if $n = 1$, then a “rectangle” is but a closed interval. As usual, consider for each $[a_i, b_i]$ a **partition** P_i as a finite set of members of $[a_i, b_i]$ such that $a_i, b_i \in P_i$ and where P_i is considered as ordered. This is often explicitly written as $P_i = \{x_{i0}, \dots, x_{ik}\}$, $a_i = x_{i0} < x_{i1} < \dots < x_{ik} = b_i$. This determines the closed one-dimensional subintervals $[x_{i(p-1)}, x_{ip}]$, $1 \leq p \leq k$. In brief, this process obtains a partition $P = P_1 \times \dots \times P_n$ of R and a finite collection of closed n -dimensional subrectangles R_q obtained by considering $([x_{10}, x_{11}] \cup \dots \cup [x_{1k-1}, x_{1k}]) \times \dots \times ([x_{n0}, x_{n1}] \cup \dots \cup [x_{nm-1}, x_{nm}])$. Each R_q has a measure, $v(R_q) \in \mathbb{R}$, assigned to it which is intuitively the product of the lengths of the sides. For simplicity of notion the definition of the measure $v(R_q)$ is left intuitively understood.

Probably the simplest partition to consider would be the one termed a **simple** partition. These are formed by selecting n nonzero natural numbers $m_1, \dots, m_n$ and dividing each interval $[a_i, b_i]$ into an equal length partition by adding to each successive partition point the number $(b_i - a_i)/(m_i)$. This concept is extended to the nonstandard world by selecting n infinite natural numbers $\Gamma_1, \dots, \Gamma_n$ and generating for each interval $[a_i, b_i]$ an internal hyperfinite partition, P_i , each subinterval of which has positive infinitesimal length $(b_i - a_i)/(\Gamma_i) = dx_i$. Then the partition $P = P_1 \times \dots \times P_n$ is a **simple fine partition** of R . Such a partition yields an internal set of hyperrectangles R_i such that $*v(R_i) = dx_1 \cdots dx_n = dX \in \mu(0)^+$. You could be much more general and consider the **fine** partitions which are internal collections of hyperfinitely many members of $*[a_i, b_i]$ such that the length of any subinterval is an infinitesimal. In Cauchy's definition he evaluated a function at specific endpoints of each subinterval. For a bounded function $f: R \rightarrow \mathbb{R}$ this evaluation concept can also be applied in the case of the subrectangles into which R is partitioned by evaluating the function f at, say, the corner nearest to the origin. However, it has become customary to be somewhat more general and include the concept of the intermediate partition. Let P be a partition of the rectangle R and assume that P determines the set of subrectangles $\{R_q | 1 \leq q \leq m\}$. An **intermediate partition**, Q , is any finite sequence of vectors $\{\vec{v}_q\}$, where $\vec{v}_q \in R_q$ for each q such that $1 \leq q \leq m$.

Definition 5.1.1. (The Integral). Let $f: R \rightarrow \mathbb{R}$ be bounded and $\mathcal{P}$ the set of simple partitions of R . Then f is said to be **INTEGRABLE** if there exists some $r \in \mathbb{R}$ and a simple fine partition, $P \in *\mathcal{P}$ such that for each of its internal intermediate partitions $\{\vec{v}_q\}$, where $1 \leq q \leq \Gamma \in \mathbb{N}_\infty$,

$$\sum_{k=1}^{\Gamma} *f(\vec{v}_q) *v(R_q) \in \mu(r).$$

Theorem 5.1.1. A bounded function $f: R \rightarrow \mathbb{R}$ is integrable if and only if it is integrable in the sense of Darboux and Riemann Sums.

Theorem 5.1.2. If bounded $f: R \rightarrow \mathbb{R}$ is integrable, then there exists a unique $r \in \mathbb{R}$ such that for every fine partition $P = \{\vec{x}_0, \dots, \vec{x}_\Omega\}$, $\Omega \in \mathbb{N}_\infty$ and every internal intermediate partition $Q = \{\vec{v}_q\}$, $1 \leq q \leq \Gamma \in \mathbb{N}_\infty$ it follows that

$$\sum_{k=1}^{\Gamma} *f(\vec{v}_q) *v(R_q) \in \mu(r).$$

The unique real number that exists by Theorem 5.1.2 is, from theorem 5.1.1, the classical value of the definite integral and hence if bounded $f: R \rightarrow \mathbb{R}$ is integrable in the sense of Definition 5.1.1, then we may write

$$\int \cdots \int_R f(\vec{x}) dx_1 \cdots dx_n = \int_R f(\vec{x}) dX = \text{st} \left(\sum_{k=1}^{\Gamma} *f(\vec{v}_q) *v(R_q) \right).$$

Observe that Theorem 5.1.2 indicates that Cavalieri's notion of the indivisible line segment as being the foundation for the definite integral is untenable. Even though the above hyperfinite sums correlate directly to the intuitive concept of the definite integral, even for possibly discontinuous

functions, this fact alone does not lead to the appropriate selection of specific integrands that will produce meaningful geometric or physical measures. *Indeed, one of the most significant aspects of modern infinitesimal analysis is that there does exist describable modeling procedures that rigorously establish that a specific integrand does provide the requisite value for a specific geometric or physical quantity.*

5.2 The Infinite Sum Theorems.

Throughout applied mathematics numerous linear functionals are utilized to discuss and predict geometric or physical qualities. Such functionals are essential to the indirect verification of many physical theories for within the laboratory environment it is the predicted values displayed by elaborate machinery that often yield the only indications that unobserved events may be occurring. As indicated in section 5.1, there are now rigorous rules that lead to an immediate adoption of the integral as the appropriate modeling structure when one mentally conceives of such events and applies experience to determine the geometric or physical properties that might establish that it is likely that certain hypothesized behavior is actually occurring.

Prior to Robinson's discovery certain vaguely described rules did appear in the mathematical literature; but none was consistently defined in a rigorous language nor did they have any particular relation to the successfully applied intuitive notions we term infinitesimal reasoning. As an intermediate step in establishing a consistent and complete approach to this problem, it is now possible to describe explicitly sufficient infinitesimal conditions that establish the integral as the correct modeling structure. These intermediate rules have become known as *The Infinite Sum Theorems*. For any bounded function $f: [a, b] \rightarrow \mathbb{R}$ a generalizations of these rules can be found in Herrmann [1985]. In appendix 5, this rule is further generalized and applied to the case that bounded $f: R \rightarrow \mathbb{R}$. Obviously, a **subrectangle**, R_S , is a rectangle that is a subset of R . Let $\mathcal{C}$ be the set of all simple partitions of R , $\mathcal{S}(P)$ the set of all subrectangles generated by $P \in \mathcal{C}$ and $\mathcal{C}_{PSR} = \{R_S | \exists P(P \in \mathcal{C} \wedge R_S \in \mathcal{S}(P))\}$ the set of all subrectangles contained in any simple partition of R .

Definition 5.2.1. (Simply Additive). Let $\{R_q | 1 \leq q \leq m\}$ be any simple partition of R and let B be any map defined on the collection $\mathcal{C}_{PSR}$ and having real number values. Then B is said to be **SIMPLY ADDITIVE** if for each $\{R_q | 1 \leq q \leq m\} = \mathcal{S}(P)$, $P \in \mathcal{C}$ it follows that

$$B(R) = \sum_{k=1}^m B(R_k).$$

Obviously, $*B$ is defined on the set $*\mathcal{C}_{PSR}$ which contains all of the infinitesimal subrectangles contained in any simple fine partition of R . I point out that our first infinitesimal sum theorem actually holds for a slightly more general partition than a simple partition of R - the special partition. However, a simple partition is a special partition and in most applied cases the simple partition suffices.

Theorem 5.2.1. (An Infinite Sum Theorem.) Let bounded $f: R \rightarrow \mathbb{R}$ and simply additive $B: \mathcal{C}_{PSR} \rightarrow \mathbb{R}$. If there exists a simple fine partition $\{R_q | 1 \leq q \leq \Gamma\}$ and for each R_q there exists some $\vec{p} \in R_q$ such that

$$*B(R_q)/dX \approx *f(\vec{p}), \quad (*)$$

then f is integrable and

$$B(R) = \int_R f(\vec{x}) dX.$$

Two observations about Theorem 5.2.1. The infinitesimal dX need not be considered the finite product of coordinate measures but may also take on the character of such physical quantities as the finite product of infinitesimal momenta, the finite product of infinitesimal probabilities and even the finite product of infinitesimal charges or infinitesimal numbers of elementary particles if such things can be conceived of in objective reality. Further, it is somewhat unfortunate for applied mathematics that the converse of Theorem 5.2.1 does not hold. N.J. Cutland [1986] has supplied your author with an example of a function defined on $[0, 1]$ that is Darboux integrable (hence integrable) but if you define $B(R_S) = \int_{R_S} f(x) dx$, then ${}^*B$ does not satisfy property (*). On the other hand, if f is continuous on R , then (*) does hold for such integrally defined functionals and, indeed, a much stronger property called *supernearness* holds as well. Let $\mathcal{C}_{SR}$ be the set of all subrectangles contained in R . **For simplicity of notation, throughout this manual, maps such as B are denoted as being, at least, defined on sets such as $\mathcal{C}_{PSR}$, $\mathcal{C}_{SR}$ etc. Simple additivity and additivity will greatly enlarge their domains of definition.**

Definition 5.2.2. (Supernearness).

Let $(x_1, \dots, x_n), (y_1, \dots, y_n) \in {}^*R$ and $y_i - x_i \in \mu(0)^+$, $1 \leq i \leq n$; and let $R_S = \{(z_1, \dots, z_n) | \forall i (1 \leq i \leq n \rightarrow x_i \leq z_i \leq y_i) \wedge (z_i \in {}^*\mathbb{R})\}$ denote an infinitesimal subrectangle of R . A map $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ is **SUPERNEAR** to bounded $f: R \rightarrow \mathbb{R}$ if for every infinitesimal subrectangle R_S of ${}^*R$ and for every $\vec{p} \in R_S$ it follows that

$${}^*B(R_S)/dX \approx {}^*f(\vec{p}), \quad (**)$$

where $dX = \prod_{i=1}^n (y_i - x_i) \in \mu(0)^+$.

Theorem 5.2.2. *A bounded function $f: R \rightarrow \mathbb{R}$ is continuous if and only if there exists a map $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ that is supernear to f .*

In the proof of Theorem 5.2.2 the following interesting integral property is established.

Corollary 5.2.2. *Suppose that $f: R \rightarrow \mathbb{R}$ is continuous. For each $R_S \in \mathcal{C}_{SR}$ define $B(R_S) = \int_{R_S} f(\vec{x}) dX$. Then B is supernear to f .*

Theorem 5.2.3. *Let bounded $f: R \rightarrow \mathbb{R}$. If $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ is supernear to f and simply additive on each simple partition of each R_S , then f is continuous on R and*

$$B(R_S) = \int_{R_S} f(\vec{x}) dX$$

for each $R_S \in \mathcal{C}_{SR}$.

In applications of the integral to geometric and physical problems it is usually assumed that the map $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ is, at least, **additive** on $\mathcal{C}_{SR}$. Recall that this means that if nonempty

$\{R_i \mid 1 \leq i \leq n\} \subset \mathcal{C}_{SR}$ is pairwise disjoint or pairwise has only boundary points in common, then $B(\cup\{R_i\}) = \sum_{i=1}^n B(R_i)$. Note that if B is additive on $\mathcal{C}_{SR}$, then B is simply additive on each member of $\mathcal{C}$ and for each simple partition of $R_S \in \mathcal{C}_{SR}$.

Corollary 5.2.3.1 *Let bounded $f: R \rightarrow \mathbb{R}$. If $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ is supernear to f and additive on $\mathcal{C}_{SR}$, then f is continuous on R and*

$$B(R_S) = \int_{R_S} f(\vec{x}) dX$$

for each $R_S \in \mathcal{C}_{SR}$.

Corollary 5.2.3.2 *Let bounded $f: R \rightarrow \mathbb{R}$. There exists one and only one map $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ that is supernear to f and either simply additive on each simple partition of each R_S or additive on $\mathcal{C}_{SR}$.*

5.3 Extensions.

In general, the bounded real valued function f need not be defined on such a convenient set as R . If $f: D \rightarrow \mathbb{R}$ is defined on a bounded set D , then the most expedient procedure to follow is to define a function $\hat{f}: R \rightarrow \mathbb{R}$, where $D \subset R$, by $\hat{f}(\vec{x}) = f(\vec{x})$ for each $\vec{x} \in D$ and $\hat{f}(\vec{x}) = 0$ for each $\vec{x} \in R - D$. With this case then, as is customary, let $\int_D f(\vec{x}) dX = \int_R \hat{f}(\vec{x}) dX$.

As far as a map such as $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ is concerned, the additivity of B may be extended to all Jordan-measurable subsets of R . The fact that B may be additive on a lesser collection of subsets of R will suffice for the basic modeling rules described in the next section. These modeling rules are very specific in character and if the proper simplistic assumptions for B are utilized, then they lead directly to the appropriate infinite sum theorem and its associated integral equivalence.

$\Rightarrow$ IMPORTANT $\Rightarrow$

$\Rightarrow$ In the following applications, the standard requirements are stated in terms of what we perceive to be global behavior of well-known ordinary functionals and their relation to standard characterizing properties. **Many of these observation are not obvious. The following applications are actually intended to foster an appreciation for the nonstandard modeling rules and procedures that appear in Chapter 6 - rules that lead more directly to the appropriate conclusions.** The reason we present the following applications is that this global approach is used in the more elementary textbooks. $\Leftarrow$

5.4 Applications and the Standard Modeling Rules.

One of the unusual aspects of the Infinite Sum Theorem 5.2.1 is that it does not view the functional B directly but, rather, a “mean value” must be considered if the attention is directed toward the concept of being infinitely close. As the derivations in the following elementary applications indicate this problem is submerged within the derivation itself and does not usually occur when the properties of the basic functional are proposed. Our applications are mostly geometric and elementary in character, while the major applications to the physical sciences will appear in their respect manuals. *Further, we concentrate upon those applications that traditionally appear in the customary core calculus, elementary differential equations and physical science courses.*

Application 5.4.1 *The 2-dimensional area between two continuous curves.*

First assume that all of the following functions are continuous on their indicated domains. Give $h: [c, d] \rightarrow \mathbb{R}$ and $k: [c, d] \rightarrow \mathbb{R}$, where we denote by the symbol $h \leq k$ the condition that $h(x) \leq k(x)$ for each $x \in [c, d]$. Let's look at the original idea behind an area function $A([c, d], h, k)$ that measures the intuitive area between these two curves and over the interval $[c, d]$. Define a **basic region** R for any pair of functions h, k , $h \leq k$ and over any interval $[c, d]$ contained in their common domain by $R = \{(x, y) | x \in [c, d] \wedge h(x) \leq y \leq k(x)\}$. Assume that we have two fixed functions f, g , $f \leq g$ defined on $[a, b]$. The follow rules (axioms) appear to model our intuitive notion of an area function.

- (i) The area function A is, at least, defined on all basic regions determined by $[c, d] \subset [a, b]$.
 - (ii) For the above two fixed functions f, g defined on $[a, b]$ the area function A is, at least, additive on the set $\mathcal{C}_{SR}$ of $[a, b]$.
 - (iii) If D, E are two basic regions in the domain of A and $D \subset E$, then $A(D) \leq A(E)$.
 - (iv) If two functions h, k are constant over any $[c, d] \subset [a, b]$, then $A([c, d], h, k) = (d - c)(k - h)$.
- These four properties for the area function A are certainly reasonable and seem to model the intuitive notions from elementary plane geometry. We now formally establish that

$$A([a, b], f, g) = \int_a^b (g(x) - f(x)) dx. \quad (I_1)$$

Derivation. Let $[x_1, x_1 + dx]$ be any hyperinterval generated by any simple fine partition of $^*[a, b]$. Note in this case $dx = (b - a)/\Gamma$, where $\Gamma \in \mathbb{N}_\infty$. By considering the *-transfer of the standard extreme value theorem for continuous functions defined on closed intervals it follows that $^*f$ and $^*g$ attain their maximum and minimum values $^*f_M$, $^*g_M$ and $^*f_m$, $^*g_m$ respectively on $[x_1, x_1 + dx]$. It is not difficult to model statements (i), (ii), (iii) (iv) set- theoretically and extend these properties to the nonstandard world. We need only consider statements (i), (ii), (iii), (iv) as intuitively *-transformed by changing the terminology to “hyper” or “*” terminology. When this is done statements (i), (iii) and (iv) yield the result that

$$(^*g_m - ^*f_M)dx \leq ^*A([x_1, x_1 + dx], ^*f, ^*g) \leq (^*g_M - ^*f_m)dx \Rightarrow \quad (1)$$

$$(^*g_m - ^*f_M) \leq (^*A([x_1, x_1 + dx], ^*f, ^*g))/dx \leq (^*g_M - ^*f_m). \quad (2)$$

Since f and g are uniformly continuous on $[a, b]$ and $^*f, ^*g$ attain their respective maximum and minimum value at members of $[x_1, x_1 + dx]$ then definition 4.4.1 implies that

$$^*f_m \approx ^*f(x_1) \approx ^*f_M, \quad ^*g_m \approx ^*g(x_1) \approx ^*g_M. \quad (3)$$

Consequently,

$$(^*g_m - ^*f_M) \approx (g(x_1) - f(x_1)) \approx (^*g_M - ^*f_m) \Rightarrow \quad (4)$$

$$(A([x_1, x_1 + dx], ^*f, ^*g))/dx \approx (g(x_1) - f(x_1)) \quad (5)$$

from the fact that f, g are bounded and Corollary 2.2.5.2. Statement (i) allows application of the Infinite Sum Theorem and integral equation I_1 is the consequence.

Once equation I_1 is obtained then it may be checked against the standard area measures for the ordinary Euclidean plane figures in order to insure that it is indeed an extension. I point out that throughout many of these applications similar modeling rules such as (i), (ii), (iii) and (iv) are essential if one wishes to achieve a formal derivation. As will be illustrated there are notable

exceptions to this general approach where one of the standard conditions (i) - (iv) may fail. However, here are two more applications where the standard functional characterizations can be formulated

Application 5.4.2. *Volume obtained by a 2-dimensional integral.*

Assume, as in the previous application, that all functions are continuous on their indicated domains. Suppose that two functions h, k are defined on a rectangle $R_0 \subset \mathbb{R}^2$ and have the property that the $h \leq k$ on R_0 . Generalizing the definition in application 5.4.1, define a basic region R_B in $\mathbb{R}^3$ for each a pair h, k by $R_B = \{(x, y, z) | (x, y) \in R_0 \wedge h(x, y) \leq z \leq k(x, y)\}$. As before, we attempt to model the concept of a volume function $V(R_0, h, k)$ over any rectangle $R_0 \subset \mathbb{R}^2$. Assume that we have two real valued fixed functions f, g , $f \leq g$ defined on a rectangle R .

(i) The volume function V is defined, at least, for all basic regions determined by rectangles that are subsets of R .

(ii) For the above two functions f, g , the function V is, at least, additive on the set $\mathcal{C}_{SR}$ of R .

(iii) If D, E are two basic regions in the domain of V and $D \subset E$, then $V(D) \leq V(E)$.

(iv) If the functions h, k are constant over any rectangle $R_0 \subset R$, then $V(R_0, h, k) = (\text{area } R)(k - h)$.

If V satisfies these rules, then

$$V(R, f, g) = \int_R (g(\vec{x}) - f(\vec{x})) dX.$$

Derivation. Except for a very slight modification, this is exactly the same as the derivation for application 5.4.1. Simply let $[x_1, x_1 + dx_1] \times [x_2, x_2 + dx_2]$ be a hyperrectangle generated by a simple fine partition of ${}^*R$. In this case, $dx_1 = (b - a)/\Gamma$, $dx_2 = (d - c)/\Omega$, $\Gamma, \Omega \in \mathbb{N}_\infty$. Since R is compact the remainder of this derivation is as in application 5.4.1.

Application 5.4.3. *Mass obtained by a 3-dimensional integral.*

Assume that $\rho(\vec{x})$ is a continuous point density function defined on a rectangle $R \subset \mathbb{R}^3$. General physical experience leads to the following characterizations for the elementary mass, $M(R, \rho)$, of R .

(i) The mass is defined for, at least, the set of all subrectangles, $\mathcal{C}_{SR}$, of R .

(ii) The mass is additive on the set $\mathcal{C}_{SR}$.

(iii) If R_S is a subrectangle of R and $\rho_1(\vec{x})$ is a continuous density function defined on R_S with the property that $\rho_1(\vec{x}) \leq \rho(\vec{x})$ for each $x \in R_S$, then $M(R_S, \rho_1) \leq M(R_S, \rho)$.

(iv) If the function ρ is constant over any rectangle $R_0 \subset R$, then $M(R_0, \rho) = \rho(v(R_0))$.

If M satisfies rules (i) - (iv), then

$$M(R, \rho) = \int_R \rho(\vec{x}) dX.$$

Derivation. As in the previous cases, one selects a simple fine partition of ${}^*R$ and lets R_q be some hyperrectangle determined by such a simple fine partition. The above characterizations are extended by * -transfer to the NSP-world. From continuity, ${}^*\rho$ attains its minimum value ${}^*\rho_m$ and maximum value ${}^*\rho_M$ at members of R_q . From (i), (iii) and (iv) it follows that

$$({}^*\rho_m)dX \leq {}^*M(R_q, {}^*\rho) \leq ({}^*\rho_M)dX \Rightarrow \quad (1)$$

$${}^*\rho_m \leq ({}^*M(R_q, {}^*\rho))/dX \leq {}^*\rho_M. \quad (2)$$

Let x_1 be any member of R_q . From the uniform continuity of ρ it follows that

$${}^*\rho_m \approx {}^*\rho(x_1) \approx {}^*\rho_M. \quad (3)$$

Consequently,

$$(*M(R_q, *\rho))/dX \approx *\rho(x_1) \quad (4)$$

and the derivation follows from the Infinite Sum Theorem.

In the above applications, the rules (i) – (v) obviously depict these functionals from a standard point of view as described completely in terms of the standard world. Except within the formal derivations, there may appear to be no consuming need for any insight into the infinitesimal NSP-world. As is be amply illustrated in the next chapter, one or more of these rules may not be self-evident when the integral is applied for both geometric and physical modeling. When this occurs, then it is often the case that certain simplistic and local aspects of the standard world are axiomatically impressed upon the pure NSP-world. This then leads to rules that do include descriptions for pure NSP-world behavior.

5.5 Extensions of the Standard Rules.

In practice, the rectangle R is too restrictive to be of much significance in applications. Fortunately, there are techniques that will allow us to remove this restriction - techniques that lead to a straightforward modification of a few of the terms that appear in such standard rules as (i) – (iv).

The appropriate alteration of these rules begins with the concepts briefly mentioned in section 5.3.

(1) Assume that $f: D \rightarrow \mathbb{R}$ is continuous on D , where D is a compact Jordan-measurable subset of $\mathbb{R}^n$. [Apostal [1957], De Lillo [1982]] The Jordan-measurable subsets of R include those that commonly appear throughout basic applications.

(2) Let $\mathcal{C}_{JR}$ denote the set of all Jordan-measurable subsets of R . Clearly, $\mathcal{C}_{SR} \subset \mathcal{C}_{JR}$ as is well-known.

(3) Assume that $B: \mathcal{C}_{JR} \rightarrow \mathbb{R}$.

(4) Now extend f to $\hat{f}$ and assume that $\hat{f}$ is integrable on R . It is easy to show that the value of $\int_D f(\vec{x}) dX = \int_R \hat{f}(\vec{x}) dX$ is independent of the choice of R .

Our next task is to see how a simple modification of the standard rules leads to a refined derivation that establishes the same integral expression. **In the following applications, the rule modifications are written in italics and $v(J)$ denotes the real Jordan content for any Jordan-measurable set $J \subset R$.**

Application 5.5.1. *Volume obtained by a 2-dimensional integral.*

Assume, as in application 5.4.2, that all functions are integrable on their indicated *compact Jordan-measurable* domains. Suppose that two functions h, k are defined on a *compact and Jordan-measurable* $J_0 \subset R \subset \mathbb{R}^2$ and have the property that the $h \leq k$ on J_0 . Generalizing the definition in application 5.4.1, define a basic region J_B in $\mathbb{R}^3$ for each a pair h, k by $J_B = \{(x, y, z) | (x, y) \in J_0 \wedge h(x, y) \leq z \leq k(x, y)\}$. As before, we attempt to model the concept of a volume function $V(J_0, h, k)$ over any $J_0 \in \mathcal{C}_{JR}$. Assume that we have two real valued fixed and continuous functions f, g , $f \leq g$ defined on *compact and Jordan-measurable* $J \subset R$.

(i) The volume function V is defined, at least, for all basic regions determined by all *compact and Jordan-measurable sets* that are subsets of R .

- (ii) For the *integrable extensions* $\hat{f}, \hat{g}$, of the above two functions the function V is, at least, additive on the set $\mathcal{C}_{JR}$ of R .
- (iii) If D, E are two basic regions in the domain of V and $D \subset E$, then $V(D) \leq V(E)$.
- (iv) If the functions h, k are constant over any $J_0 \in \mathcal{C}_{JR}$, then $V(J_0, h, k) = (v(J_0))(k - h)$. If V satisfies these rules, then

$$V(J, f, g) = \int_R (\hat{g}(\vec{x}) - \hat{f}(\vec{x})) dX = \int_J (g(\vec{x}) - f(\vec{x})) dX.$$

Derivation. Let $I = [x_1, x_1 + dx_1] \times [x_2, x_2 + dx_2]$ be a hyperrectangle generated by a simple fine partition of ${}^*R$. In this case, $dx_1 = (b - a)/\Gamma$, $dx_2 = (d - c)/\Omega$, $\Gamma, \Omega \in \mathbb{N}_\infty$. By * -transfer, we transfer the general results concerning Jordan-measurable sets to the NSP-world. Assume that ${}^*J \cap I = K \neq \emptyset$. Since $I \in {}^*\mathcal{C}_{JR}$ then $K \in \mathcal{C}_{JR}$. Further since I is * -compact then K is * -compact. Noting that ${}^*f = {}^*\hat{f}$, ${}^*g = {}^*\hat{g}$ on K , then the * -extreme value theorem implies that ${}^*\hat{f}$, ${}^*\hat{g}$ attain their maximum and minimum values ${}^*\hat{f}_M$, ${}^*\hat{g}_M$ and ${}^*\hat{f}_m$, ${}^*\hat{g}_m$ respectively on K . Now by application of * -additivity and (iv) and the fact that $K, I, I - K$ are * -Jordan-measurable (if $I - K = \emptyset$, then we still let it be measurable with content equal to zero) we have that ${}^*V(I, {}^*\hat{f}, {}^*\hat{g}) = {}^*V(K, {}^*\hat{f}, {}^*\hat{g}) + {}^*V(I - K, {}^*\hat{f}, {}^*\hat{g}) = {}^*V(K, {}^*\hat{f}, {}^*\hat{g})$.

Next we also apply (iii) and obtain

$$\begin{aligned} ({}^*\hat{g}_m - {}^*\hat{f}_M) {}^*v(K) &\leq ({}^*\hat{g}_m - {}^*\hat{f}_M) dX \leq {}^*V(I, {}^*\hat{f}, {}^*\hat{g}) = \\ {}^*V(K, {}^*\hat{f}, {}^*\hat{g}) &\leq ({}^*\hat{g}_M - {}^*\hat{f}_m) {}^*v(K) \leq ({}^*\hat{g}_M - {}^*\hat{f}_m) dX. \end{aligned} \quad (1)$$

Since $f = \hat{f}$, $g = \hat{g}$ are uniformly continuous on K and $K \neq \emptyset$ then for $x_1 \in K$ we have that ${}^*\hat{f}_m \approx {}^*\hat{f}(x_1) \approx {}^*\hat{f}_M$, ${}^*\hat{g}_m \approx {}^*\hat{g}(x_1) \approx {}^*\hat{g}_M$. Consequently,

$$({}^*\hat{g}_m - {}^*\hat{f}_M) \approx ({}^*\hat{g}(x_1) - {}^*\hat{f}(x_1)) \approx ({}^*\hat{g}_M - {}^*\hat{f}_m). \quad (2)$$

Application of expressions (1) and (2) yields

$${}^*V(I, {}^*\hat{f}, {}^*\hat{g})/dX \approx ({}^*\hat{g}(x_1) - {}^*\hat{f}(x_1)). \quad (3)$$

For the case that $K = \emptyset$ it is obvious that ${}^*V(I, {}^*\hat{f}, {}^*\hat{g}) = 0 = ({}^*\hat{g}(x_1) - {}^*\hat{f}(x_1))dX$ $x_1 \in I$. In this case we also have that expression (3) holds. Application of the Infinite Sum Theorem completes the derivation.

The above example suffices to show how all of the previous standard modeling rules and applications can be extended to the case of the Jordan-measurable subsets and integrable functions. However, for many applications of integral modeling to geometric theories and natural system behavior such listed axioms for the behavior of the conjectured functionals are often not evident. This is particularly so for standard axioms such as (iii) and (iv). To eradicate this difficulty, a direct appeal is made to the NSP-world, either to the infinitesimal terms of the hyperfinite sum that appears in definition 5.1.1 or to statement (*) of Theorem 5.2.1.

Chapter 6.

NONSTANDARD RULES FOR INTEGRAL MODELING

6.1 Historical Examples.

In 1855, Maxwell [1890] presented his fluid flow analogue model for Faraday's concept of both magnetic and electric lines of force. An analysis of Maxwell's imagery relative to our present understanding of the behavior of infinitesimal quantities is very enlightening. Maxwell considers "tubes" of moving points (not particles) of fluid and their paths of motion as a pure imaginary picture of what one might conceive of as line a force. Obviously, the idea was not to consider the concept of "force" as an independent entity but, rather, to first picture "something" - the points of fluid material and their paths of motion - as representing the effects of unknown forces. *"The direction of motion of the fluid will in general be different at different points of the space which it occupies, but since the direction is determined for every such point, we may conceive a line to begin at any point and to continue so that every element of the line indicates by its direction the direction of motion at that point in space. Lines drawn in such a manner that their direction always indicates the direction of fluid motion are called lines of fluid motion."* [Maxwell, 1890:160]

Maxwell then imagines a closed curve on a surface - a surface that "cuts" the lines of fluid motion - and the lines of fluid motion that intersect this surface curve. These curve generated flow lines then produce, in his mind, a tubular surface which he calls *a tube of fluid motion*. He then fills the interior of these tubular surface with the flow lines that intersect that portion of the surface which would have the curve as its boundary. He also assumes that the fluid is incompressible. This yields another postulated property of these flow lines. *"The quantity of fluid which in a unit of time crosses any fixed section of the tube is the same at whatever part of the tube the section is taken....and no part runs through the sides of the tube, therefore the quantity which escapes from the second section is equal to that which enters through the first."* [Maxwell 1890:161]

Maxwell then supplies a paramount nonstandard rule to the methods of infinitesimal model - a rule that has recently been called the concept of the infinitesimal microscope. *"An infinite number of lines would have to be drawn at indefinitely small intervals; but since the description of such a system of lines would involve continual reference to the theory of limits, it has been thought better to suppose the lines drawn at intervals depending on the assumed unit, and afterwards to assume the unit as small of we please by taking a small submultiple of the standard unit."* [Maxwell 1890:161] Notice that Maxwell's statement about the necessity of limit theory is now known to be false. He may have made such a statement, as did Kepler before him, to placate those who might be more attuned to rigorous derivations. Further, following general scientific practices, Maxwell does not establish his limit theory conclusions but, rather, ascribes to an infinitesimalizing approach he claims is equivalent to the physical limit theory. More importantly we have our first vague nonstandard rule for infinitesimal modeling

VR1.

These infinitesimalizing ideas are equivalent to a type of infinite magnification of a infinitely small portion of the fluid - a magnification that yields finitely many lines in our field of view that appear to be drawn at real finite distances apart.

I mention that such geometric notions as expressed in VR1 can indeed be formalized within the geometric theory of ${}^*\mathbb{R}^3$.

Within Maxwell's research reports he states numerous times that the reasons for his derivations are "evident." Thus, he often gives no specific causes for his logical conclusions and leaves them axiomatic in character. It is, however, the vague methods of infinitesimalizing that continue to interests us - methods that are also often assumed to be "evident" to Maxwell's audience. Intuitively, as it will be illustrated, physical and geometric infinitesimal integral modeling also displays yet another vague nonstandard rule.

VR2.

Infinitesimal integral modeling often makes a direct appeal to the Infinite Sum Theorems as well as a simplified interpretation of the equation (Theorem 5.1.2)

$$\int \cdots \int_R f(\vec{x}) dx_1 \cdots dx_n = \int_R f(\vec{x}) dX = \text{st} \left(\sum_{k=1}^{\Gamma} {}^*f(\vec{v}_q) {}^*v(R_q) \right).$$

In the infinite sum theorems the equivalence relation $\approx$ is replaced by an equality while the standard part operator is ignored and the integral is made equal to some type of summation process which is often conceived of as finite in character.

How does Maxwell apply his fluid motion analogy to assumed continuously varying magnetic properties? Surprising, he views them as discrete and constant with relation to his magnified portion the of fluid material. *"The quantity of magnetism in any section of a magnetic body is measured by the number of lines of magnetic force that pass through it."* Maxwell [1890:182] This "number" is assumed to be a standard natural number. *"If i be the quantity of the magnetization at any point, or the number of lines of force passing through unit of area in the section of the solenoid, then the total quantity of magnetization in the circuit is the number of lines which pass through a section, $I = \sum i dy dz$, where $dy dz$ is the element of the section, and the summation is performed over the whole section."* [Maxwell 1890:183] In the magnified view, the section is to be conceived of as a rectangle with actual real number area that is then made arbitrary small by Maxwell's small unit convention. The constant numbers i are not assumed to be altered as a physical quantity by the small unit convention but are fixed constants. The cardinality (i.e. intuitively the number of terms in the summation) is ignored and this "summation" is equated to the integral value I .

Maxwell appears to have arrived at his conclusions by considering the simplest of laboratory experiences for assumed constant quantities and geometric configurations. He has then impressed these experiences upon the infinitesimal nonstandard world. By assuming that the outcome is somehow additive in character this leads directly to the integral model.

With respect to pure geometric nonstandard modeling the same general process, with certain exceptions, is also applied. It was seen in Example 4.4.1.A on page 33 that the length of an n -dimensional curve is viewed globally as approximated by a hyperpolygonal curve and following this the length of the curve is defined to be the hyperfinite sum $\sum_{i=0}^{\Gamma-1} \|\vec{v}_i\| = |\mathcal{P}_\Gamma| \in {}^*\mathbb{R}$. However, for curves that are continuously differentiable Robinson formally showed that each term of this

summation could be replaced by the term

$$\sqrt{\sum_{j=1}^n *f'_j(t_i)^2} dt$$

where this replacement term can be conceived of as the infinitesimal length of an infinitesimal line segment infinitely close (in the sense of the infinitesimal sum theorem) to the original line segments that comprise the hyperpolygonal curve. **A proof of this can be found in appendix 6.** On the other hand, one could proceed backwards, as some geometers have done, and discuss the vague notion of the “element of length,” ds , while forcing the length of a curve to be the “sum” of such elements ds . Indeed, one of the greatest of all infinitesimal geometers does all of his analysis in terms of the still vague “element” concept and uses, what would be today, totally unacceptable non-rigorous derivations. *“Let us now examine the integral curvature of this triangle, which is equal to $\int k d\sigma$, $d\sigma$ denoting a surface element of the triangle. Wherefore, since this element is expressed by $mdp \cdot dq$, we must extend the integral $\iint mdp \cdot dq$ over the whole surface of the triangle.”*[Gauss 1827; Art 20] Gauss in his papers also states that various quantities are equal when in reality they are but infinitely close. Fortunately, the algebraic manipulations of infinitesimal quantities as employed by Gauss and Maxwell were restricted to their ring properties. Unfortunately, Gauss’ derivations are highly non-rigorous in character.

By comparing Robinson’s techniques with many of significant mathematical models developed over the past few centuries by application of intuitive infinitesimal modeling, it is now possible to gain a little better insight into the methods used, to make these methods somewhat more acceptable in rigor and to improve considerably upon the notions outlined in the vague nonstandard rules VR1 and VR2.

6.2 The Monadic Environment.

Vague rule 1 can now be made essentially rigorous in character. When we model the most elementary geometric or simplistic physical behavior it is usually conceived of within bounded portions of $\mathbb{R}^n$. Even though one may later remove the boundedness concept, such modeling often begins with such a restriction. Furthermore, this holds true whether or not one is concerned with real or complex variables. What is conceived of as “elementary geometric” or “simplistic physical” behavior is most often fostered by individual experiences and a personal development of discipline intuition. Simplistic diagrams and sketching develop an intuition for geometric concepts, while basic laboratory experimentation tends to yield to the conscientious investigator an intuitive understanding of basic natural system behavior. It is this phenomenological approach to simplistic physical behavior that has led to the development of most of our present day intricate mathematical models that, even though they may tend to predict observed behavior, may not correspond in their entirety to physical reality.

IR1

Elementary geometric or simplistic physical behavior takes place within an m -dimensional monad, $\mu(\vec{p}) \subset * \mathbb{R}^m$, where standard $\vec{p} \in \mathbb{R}^m$. Such behavior may be intuitively $*$ -transferred to similar behavior within $\mu(\vec{p})$.

It might be argued that IR1 is too restrictive for many of our modern applied mathematical structures. It is interesting to note that many monadic properties generalize not only to general topological spaces (see many of papers published by Robert A. Herrmann from 1975 - 1984) but even to the more general pre and pseudo topological spaces [Herrmann 1980]. Can we identify for ${}^*\mathbb{R}^m$ what, at least partially, constitutes basic simplistic behavior? The first step in this process is to study in $\mathbb{R}^n$ or the laboratory geometric or physical behavior restricted to entities termed “elements.” Further, one is often only concerned with specific functionals associated with such objects. As previous mentioned, in applied infinitesimal analysis, infinitesimals that are denoted by such symbols as dx need not correspond to geometric concept of length. Indeed, they can be negative or correspond to such notions as velocity, acceleration or even infinitesimal “numbers of molecules.” Nevertheless, it is convenient to regard the basic elements as geometric in character and not to be concerned with their specific functional or physical interpretation.

IR2

Let internal $L: {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}^m$ be a designated linear transformation and $S \subset {}^*\mathbb{R}^n$ an infinitesimal subrectangle of ${}^*R$, where rectangle $R \subset \mathbb{R}^n$. **An m-dimensional element is the configuration $L[S]$.** The **basic element** is the infinitesimal subrectangle itself.

For all such designated L considered in IR2 the elements can be characterized as m-dimensional parallelepipeds, where the 1-dimensional parallelepiped is to be conceived of as a line segment. Further, we have the following little theorem that shows the relationship between infinitesimal subrectangles and monads.

Theorem 6.2.1. *Let A be a compact subset of $\mathbb{R}^n$ and the infinitesimal subrectangle $R \subset {}^*A$. Then there exists some $p \in A$ such that $R \subset \mu(p)$.*

⇓ IMPORTANT ⇓

⇒ The hypotheses of many of the following rules and theorems are stated in terms of a Jordan-measurable set J . Such premises may be weakened by restricting the functional to $\mathcal{C}_{SR}$ and assuming that J is but a subrectangle. **I mention the important but well-known fact that all of the usual geometric configurations utilized in the basic calculus and differential equation undergraduate courses are Jordan-measurable. Further, the elementary prototype used in place of the general Jordan-measurable set is the subrectangle itself. Most experimental investigations do not go beyond the subrectangle. After the hypotheses that appear in the following rules and theorems are restricted to subrectangles and their conclusions are verified, then their consequences are often extended to J without further confirmation.**

⇑

In order to use the infinite sum theorem directly for the **basic elementary integral** it is assumed that the internal linear map L is the extended identity may ${}^*I$. Then the following are

gleaned from the basic definitions or the intuitive methods of the geometer as well as from simple experiments on natural system behavior whether it be objectively real or imaginary.

IR3 – Infinitesimal Max. and Min. Rule

(1) We wish to measure a quantity M for a compact Jordan-measurable set $J \subset R \subset \mathbb{R}^n$, where M is defined on and, at least, additive over members of the set $\{\mathcal{C}_{SR}, R - J, J\}$. Further, if subrectangle $S \subset R - J$, $M(S) = 0$, and $M(R - J) = 0$. Let $v(J)$ denote the Jordan content.

(2) There is a generating function $f(\vec{x})$ that is related to the functional M in the following manner:

- (i) The functions f is continuous on J .
 - (ii) Let P be some simple fine partition, $S \in {}^*\mathcal{S}(P)$ and $K = {}^*J \cap S \neq \emptyset$. Then there exist $\vec{x}_m \in K$ and $\vec{x}_M \in K$ such that ${}^*f_m = {}^*f(\vec{x}_m) = {}^*\inf\{{}^*f(\vec{x})|\vec{x} \in K\} = \inf\{{}^*f(\vec{x})|\vec{x} \in K\}$ and ${}^*f_M = {}^*f(\vec{x}_M) = {}^*\sup\{{}^*f(\vec{x})|\vec{x} \in K\} = \sup\{{}^*f(\vec{x})|\vec{x} \in K\}$ and
 - (iii) $({}^*f_m) {}^*v(S) \leq {}^*M(S) \leq ({}^*f_M) {}^*v(S)$. [Note: this is the case where ${}^*L$ is the identity map.]
-
-

Obviously the rules in IR3 are closely related to those that appear in chapter 5, section 5.4. Except for 2(iii) these rules are usually tacitly assumed. It is useful to repeat the observation made in section 5.4. *Certain aspects of rule IR3 could be relaxed if statement 2(iii) is formulated in terms of an infinitesimal “mean value” property for the functional M . However, it is the customary linear form in 2(iii) that appears throughout all of the traditional definitions - a form that we are trained to seek and experimentally justify.* Notice that the terms of a Riemann sum are represented in the inequality in 2(iii) by either ${}^*f_m$ or ${}^*f_M$. The following theorem is almost obvious.

Theorem 6.2.2. *If IR3 holds, then*

$$M(J) = \int_J f(\vec{x}) dX.$$

Theorem 6.2.3. *Let compact Jordan-measurable $J \subset R \subset \mathbb{R}^n$. If continuous $f: J \rightarrow \mathbb{R}$, then for any partition P of R and any $S \in \mathcal{S}(P)$, where $K = J \cap S \neq \emptyset$ there exist $\vec{x}_m \in K$ and $\vec{x}_M \in K$ such that $f_m = f(\vec{x}_m) = \inf\{f(\vec{x})|\vec{x} \in K\}$ and $f_M = f(\vec{x}_M) = \sup\{f(\vec{x})|\vec{x} \in K\}$.*

The title of this chapter is actually somewhat misleading. Even though all of the previous rules relative to the monadic environment are stated in terms of this monadic world, generally for the integral, individuals still rely upon standard world observations. It is the generation of differential equation models that utilize this infinitesimal locale exclusively. What mental or experimental procedures does a researcher employ in order to arrive at the conclusion expressed in IR3 part 2(iii)? **Evidently, these standard procedures must be closely related to the rule IR1 - IR3 even if they are expressed in the language of a standard mathematical structure.** In derivations written prior to 1981, we read that such conclusions are “self-evident” or “obvious.” These self-evident features of informal infinitesimal modeling are, of course, some type of unmentioned infinitesimal reasoning process. Formally, the unmentioned infinitesimalizing procedure can now be characterized as formal * -transfer and what needs to be determined in the large scale or

macroscopic environment prior to such infinitesimalizing is contained in the premises of the next somewhat obvious proposition. We call a set, $\mathcal{P}$, of simple partitions of R **acceptable** if there exists a simple fine partition $P \in {}^*\mathcal{P}$.

Theorem 6.2.4. (Self-evident Max. and Min.) *Let the rectangle $R \subset \mathbb{R}^n$ and suppose that compact Jordan-measurable $J \subset R$. Let M be defined as in (1) of IR3, continuous $f: J \rightarrow \mathbb{R}$, $\mathcal{P}$ an acceptable set of partitions of R and any $P \in \mathcal{P}$. If for any $S \in \mathcal{S}(P)$ such that $J \cap S \neq \emptyset$ it follows that $(f_m)v(S) \leq M(S) \leq (f_M)v(S)$, then the infinitesimalizing process IR3 holds.*

In order to better apply the self-evident theorem an intuitive discussion of its content is in order. The values f_m and f_M are values for the original defining function restricted to $J \cap S$. Thus essentially for the boundary type subrectangles S (i.e. $S \not\subset J$ but $S \cap J \neq \emptyset$) these values have been extended to the entire rectangle S . What has been done should be viewed as an application of these bounding values of f to the extended configurations $\bigcup\{S | S \in \mathcal{S}(P) \wedge J \cap S \neq \emptyset\}$. **[Important. See the simplest rules IR5, IR6 in Appendix 6 where a continuity concept eliminates this difficulty.]**

6.3 Simple Applications.

Assuming in the macroscopic world a continuous distribution of matter throughout a space region Synge and Griffith define the moment of inertia as $I = \int r^2 dm$ and then state that “ dm is the mass of an infinitesimal element...” [Synge and Griffith [1959:173]]. For a uniform rod of total mass m and length $2a$ positioned on the x -axis, they state that $dm = (m dx)/(2a)$. It is often the case that this concept is extended to the case the mass is not uniform but is rather determined by the continuous point density function $\rho(x)$. In this particular case it is claimed that $dm = \rho(x) dx$. However, from the corrected notions of infinitesimal modeling this last statement is simply incorrect. Based upon Corollaries 5.2.3.1 and 5.2.3.2 and the definition of supernearness we can only be assured that for an infinitesimal subrectangle R and for any $x \in R$, $dm = {}^*M(R, {}^*\rho)$ is infinitely close to ${}^*\rho(x) dx$. Indeed, the notion of “infinitely close” in this context is not sufficient for a proper understanding of the relationship between dm and ${}^*\rho(x) dx$. As well be discussed later these two objects must, for a given dx , be “closer” then indicated by the general infinitely close concept. It is, therefore, necessary to establish various elementary applications of these infinitesimal rules in a manner distinctly different from the customary ones. I envision that many of our present day texts that claim to teach the infinitesimal modeling of natural system behavior will need to be completely revised if rigor is to be incorporated.

Application 6.3.1. *The quantity of an incompressible fluid passing through a 2-dimensional rectangle [resp. Jordan-measurable plane region J].*

Suppose that we have for a macroscopic environment a function $V(x, y)$ that represents the point velocity, in a normal direction, of the incompressible fluid passing through a 2-dimensional rectangle R . Then the amount of fluid passing through R per unit time, $Q(R)$, is

$$\int_R v(x, y) dX.$$

Derivation. Before we start this derivation observe that it must be considered slightly less rigorous than those that appear in chapter 5. What does experience indicates about such a value $Q(R)$? First, since the fluid is incompressible then Q is additive on $\mathcal{C}_{SR}$. Observation also indicates

that for a subrectangle R_S of any simple partition $(V_m) v(R_S) \leq Q(R_S) \leq (V_M) v(R_S)$. If you agree to this experiential argument, then by Theorems 6.2.2 and 6.2.3 the result follows.

Application 6.3.2. *Moment of inertia of a circular disc of radius a about a line ℓ through the center perpendicular to the plane of the disc.*

[Point mass method.] Consider the disc J centered at the origin of our 2-dimensional Cartesian coordinate system. Let $\rho(x, y)$ represent a continuous point density function for J . The moment of inertia, I_1 , for a finite system of point masses, m_i , located on the disc each at a distance of r_i from ℓ is $I_1 = \sum r_i^2 m_i$. This may be rewritten as $I_1 = \sum (x_i^2 + y_i^2) m_i$. The first requirement for application of IR3 is that this idea be extended to a continuous density function, $\rho(x, y)$, of point masses. This implies that we consider the continuous moment of inertia generating function $f(x, y) = \rho(x, y) (x^2 + y^2)$. Using this assumed function it follows that

$$Q(J) = \int_J f(x, y) dX.$$

If ρ is a constant, then

$$Q(J) = \int_J \rho(x, y) (x^2 + y^2) dX = \int_0^a 2\pi\rho r^3 dr = (\pi\rho a^4)/2 = (m/2)a^2.$$

Derivation. It's clear that we have selected the basic aspects of this application to force it to have the properties expressed in IR3. However, what experiences do we have with moments of inertia that will allow us to conclude that the inequality $(f_m) v(S) \leq I(S) \leq (f_M) v(S)$ holds? If you have such experiences, then the result is immediate. If you don't or you cannot adequately explain you intuition, then another derivation method would be required.

Application 6.3.2 and the like seem to have a very weak derivations. There are two notations that will aid in eradicating these derivation difficulties. Indeed Maxwell explicitly states one of these procedures and there is a statement within the premises of application 6.3.2 that is significant and often appears when extensions are considered. **This leads to two often used, significant and powerful infinitesimal modeling procedures. The first is the extension of constants values and the second for, point definable quantities, is the extension of the finite to the hyperfinite.**

6.4. The Method of Constants.

The method of constants makes a direct appeal to expression (*) of the Infinite Sum Theorem (5.2.1) as it appears on page 37. Individuals observe simple properties about a functional relative to the assumption that the generating function can have constant values. These observations and a simplified physical theory are then transferred to the NSP-world. They then assume that these simple properties hold for the NSP- world and then without justification state that for a continuous generating function the value of the functional is but a "sum " of the appropriate (infinitesimal) quantities. A rigorous procedure is now possible.

IR4- Method of Constants

(1) In what follows, let for any $A \subset \mathbb{R}^n$ "int" denote the interior of A . Let $\mathcal{A} = \{C_{SR}, \{\text{int}(J \cap S) \neq \emptyset \mid S \in C_{SR}\}\}$. We wish to measure a quantity M for a Jordan-measurable set $J \subset R \subset \mathbb{R}^n$, where M is, at least, defined on and additive over the members of the set $\{\mathcal{A}, R - J, J\}$ and for a

subrectangle S , $S \subset R - \text{int}(J)$ it follows that $M(S) = 0$ and $M(R - J) = 0$. Let $v(J)$ denote the Jordan content.

(2) There is a generating function $f(\vec{x})$ that is related to the functional M in the following manner:

- (i) The function f is bounded on J .
 - (ii) Let P be any arbitrary simple fine partition, $S \in {}^*S(P)$ an arbitrary infinitesimal subrectangle and $\emptyset \neq K = {}^*\text{int}({}^*J \cap S)$.
 - (iii) There exists some $\vec{x} \in K$ such that ${}^*M(S) = {}^*f(\vec{x}) {}^*v(S)$ or ${}^*M(S)/{}^*v(S) \approx {}^*f(\vec{x})$.
-
-

Please note carefully where IR4 differs from IR3. First, f is only assumed to be bounded and J need not be compact. Also note that for $\emptyset \neq K$, ${}^*v(K) \neq 0$ since in the standard case nonempty and Jordan-measurable $\text{int}(J \cap S)$ contains a rectangle.

Theorem 6.4.1. *If IR4 holds, then*

$$M(J) = \int_J f(\vec{x}) dX.$$

Once again we need some sort of infinitesimal reasoning process that leads to application of IR4. Within the laboratory or an imaginary mind experiment individuals often tacitly observe that the hypotheses of the next “self-evident” theorem hold and, without stating it, take an intuitive * -transform and obtain IR4. It is IR4, or something akin to it, that appears in the literature. However, certain aspects of the next result are necessary due to technical difficulties and one useful requirement is some what unexpected.

Theorem 6.4.2. (Self-evident Method of Constants) *Let the rectangle $R \subset \mathbb{R}^n$ and suppose that Jordan-measurable $J \subset R$. Let M be defined as in (1) of IR4, continuous $f: R \rightarrow \mathbb{R}$, $\mathcal{P}$ an acceptable set of partitions of R and any $P \in \mathcal{P}$. If for any $S \in \mathcal{S}(P)$ such that $\emptyset \neq K = \text{int}(J \cap S)$ there exists some $\vec{x} \in K$ and some $\vec{y} \in S$ such that (i) $M(K) = f(\vec{x}) v(K)$ and (ii) $M(S) = f(\vec{y}) v(S)$, then the infinitesimalizing process IR4 holds for f restricted to J .*

As with IR3 the intuitive notion behind infinitesimal reasoning Theorem 6.4.2 is the extension of the generating function values to the boundary type subrectangles S . Both of our self-evident theorems display an interesting phenomenon. The rules IR3 and IR4 require the existence of but one simple fine partition for application; but, the self-evident theorems need an infinite collection of such partitions with the requisite properties in order to guarantee that such a partition exists in the NSP-world. In practice the requirement that f be continuous on R may be relaxed to piecewise continuity on a set of J covering subrectangles of R that at most overlap only on their boundaries. **Of interest is the necessity for premise (ii). In applications it is simply assumed from observation that there is a partition of subrectangles, S , that are “small enough” in size that there exists such a constant $f(\vec{x})$, $\vec{x} \in \text{int}(J \cap S)$ such that (i) holds. The same methodology should also indicate that there exists a constant $f(\vec{y})$, $\vec{y} \in S$ such that (ii) holds. However, the (ii) premise is never mentioned as a requirement.**

Application 6.4.1. (Application 6.3.1 revisited.) *The quantity of an incompressible fluid passing through a 2-dimensional rectangle [resp. Jordan-measurable plane region J]*

Suppose that we have for a macroscopic environment a continuous function $V(x, y)$ that represents the point velocity, in a normal direction, of the incompressible fluid passing through a 2-dimensional rectangle R . Let J be any Jordan-measurable subset of R . Then the amount of fluid passing through J per unit time, $Q(J)$, is

$$\int_J v(x, y) dX.$$

Derivation. It appears that the method of constants yields a more satisfactory derivation of this integral formula. Consider any simple partition P and any $S \in \mathcal{S}(P)$. Let $\emptyset \neq K = \text{int}(J \cap S)$. Then experience indicates that there is some $\vec{x} \in K$ such that $Q(K) = V(\vec{x}) v(K)$. Indeed, this can be most easily argued by the intermediate value theorem if J is connected. For the same reason there is some $\vec{y} \in S$ such that $Q(S) = V(\vec{y}) v(S)$. If you agree to these seeming innocuous statements relating scalar constant velocity normal to J , then the result follows from the Method of Constants.

Originally the basic infinitesimal reasoning behind Theorem 6.4.2 was simply that in the NSP-world $*M(K) \approx *M(S)$, where $\emptyset \neq * \text{int}(*J \cap S)$ and S is a infinitesimal subrectangle of $*R$. However, technically, there seems to be no way to avoid that fact that this idea is incorrect. It may be a good starting point but, infinitesimally, it is necessary that $*M(K)/*v(K) \approx *M(S)/*v(S)$.

Theorem 6.4.3. (Extended Self-evident Method of Constants) *Let the rectangle $R \subset \mathbb{R}^n$ and suppose that Jordan-measurable $J \subset R$. Let M be defined as in (1) of IR4, continuous $f: R \rightarrow \mathbb{R}$, continuous $g: R \rightarrow \mathbb{R}$, $\mathcal{P}$ an acceptable set of partitions of R and any $P \in \mathcal{P}$. If for any $S \in \mathcal{S}(P)$ and $\emptyset \neq K = \text{int}(J \cap S)$ there exists some $\vec{x}_1, \vec{x}_2 \in K$ and some $\vec{y}_1, \vec{y}_2 \in S$ such that (i) $M(K) = f(\vec{x}_1)g(\vec{x}_2)v(K)$ and (ii) $M(S) = f(\vec{y}_1)g(\vec{y}_2)v(S)$, then the infinitesimalizing process IR4 holds for fg restricted to J .*

Why do we need the Extended Self-evident theorem? First, many physical and geometric quantities are defined as the product of other previously defined generating functions. Moreover, it is often the case that these quantities are actually defined for the sole purpose of applying the differential and integral calculus. One example of this should suffice. Consider the following supposedly non-calculus approach to the moment of inertia of a circular disc of radius a about a line ℓ through the center perpendicular to the plane of the disc. In the book *Mechanics of Engineers* [Morley [1942]] the following approach is used.

Assume that the density is unity. . . , consider the disc divided into n “ring-shaped strips such as PQ , each of width a/n .” Morley then takes the distance of the p th strip from the center O as $p \times a/n$. He then lets the area be the same as that of the rectangle $2\pi \times pa/n \times a/n = 2\pi pa^2/n^2$. He then states that the moment of inertia about ℓ would be this area times the distance of the outer edge $p \times a/n$ which yields

$$2\pi a^4 p^3 / n^4. \tag{1}$$

If now we add these n strips this would yield, after simplification, the expression

$$(\pi a^4 / 2) \left(1 + \frac{2}{n} + \frac{1}{n^2}\right). \tag{2}$$

However, on the other hand, if we choose PO to be the distance of the strip from the line ℓ then assuming the everything else holds the expression for the sum all the n strips would be

$$(\pi a^4/2)\left(1 - \frac{2}{n} + \frac{1}{n^2}\right). \quad (3)$$

Observe that for either (2) or (3) the limit as $n \rightarrow \infty$ is $\pi a^4/2$. Morley's definition requires that such a limit be taken. If these ideas are now applied to parallel lines interior to the strips, then the same conclusion would follow. I discuss aspects of this example more fully at the beginning of the next section.

Remark 6.4.1 Closer examination of the Self-evident Theorems 6.4.2 is useful. For a particular linear function M defined as in IR4, the relation between the values $M(S) = 0$, where the subrectangle $S \subset R - \text{int}(J)$, is of a special nature. These values, in general, are assumed to be unaffected by the function f . It is as if the function f has degenerated to the zero function when it is observed by subrectangles exterior to $\text{int}(J)$. In most practical cases this is exactly how it should be. Since we are not interested in the behavior of f on such exterior objects. Two things often happen in practice. The function $f: J \rightarrow (a, b) \subset \mathbb{R}$ is only assumed to be continuous on J . It does not matter whether or not f is the restriction of a function that is continuous on R . The other case is that the function f is explicitly given and it is indeed continuous on some bounding set R . If J is compact and $f: J \rightarrow (a, b)$ is continuous and not explicitly expressed, M is defined as in IR4 and the remaining hypotheses hold for any continuous extension $F: R \rightarrow (a, b)$ of f to R , then by the Teitze Extension Theorem $M(J) = \int_J f(\vec{x}) dX$. The same modifications could be made, if necessary, to the Extended Self-evident Theorem 6.4.3.

Application 6.4.2. (Application 6.3.2 revisited.) *Moment of inertia of a circular disc of radius a about a line ℓ through the center perpendicular to the plane of the disc.*

Consider the disc J centered at the origin of our 2-dimensional Cartesian coordinate system. Let $\rho(x, y)$ represent a continuous point density function for J which is assumed to be continuous on some $R \subset \mathbb{R}^2$ such that $J \subset R$. Then the moment of inertia, $Q(J)$, of J about ℓ a line perpendicular to the disc at its center is

$$Q(J) = \int_J f(x, y) dX,$$

where $f(x, y) = \rho(x, y) (x^2 + y^2)$. Using this assumed function it follows that if ρ is a constant, then

$$Q(J) = \int_J \rho(x, y) (x^2 + y^2) dX = \int_0^a 2\pi \rho r^3 dr = (\pi \rho a^4)/2 = (m/2)a^2.$$

Derivation. Kinetic energy experimentation with a flywheel with movable weights attached to its surface indicates that for actually physical regions J , there does exist a “small enough” simple partition rectangle S such that there are two points $\vec{x}, \vec{y} = (y_1, y_2) \in \text{int}(J \cap S) \neq \emptyset$ and $Q(J \cap S) = \rho(\vec{x}) (y_1^2 + y_2^2)$. For this particular subrectangle S it is also observed that their exist two such points in S with the same property for $Q(S)$. The same conclusion apparently would hold for all other simple partition rectangles with a “smaller diagonal length.” [This generalization to “all” such rectangles appears reasonable.] Let $(m_1, m_2) \in \mathbb{N}^2$ be the generator of the simple partition P such that $S \in \mathcal{S}(P)$. Let the set of partitions $\mathcal{P}$ generated by the pair $\{(x, y) \mid (x, y) \in \mathbb{N}^2 \wedge (x \geq m_1) \wedge (y \geq m_2)\}$. Then $\mathcal{P}$ is an acceptable partition. Letting $f(x, y) = (x^2 + y^2)$, which is continuous on R , then the result follows from the Method of Constant Theorem 6.4.3.

6.5. The Hyperfinite Method.

Except when the operational approach to physical quantities is used, most authors when discussing the properties of rigid body motion immediately force upon the reader the imaginary notion of the point “particle.” Synge and Griffith, in the text mentioned, use exclusively this technique. Also, even though it may not be apparent from his example in the previous section, Morley motivates all of his derivations similarly. It is now possible to establish in a somewhat rigorous fashion that the technique of particles is adequate.

As a prototype, we again concentration upon a nonempty plane Jordan-measurable region J . Letting ρ be an appropriate density function and d and appropriate distance function, from line ℓ of rotation one establishes that the moment of inertia is

$$I(J) = \int_J \rho(\vec{x}) d^2(\vec{x}) dX. \quad (1)$$

By *-transfer of the hypotheses of Theorem 6.4.3 it follows that for a simple fine partition $P = \{S_1, \dots, S_\Gamma\}$ of ${}^*R$ and two hyperfinite sequences, $Q'_1 = \{\vec{x}_1, \dots, \vec{x}_\Gamma\}$, $Q'_2 = \{\vec{y}_1, \dots, \vec{y}_\Gamma\}$, $\vec{x}_i, \vec{y}_i$ members of $\emptyset \neq K_i = {}^*\text{int}({}^*J \cap S_i)$ such that ${}^*I(K_i) = {}^*\rho(\vec{x}_i) {}^*d(\vec{y}_i) {}^*v(K_i)$. Observe that properties of Jordan-measurable sets and the measure v imply that $\text{int}({}^*J \cap S_i)$ is *-Jordan-measurable and that ${}^*v(K_i) = {}^*v({}^*J \cap S_i)$; which leads to ${}^*I({}^*J \cap S_i) = {}^*\rho(\vec{x}_i) {}^*d(\vec{y}_i) {}^*v(K_i) = {}^*\rho(\vec{x}_i) {}^*d(\vec{y}_i) {}^*v({}^*J \cap S_i) = {}^*I(J \cap S_i)$. There is a nonempty internal $\mathcal{A} \subset {}^*\mathbb{N}$ such that $\mathcal{A} = \{n \mid n \in {}^*\mathbb{N} \wedge 1 \leq n \leq \Gamma \wedge J \cap S_n \neq \emptyset\}$. From the additivity of I it follows that

$${}^*I(J) = \sum_{j \in \mathcal{A}} {}^*\rho(\vec{x}_j) {}^*d(\vec{y}_j) {}^*v({}^*J \cap S_j). \quad (2)$$

In the standard case for every $P \in \mathcal{P}$ and $\{S_1, \dots, S_n\} = \mathcal{S}(P)$ there are, of course, two finite sequences $Q_1 = \{\vec{x}_1, \dots, \vec{x}_n\}$, $Q_2 = \{\vec{y}_1, \dots, \vec{y}_n\}$, $\vec{x}_i, \vec{y}_i$ members of $\emptyset \neq \text{int}(J \cap S_i)$ such that

$$I(J) = \sum_{j=1}^n \rho(\vec{x}_j) d(\vec{y}_j) v(J \cap S_j) = \text{st}({}^*I(J)). \quad (3)$$

It's equation (3) above that yields the concept of the particle point masses by defining $m_j = \rho(\vec{x}_j) v(J \cap S_j)$ and, hence,

$$I(J) = \sum_{j=1}^n m_j d(\vec{y}_j) v(J \cap S_j) \quad (4)$$

If now one assumes the particle point mass equation (4) holds for each member of $\mathcal{P}$ then *-transfer yields

$${}^*I(J) = \sum_{j \in \mathcal{A}} {}^*m_i {}^*d(\vec{y}_j) {}^*v({}^*J \cap S_j). \quad (5)$$

The process of introducing (i) the additivity of I , (ii) simplifying the terms of the sum, if possible, by reducing to other geometrical quantities, and (iii) extending to the hyperfinite, is called the **hyperfinite method**. However, this method is unnecessary since it is but a simple extension of the Method of Constants. Note that for non-discrete quantities it is, technically, not correct to say that we “extend the finite sum of quantities to an infinite sum of such quantities” or some similar expression as is often done by Maxwell and others. The sum is not, in general, an (external) infinite sum, but, is an internal hyperfinite sum. The conceptual and formal differences between these two concepts are considerable.

But looking at our definition 5.1.1 for the integral, we are lead to the, usually not appreciated, approximation methods. It follows that for any position real r there is a simple partition P and a finite set of subrectangles $\{S_1, \dots, S_2\} = \mathcal{S}(P)$ such that for any set of intermediate partition points $Q = \{\vec{x}_1, \dots, \vec{x}_n\}$,

$$\left| \sum_{i=1}^n \hat{\rho}(\vec{x}_i) d^2(\vec{x}_i) v(S_i) - I(J) \right| < r. \quad (6)$$

Now each S_i is of one of three types. (a) $S_i \subset R - \text{int}(J)$, (b) $K = \text{int}(J \cap S_i) = \text{int}(J) \cap \text{int}(S_i) \neq \emptyset$ and $\text{int}(S_i) \not\subset \text{int}(J)$, (c) $\text{int}(S_i) \subset \text{int}(J)$. Let ∂A , $A \subset \mathbb{R}^n$ denote the **boundary points** of A . In case (a) since $v(\partial J) = 0$, $v(J) = v(\text{int}(J))$, $\partial J \subset R - \text{int}(J)$ and $v(S) \neq 0$ then there exists some $\vec{x} \in S$ such that $\vec{x} \notin J$. In case (b) and (c) there is some $\vec{x} \in \text{int}(J) \cap S$. Consequently, there is a finite sequence of points $S = \{\vec{y}_1, \dots, \vec{y}_n\}$ that contains a subsequence $S_0 = \{\vec{y}_{k(1)}, \dots, \vec{y}_{k(m)}\}$ such that

$$\left| \sum_{i=1}^m \rho(\vec{y}_{k(i)}) d^2(\vec{y}_{k(i)}) v(S_i) - I(J) \right| < r, \quad (7)$$

where the S_i are all of type (b) or (c) and each $\vec{y}_{k(i)} \in \text{int}(J)$. Yet there exists a simple partition and another sequence $S' = \{\vec{z}_1, \dots, \vec{z}_p\}$ that contains a subsequence $S'_0 = \{\vec{z}_{k(1)}, \dots, \vec{z}_{k(q)}\}$ such that

$$\left| \sum_{i=1}^q \rho(\vec{z}_{k(i)}) d^2(\vec{z}_{k(i)}) v(S_i) - I(J) \right| < r, \quad (8)$$

and all of the rectangles S_i are interior rectangles of type (c) and each $\vec{z}_{k(i)} \in \text{int}(J)$.

Of course these special selections of members in $\text{int}(J)$ extended to the NSP-world and equation (8) reads as

$$\sum_{i=1}^{\Gamma} {}^*\rho(\vec{z}_{k(i)}) {}^*d^2(\vec{y}_{k(i)}) {}^*v(S_i) \approx I(J),$$

and all of the infinitesimal subrectangles S_i are interior subrectangles of type (c) and each $\vec{z}_{k(i)} \in {}^*\text{int}(J)$.

Thus there are many different NSP-world configurations and expressions that have the same N-world effects. These effects are obtain by application of the standard part operator. However, even though it is obvious that all of the procedures discussed in this section and the above numbered expressions give a very detailed and analytically correct approach to infinitesimal modeling - an approach that reveals much about the nature of the NSP-world model - an early introduction of these infinitesimal concepts into an elementary exposition for physical modeling is often unnecessary and, indeed, they will tend to submerge the new discoveries expressed by the self-evident theorems. On the other hand, under certain circumstances, it appears necessary to include these infinitesimal notions. In the next section, I diverge briefly into certain instructional aspects of these discoveries.

6.6. Instruction.

The basic methods of infinitesimal modeling revolve about the application of simplified physical or geometric theories to “simple configurations” within the N-world environment. For the physical sciences, natural system behavior is viewed locally with many of the requirements of the self-evident theorem assumed. Physical modeling is viewed as objective while geometric modeling is subjective. Physical modeling deals with observation and experience with the behavior of natural systems. Geometric modeling, while originally motivated by physical concerns, has become a subject of abstract

definitions. This is obvious from the applications made in chapter 5 for there, the properties of the geometric measures are paramount and are global properties obtained from mathematical experience. The local self-evident rules and even the infinitesimal rules in this chapter are more relevant to natural system behavior. However, both the geometric and physical may be more closely associated with IR3 and IR4 then first assumed.

Experience dictates that the simplest and first level of comprehension for physical concerns is obtained from the two theorems on the Self-evident Method of Constants. They would be the easiest to apply for the neophyte. There is a reason, however, why in certain cases the actual infinitesimal rules IR3 and IR4 need to be applied. This is especially the case for geometric measures. These rules would be a second level of comprehension and this entails a certain basic familiarity with infinitesimal concepts. Moreover, the generating functions for a perceived linear functional need not be the integrand utilized. Indeed, many of the rules and integral notions within chapter 5 and chapter 6 can be expressed by hyperfinite summation that yields internal functionals not just the standard extension of a standard concept. An example of this is our proof of the “length of a curve integrand” in appendix 6 on pages 209–210.

In the proof for the length of a continuously differentiable curve $c: [0, 1] \rightarrow \mathbb{R}^n$ an internal functional is defined for the length of a, possibly broken, hyperpolygonal line determined by a hypercurve $*c: *[0, 1] \rightarrow *\mathbb{R}^n$ with coordinates functions $*f_j: *[0, 1] \rightarrow *\mathbb{R}$. This internal function may be consider defined as follows: let $\{t_i\}$ be any hyperfinite partition of $*[0, 1]$. Let T be any nonempty internal, hence hyperfinite, subset of $\{t_i\}$ containing two or more members. For each pair of elements in T generate the hyperline segments determined by the curve $*c$ and consider $\|\vec{v}_i\|$ from Example 4.4.1.A. Finally, consider the internal hyperfinite sum operator over the internal set of the $\|\vec{v}_i\|$ determined by T . Let L denote this internal functional. Now for each internal subinterval S determined by T define $L(S) = L(*\text{int}(S))$. This is our $*$ -additive functional defined over the set of all such internal functions c . Under the given hypotheses it is shown that for any internal subrectangle S , taken from a simple fine partition of $*[0, 1]$ there exists some $t_i \in S$ such that $\|\vec{v}_i\|/dt \approx \sqrt{\sum_{j=1}^n (*f'_j(t_i))^2}$. The expression on the right is the standard extension of the standard function $\sqrt{\sum_{j=1}^n (f'_j(t_i))^2}$ while the expression on the left is $L(S)/dt$. Thus our basic definition for the length of a curve - one that with a slight generalization is also shown to be equivalent to rectifiability - almost satisfies IR4 with the exception that IR4 is written in terms of a standard additive functional not an internal $*$ -additive nonstandard entity. It is obvious how IR3 and IR4 would be modified to include such internal functionals.

The fact that geometric definitions are subjective in character would allow us to define many such concepts entirely in terms of infinitesimal concepts. This would be an aid in developing many integral statements about n -dimensional geometric properties in terms of integrals defined on k -dimensional spaces, where $k < n$.

6.7 Realism.

I will not, at this point, dwell upon the philosophical modeling concept termed “realism.” This concept assumes that all mathematical objects within the mathematical formalism correspond to objects within objective (physical) reality. There is, however, an interesting historical fact relative to the rejection of realism. Even though Bohr rejected complete realism for his concept of quantum mechanics in order to force a type of physical completeness upon this theory - a completeness that is now known to be in error [Aerts [1984]] - he was not the first to do so. In his 1909 treatise on his theory of electrons, Lorentz accepted the notion of an ether but rejected complete realism. “I

should add that, while thus denying the real existence of ether stresses, we can still avail ourselves of all the mathematical transformations by which application of the formula (43) may be made easier. We need not refrain from reducing the force to a surface-integral, and for convenience's sake we may continue to apply to the quantities occurring in this integral the name of stresses. Only we must be aware that they are only imaginary ones, nothing else than auxiliary mathematical quantities. Perhaps all this that has been said about the absolute immobility of the ether and the non-existence of the stresses, may seem somewhat startling." Lorentz [1952:31]

As far as geometric modeling is concerned the concept of realism is not as significant as when it is applied to physical problems. Infinitesimal modeling may be considered as a partial realism. The notion of a continuum model in an ultimately discrete world seems non-realistic. However, what is being modeled is macroscopic and large scale behavior, macroscopic and large scale effects upon observers and other objects. It is not the actual physical entities that are being modeled; but, rather, the effects these entities produce relative to a specific set of scenarios.

However, on another level, modern physical theories often deal with entities that are not directly observable, only their indirect effects emerge within the laboratory setting. Technically, these entities are speculations that may in time be replaced by yet other speculative objects. Indeed, using special techniques, the language that actually expresses physical theories is mathematically embedded into the natural numbers by a Gödel coding; which is then embedded into a nonstandard structure. The linguistics of the physical theory produces a new collection of statements that tells us about the behavior of a new world, called in general, the NSP-world. [Herrmann 1987] Within this world we are not speculating ad hoc about infinitesimal objects, it is the standard theory that generates their logical existence and even predicts some of their extraordinary properties. Whether or not such entities are accepted as "real" or not depends upon their usefulness. In case you may have missed it previous, I again quote the following from the first paper Robinson published relative to his formal theory of infinitesimals. "*For phenomena on a different scale, such as considered in Modern Physics, the dimensions of a particular body or process may not be observable directly. Accordingly the question whether or not a scale of non-standard analysis is appropriate to the physical world really amounts to asking whether or not such a system provides a better explanation of certain observable phenomena than the standard system of real numbers. The possibility that this is the case should be borne in mind.*" Fine Hall, Princeton University. [Robinson 1961]

Chapter 7.

**PURE INFINITESIMAL
INTEGRAL MODELING**

7.1 Brief Discussion.

This basic manual is intended to present brief accounts of various approaches to infinitesimal modeling so that individuals may select the method the is most appropriate for their discipline. In Chapter 5, the standard and global type rules for integral modeling are introduced in a piecemeal fashion. Each of the applications in that chapter depends upon an in depth knowledge of the properties of a specific functional that measures a specific geometric or physical quantity. Actually, as is clear from the derivations, it is more likely that the properties of the specific functional are selected in concert with the requirements of the Infinite Sum Theorem. As for Chapter 6, although the Self-evident theorems are more general in character, they still rely upon the Infinite Sum Theorem as the proofs of Theorems 6.2.2 and 6.4.1 indicate. However, in general, neither the approach of Chapter 5 nor the approach of Chapter 6 appears within the pre-1960 literature. Previous appliers of the concepts of infinitesimal reasoning relied heavily upon a pure infinitesimal approach that is highly discipline orientated and which makes a direct appeal to the Riemann styled sum and Definition 5.1.1. This pure infinitesimal approach relies upon expedient definitions for elusive geometric or physical entities within the NSP-world - entities called by the general term “elements.” The collection for geometric modeling includes the basic elements of IR² in Chapter 6 and various Euclidean compositions and decompositions of these configurations.

For each discipline, the collection of elements is sequentially defined from the most basic through the more complex in a, hopefully, consistent manner. It is almost always the case that the defined objects are taken from those in the N-world that behave in the simplest possible discipline defined manner and this simplistic behavior is then impressed by *-transfer upon the NSP-world. For geometry, the selection of an acceptable set of elements is somewhat more arbitrary than for physical disciplines. The experimental or observational disciplines rely upon the concept of what might be termed as simple or idealized behavior and well-grounded physical methods of approximation. However, after the infinitesimal elements have been selected then, in all cases of which I am aware, the rigorous derivations that these elements lead to an integral model are very similar. Thus, there may be an unconscious interplay between the appropriate element, as axiomatically selected, and the rigorous method. Examination of the literature leads to the following general observations.

(i) The infinitesimal elements are axiomatically selected, per discipline, and the concept of (hyperfinite) summation is applied. Whether or not this summation can be calculated by means of a Riemann styled integral is somewhat secondary.

(ii) In anticipation of an analytical approach, it has become customary to employ certain symbols and terms that tend to describe inaccurately the actual situation. As a prototype consider the symbol ds . In infinitesimal modeling this is a **general symbol** that represents any of the lengths of hyperline segments that comprise any of the hyperpolygonal curves that determine the length of a geometric curve. With respect to our notation $ds = \|\vec{v}_i\|$. Only when actual attempts are made to calculate the length of the entire curve are additional analytical constraints placed upon the representing functions. For modeling purposes, the geometric or physical elements are considered more basic in character in order to avoid, if possible, all of the well-known analytical difficulties.

(iii) Restriction to the monadic environment takes place only when an n-dimensional quantity

is being measured with respect to an n -dimensional partition. Otherwise, the infinitesimal elements span both the N -world and the NSP -world; portions are monadic and portions are not.

(iv) The selected infinitesimal elements are not unique, even in their general character.

In the next section, certain elements will be selected for problems in elementary infinitesimal geometry. Following this, analytical constraints will be applied so that elementary integral expressions can be rigorously derived. These rigorous derivations display common derivation features that should be apparent. Indeed, the common features already appear in the proof on page 127 that gives the length of a continuously differentiable curve relative to the hyperpolygonal approximating curves discussed in examples 4.4.1.A and 4.4.1.B.

7.2 Geometric Elements.

The first and most basic measure that appears in Euclid's geometry book is the length of a line segment. I repeat, with slight modifications, the information in example 4.4.1.A for the generalization of this to curves.

Recall that a curve is a continuous map $c: [0, 1] \rightarrow \mathbb{R}^n$. This is equivalent to considering c as determined by n continuous coordinate functions $x_i = f_i(t)$, $1 \leq i \leq n$ each defined on $[0, 1] \subset \mathbb{R}$. The geometric curve C determined by these functions is usually considered as the set $\{(x_1, \dots, x_n) | t \in [0, 1]\}$. The $*$ -transform of these defining functions leads to the functions $x_i = {}^*f_i(t)$, $1 \leq i \leq n$, each defined on ${}^*[0, 1] \subset {}^*\mathbb{R}$ and they generate the "hypercurve" ${}^*C \subset {}^*\mathbb{R}^n$.

Let Q be any fine partition of ${}^*[0, 1]$. By $*$ -transfer, Q behaves like an ordered partition of the interval $[0, 1]$ as defined in the standard sense and we write $Q = \{t_i \mid 0 \leq i \leq \Gamma\}$. The internal hyperfinite set Q generates the internal set of "points" $P = \{({}^*f_1(t_i), \dots, {}^*f_n(t_i)) \mid t_i \in Q\}$ that are members of the hypercurve ${}^*C$. Now for each $i = 0, \dots, \Gamma - 1$, and each j , $0 \leq j \leq n$ let ${}^*f_j(t_{i+1}) - {}^*f_j(t_i) = d(j, i)$. Since c is continuous then each $d(j, i) \in \mu(0)$. For each $i \in {}^*\mathbb{N}$ such that $0 \leq i \leq \Gamma - 1$, the internal set $\ell_i = \{(x_1, \dots, x_n) \mid \forall j \in {}^*\mathbb{N}, 0 \leq j \leq n, x_j = {}^*f_j(t_i) + t(d(j, i)) \wedge t \in {}^*[0, 1]\}$ is a hyperline segment connecting the two points $({}^*f_1(t_i), \dots, {}^*f_n(t_i))$, $({}^*f_1(t_{i+1}), \dots, {}^*f_n(t_{i+1}))$ on the curve ${}^*C$. From this one obtains the internal hyperpolygonal curve $\mathcal{P}_Q = \bigcup \{\ell_i \mid 0 \leq i \leq \Gamma - 1\}$. As to the length of $\mathcal{P}_Q$ simply extend the concept of length in the classical sense by defining for each $i = 0, \dots, \Gamma - 1$ the vector $\vec{v}_i = (d(1, i), \dots, d(n, i)) \in {}^*\mathbb{R}^n$. Then let the hyperfinite sum $\sum_{i=0}^{\Gamma-1} \|\vec{v}_i\| = |\mathcal{P}_Q| \in {}^*\mathbb{R}$. **For the curve, the geometric element is the hyperline segment ℓ_i .** Even though, in general, you would have a different hyperpolygon with a different hyperreal length for $\forall \Gamma \in \mathbb{N}_\infty$ the following Proposition is proved in appendix 6.

Theorem 7.2.1. *Consider continuous $c: [0, 1] \rightarrow \mathbb{R}^n$. Then c is rectifiable if and only if there exists some $L \in \mathbb{R}$ such that for every fine partition Q of ${}^*[0, 1]$*

$$\text{st}({}^*|\mathcal{P}_Q|) = L.$$

Application 7.2.1. *The 2-dimensional area between two continuous curves using the 1-dimensional integral.*

Give continuous $f: [a, b] \rightarrow \mathbb{R}$ and continuous $g: [a, b] \rightarrow \mathbb{R}$, where $f(x) \leq g(x)$ for each $x \in [a, b]$. Then the area between the curves f and g is

$$\int_a^b (g(x) - f(x)) dx.$$

Derivation. From the NSP-world view point each curve is viewed as any hyperpolygonal line created by any simple fine partition of $^*[a, b]$. Let Q be such a simple fine partition. For a given $I_i = [x_i, x_{i+1}] \in ^*\mathcal{S}(Q)$, let $\ell(f)_i$ be the corresponding hyperline segment determined by the curve f and $\ell(g)_i$ that determined by g for the interval I_i . Select as the infinitesimal element the **hypertrapezoid**, T_i , with sides $\ell(f)_i$, $\ell(g)_i$ and the hyperline segments with end points $\{(x_i, ^*g(x_i), (x_i, ^*f(x_i)))\}$ and $\{(x_{i+1}, ^*g(x_{i+1}), (x_{i+1}, ^*f(x_{i+1})))\}$. Observe that this hypertrapezoid selection is consistent with the NSP-world view of a curve and the hypertrapezoid is composed of * -Euclidean composition or decomposition of our previously defined basic elements. By * -transfer the hypertrapezoid's hyperarea $^*A(T_i) = (1/2)((^*g(x_i) - ^*f(x_i)) + (^*g(x_{i+1}) - ^*f(x_{i+1}))) dx$. As was done for the length of a curve, let $T(Q) = \bigcup\{T_i \mid 0 \leq i \leq \Gamma - 1\}$ and consider the hyperfinite sum

$$\sum_{I_i \in ^*\mathcal{S}(Q)} ^*A(T_i) = ^*A(T(Q)). \quad (1)$$

From this definition it follows that

$$\text{st}(^*A(T(Q))) = \int_a^b (g(x) - f(x)) dx. \quad (2)$$

$\triangle$ Rather than relegate the proof that there exists a unique $r \in \mathbb{R}$ such that for any simple fine partition Q of $^*[a, b]$, $^*A(T(Q)) \in \mu(r)$ to the appendix we present it as follows by investigating the behavior of the expression $((^*g(x_i) - ^*f(x_i)) + (^*g(x_{i+1}) - ^*f(x_{i+1})))$. From the uniform continuity of g, f it follows that $(^*g(x_i) - ^*f(x_i)) \approx (^*g(x_{i+1}) - ^*f(x_{i+1}))$. Hence, $^*A(T_i) = ((^*g(x_i) - ^*f(x_i)) + \delta_i) dx$, where $\delta_i \in \mu(0)$. Consequently,

$$\sum_{I_i \in ^*\mathcal{S}(Q)} ^*A(T_i) = \sum_{I_i \in ^*\mathcal{S}(Q)} ((^*g(x_i) - ^*f(x_i)) dx + \sum_{I_i \in ^*\mathcal{S}(Q)} \delta_i dx. \quad (3)$$

By considering the finite case it follows that there exists some $\delta \in \{|\delta_i| \mid 0 \leq i \leq \Gamma - 1\}$ such that $\delta = \max\{|\delta_i| \mid 0 \leq i \leq \Gamma - 1\}$ and, hence, $\delta \in \mu(0)$. By * -transfer we have that

$$\left| \sum_{I_i \in ^*\mathcal{S}(Q)} \delta_i dx \right| \leq \sum_{I_i \in ^*\mathcal{S}(Q)} |\delta_i| dx \leq \delta \sum_{I_i \in ^*\mathcal{S}(Q)} dx = \delta(b - a) \in \mu(0). \quad (4)$$

Thus

$$\sum_{I_i \in ^*\mathcal{S}(Q)} \delta_i dx = \lambda \in \mu(0). \quad (5)$$

Therefore, (2) can now be written as

$$\sum_{I_i \in ^*\mathcal{S}(Q)} ^*A(T_i) = \sum_{I_i \in ^*\mathcal{S}(Q)} ((^*g(x_i) - ^*f(x_i)) dx + \lambda \quad (6)$$

Since the function $(g(x) - f(x))$ is bounded and integrable and the partition Q can also be considered as an internal immediate partition then Theorem 5.2.2 yields

$$\text{st}\left(\sum_{I_i \in ^*\mathcal{S}(Q)} ^*A(T_i)\right) = \text{st}\left(\sum_{I_i \in ^*\mathcal{S}(Q)} ((^*g(x_i) - ^*f(x_i)) dx\right) = \int_a^b (g(x) - f(x)) dx. \quad (7)$$

and this completes the proof. $\triangle$

The above derivation certainly appears interesting except for the obvious fact that the **hypertrapezoid configuration is not a unique geometric element**. The key to this derivation is the process that begins with the statement that the values are infinitely close ($\approx$) and the expressions (3)(4)(5)(6)(7). The derivation would hold for hyperrectangles, hypertriangles and the like. Hyperrectangles yield the same 1-dimensional integral expressions. Hypertriangles yield 1/2 the indicated integral expression, and so forth. This non-uniqueness for elements is sometimes considered one of the basic difficulties with infinitesimal modeling both for the geometric and physical. However, since the concept of area obtained by integrals is supposed to be an extension of the Euclidean area measure then any element that does not yield such an extension can be rejected. *For elementary calculus, the use of the hypertrapezoid is very expedient when the usual 1-dimensional integral expressions for the basic geometric measures are considered. It will be our element of choice in this case.*

Application 7.2.2. *Volume of revolution using the 1-dimensional integral.*

Given continuous $f: [a, b] \rightarrow \mathbb{R}$ where $f(x) \geq 0$ for each $x \in [a, b]$. Then the volume obtained by rotating this curve about the x -axis is

$$\int_a^b \pi f^2(x) dx.$$

Derivation. Let Q be a simple fine partition of $^*[a, b]$. For a given $I_i = [x_i, x_{i+1}] \in {}^*\mathcal{S}(Q)$, let $\ell(f)_i$ be the corresponding hyperline segment determined by f . Consider the hypertrapezoid composed of hyperline segments $\ell(f)_i$, I_i and hyperline segments with end point $\{(x_i, 0), (x_i, {}^*f(x_i))\}$, $\{(x_{i+1}, 0), (x_{i+1}, {}^*f(x_{i+1}))\}$. The rotation of this hypertrapezoid about the x -axis is the frustum of a right cone, F_i . By * -transfer, the volume of such a * -geometric configuration is ${}^*V(F_i) = (1/3)\pi(f^2(x_i) + f(x_i)f(x_{i+1}) + f^2(x_{i+1}))$. Assuming the usual * -additivity, the volume of the entire configuration $F(Q) = \bigcup \{F_i \mid 0 \leq i \leq \Gamma - 1\}$ is

$$\sum_{I_i \in {}^*\mathcal{S}(Q)} {}^*V(F_i) = {}^*V(F(Q)). \quad (8)$$

From this definition it will follow that

$$\text{st}({}^*V(F(Q))) = \int_a^b \pi f^2(x) dx. \quad (9)$$

Noting that $f^2(x_i) + f(x_i)f(x_{i+1}) + f^2(x_{i+1}) \approx f^2(x_i) + f(x_i)f(x_i) + f^2(x_i) = 3f^2(x_i)$ then (9) follows in the same manner as in the derivation for application 7.2.1.

Application 7.2.3. *The surface area of revolution using the 1-dimensional integral.*

Given continuous differentiable $f: [a, b] \rightarrow \mathbb{R}$ where $f(x) \geq 0$ for each $x \in [a, b]$. Then the surface area obtained by rotating this curve about the x -axis is

$$\int_a^b 2\pi f(x) \sqrt{1 + (f'(x))^2} dx.$$

Derivation. Let Q be a simple fine partition of $^*[a, b]$. For a given $I_i = [x_i, x_{i+1}] \in {}^*\mathcal{S}(Q)$, let $\ell(f)_i$ be the corresponding hyperline segment determined by f . Consider the hypertrapezoid composed of hyperline segments $\ell(f)_i$, I_i and hyperline segments with end point

$\{(x_i, 0), (x_i, {}^*f(x_i))\}, \{(x_{i+1}, 0), (x_{i+1}, {}^*f(x_{i+1}))\}$. The rotation of this hypertrapezoid about the x -axis is the frustum of a right cone, F_i . By $*$ -transfer, the surface area of such a $*$ -geometric configuration is $S(F_i) = \pi({}^*f(x_i) + {}^*f(x_{i+1})) {}^*|\ell(f)_i|$, where ${}^*|\ell(f)_i|$ is the $*$ -length of the hyperline segment. In appendix 6, ${}^*|\ell(f)_i| = \|\vec{v}_i\|$. Assuming the usual $*$ -additivity, the surface area of the entire configuration $F(Q) = \bigcup \{F_i \mid 0 \leq i \leq \Gamma - 1\}$ is

$$\sum_{I_i \in {}^*\mathcal{S}(Q)} {}^*S(F_i) = {}^*S(F(Q)). \quad (10)$$

From this definition it will follow that

$$\text{st}({}^*S(F(Q))) = \int_a^b 2\pi f(x) \sqrt{1 + (f'(x))^2} dx. \quad (11)$$

Now the complete derivation uses the proof of the integral length formula that appears in appendix 6. There it is shown that $\|\vec{v}_i\| = \sqrt{1 + ({}^*f'(x_i))^2} dx + \delta_i dx$, where $\delta_i \in \mu(0)$. But, ${}^*f(x_i) = {}^*f(x_{i+1}) + \lambda_i$, $\lambda_i \in \mu(0)$. Therefore, ${}^*f(x_i) + {}^*f(x_{i+1}) = 2{}^*f(x_i) + \lambda_i$. Since f is bounded then ${}^*S(F_i) = \pi({}^*f(x_i) + {}^*f(x_{i+1})) {}^*|\ell(f)_i| = 2\pi f(x_i) \sqrt{1 + ({}^*f'(x_i))^2} dx + \gamma_i dx$, $\gamma_i \in \mu(0)$. The derivation is completed by application of steps (3)(4)(5)(6)(7) as demonstrated in application 7.2.1.

⇓ IMPORTANT ⇓

⇒ Please note that for applications 7.2.1, 7.2.2, and 7.2.3 the actual defining geometric quantities are given by equations (1), (8) and (10), respectively. The geometric configurations, $T(Q)$ and $F(Q)$ composed of the hyperfinite union of the respective infinitesimal elements may be considered as members of the nonstandard extension of the set of all ordinary Euclidean configurations. Configurations $T(Q)$, $F(Q)$ are internal subsets of appropriate $*$ -Euclidean entities that do exhibit a standard area or volume measure. By $*$ -transfer of the standard properties of geometric measures relative to subsets it follows that the standard part of each of these expressions exists as a real number. What is established in these applications under the analytical constraints given is that the standard part is expressible by the indicated integral. ⇐

Relative to the infinitesimal geometric elements, I have not altered the requirement that for the n -dimensional integral these elements be the infinitesimal subrectangles. In the latter sections of this manual, since it is written for individuals with a strong undergraduate mathematics background, the Jordan-measurable generalization for the basic rectangular region was used. As far as a Jordan-measurable J is concerned do we ever need to consider any subrectangle that is not in the interior of J ?

Let Jordan-measurable $J \subset R \subset \mathbb{R}^n$ and let Q be any fine partition of R . It is sometimes useful to assume that J is a closed subset of R (hence, compact). For, if J is not closed, then recall that the Jordan-content of J , $v(J) = v(\overline{J})$. **Intuitively, think of J as any of the ordinary regions studied in elementary calculus with their intuitive boundary, inner and exterior portions.** Using these ideas of boundary, inner and exterior portions of J , the set of all subrectangles determined by Q , which is denoted by ${}^*\mathcal{S}(Q)$, may be separated formally into three disjoint hyperfinite collections. Let the set of **boundary subrectangles** be

$\partial(Q) = \{S \mid S \in {}^*\mathcal{S}(Q) \wedge S \cap {}^*J \neq \emptyset \wedge S \cap ({}^*\mathbb{R}^n - {}^*J) \neq \emptyset\}$. Observe that if $S \in \partial(Q)$, then $S \subset \mu(p)$ and $\mu(p) \cap {}^*J \neq \emptyset$ and $\mu(p) \cap ({}^*\mathbb{R}^n - {}^*J) \neq \emptyset$ imply that $p \in \partial(J)$. Now consider the set of **exterior subrectangles** $\text{ext}(Q) = \{S \mid S \in {}^*\mathcal{S}(Q) \wedge S \subset ({}^*\mathbb{R}^n - {}^*J)\}$. Finally, the set of **inner subrectangles** is $\text{inn}(Q) = \mathcal{S}(Q) - (\partial(Q) \cup \text{ext}(Q))$. Observe that the boundary subrectangles generate boundary points of J . However, by considering J to be a rectangle or the interior of a rectangle then it is clear that there may exist some $S \in \text{inn}(Q)$ or $S \in \text{ext}(Q)$, respectively, such that $S \subset \mu(p)$ and $p \in \partial(J)$. It is definitely the case, however, that $S \subset {}^*J$ if and only if $S \in \text{inn}(Q)$. Of course the sets $\partial(Q)$, $\text{inn}(Q)$ and $\text{ext}(Q)$ are mutually disjoint. This leads to the very useful

Theorem 7.2.2. *Let Jordan-measurable $J \subset R \subset \mathbb{R}^n$, bounded $f: J \rightarrow \mathbb{R}$, and Q be any fine partition of ${}^*R$. Let hyperfinite ${}^*\mathcal{S}(Q) = \{S_i \mid 0 \leq i \leq \Gamma - 1\}$. Assume that there exists a hyperfinite sequence U_i , where $U_i \in S_i$ for each $S_i \in \mathcal{S}(Q)$. Then*

$$\text{st}\left(\sum_{S(i) \in \text{inn}(Q)} {}^*f(U_i) {}^*v(S_i)\right) = \text{st}\left(\sum_{S(i) \in \mathcal{S}(Q)} {}^*\hat{f}(U_i) {}^*v(S_i)\right).$$

Theorem 7.2.2 indicates, as expected, that only the interior infinitesimal subrectangles are significant when modeling with respect to the Jordan-content of any set.

Application 7.2.4. *Volume obtained by 2-dimensional integral.*

Suppose that compact Jordan-measurable $J \subset R \subset \mathbb{R}^n$. Let continuous $f: J \rightarrow \mathbb{R}$. For each $\vec{x} \in J$ let $f(x) \geq 0$. Then the volume between the surface determined by f and the xy -plane is

$$\int_J f(x) dX.$$

Derivation. Let Q be a simple fine partition of R and consider some $S_i \in \mathcal{S}(Q)$. From the NSP-world viewpoint, the surface curves obtained by intersecting the surfaces with planes parallel to the coordinate planes are hyperpolygonal lines. Since the choice of the particular hyperline segments is arbitrary and $S_i = [x_i, x_{i+1}] \times [y_i, y_{i+1}]$ then consider the * -Euclidean configuration $TR(S_i)$ composed of the truncated hyperrectangular solid with S_i as its base and its top a parallelogram with adjacent sides the hyperline segments $\{(x_i, y_i, {}^*\hat{f}(x_i, y_i), (x_i, y_{i+1}, {}^*\hat{f}(x_i, y_{i+1}))\}$ and $\{(x_i, y_i, {}^*\hat{f}(x_i, y_i), (x_{i+1}, y_i, {}^*\hat{f}(x_{i+1}, y_i))\}$. Using the * -Euclidean measure for the volume ${}^*V(T(S_i))$ then once again * -additivity yields for the configuration $TR(Q) = \bigcup \{TR(S_i) \mid 0 \leq i \leq \Gamma - 1\}$

$$\sum_{S(i) \in \mathcal{S}(Q)} {}^*V(TR(S_i)) = {}^*V(TR(Q)). \quad (12)$$

Investigating the various configurations $TR(S_i)$, and using the * -Euclidean measure of these configurations then Theorem 7.2.2 yields that

$$\sum_{S(i) \in \text{inn}(Q)} {}^*V(T(S_i)) \approx \sum_{S(i) \in \mathcal{S}(Q)} {}^*V(S_i). \quad (13)$$

Since J is compact then

$${}^*f(x_i, y_i) \approx {}^*f(x_i, y_{i+1}) \approx {}^*f(x_{i+1}, y_i). \quad (14)$$

Applying the * -Euclidean measure formula yields that ${}^*V(T(S_i)) = ({}^*f(x_i, y_i) + \delta_i) dX$ and the method of Application 7.2.1 yields

$$\text{st}\left(\sum_{S(i) \in \text{inn}(Q)} {}^*V(T(S_i))\right) = \text{st}\left(\sum_{S(i) \in \mathcal{S}(Q)} {}^*V(S_i)\right) = \int_J f(x) dX. \quad (15)$$

For integral modeling, physical infinitesimal elements are based upon the Method of Constants. It assumes that the physical quantity being considered has the same effect as if it were concentrated at some point within subrectangle or another similar object. The method is essentially outlined by Maxwell in his previously quoted descriptions. Clearly, such infinitesimal physical modeling is highly discipline orientated and is closely associated with simple mind experiments.

Application 7.2.5. *The value of the electric field vector at a point P exterior to the plane of a charged 2-dimensional closed and bounded Jordan-measurable region.*

Consider a Jordan-measurable region J , a point $P = (a, b, c)$, ($c \neq 0$) in space exterior to the plane. Let $\rho(x, y, 0)$ be a continuous charge density function defined on J . The scalar value of the electric field at P is

$$|\vec{E}(a, b, c)| = \int_J \frac{\rho(x, y, 0)}{(x - a)^2 + (y - b)^2 + c^2} dX.$$

Derivation. (The method of point charges.) For a single point charge q at a distance r from P , the definition of the scalar value of the electric field is $|\vec{E}| = q/r^2$. Let $J \subset R \subset \mathbb{R}^2$, $(x_i, y_i, 0) \in J$, and consider a simple fine partition Q of R . If one considers a point charge with value ${}^*\rho(x, y, 0) {}^*v(S_i)$; $(x, y, 0) \in S_i \in \text{inn}(Q)$, then $|\vec{E}(a, b, c)| = ({}^*\rho(x', y', 0) {}^*v(S_i)) / ((x - a)^2 + (y - b)^2 + c^2)$. But the function $h(x, y) = \rho(x, y, 0) / ((x - a)^2 + (y - b)^2 + c^2)$ is uniformly continuous on J . Thus

$$\begin{aligned} {}^*\rho(x', y', 0) {}^*v(S_i) &= ({}^*\rho(x_i, y_i, 0) + \delta_i) {}^*v(S_i) = \\ &{}^*\rho(x_i, y_i, 0) {}^*v(S_i) + \delta_i {}^*v(S_i), \end{aligned}$$

where $\delta_i \in \mu(0)$. Assuming that the scalar value of the electric field for any nonempty finite set of point charges is the sum of the individual values and that within any 2-dimensional rectangle, S , with charge density $\rho(x, y, 0)$ there is some point where the entire charge can be considered as concentrated, or that for a constant charge density ρ the total charge is $\rho v(S)$, then $*$ -transfer and the method used in applications 7.2.1 and 7.2.4 imply that the scalar value of the electric field at P due to the charge on J is

$$\begin{aligned} |\vec{E}_J(P)| &= \text{st}(|{}^*\vec{E}_J(P)|) = \\ \text{st}\left(\sum_{S(i) \in \text{inn}(Q)} ({}^*\rho(x', y', 0) {}^*v(S_i)) / ((x - a)^2 + (y - b)^2 + c^2)\right) &= \\ \int_J \frac{\rho(x, y, 0)}{(x - a)^2 + (y - b)^2 + c^2} dX. \end{aligned}$$

Remark: In 7.2.5, it is not correct to simply consider the points $(x', y', 0) \in S_i$. It is always necessary that the chosen intermediate partition be internal. Further, extending the concept of the constant charge density is a major approach to infinitesimal modeling since if $\rho(x, y, 0)$ is continuous on compact J , then ρ behaves in a constant-like manner in that if infinitesimal $S \subset J$ and $p, q \in S$, then $p \approx q$ implies that $\text{st}(\rho(p)) = \text{st}(\rho(q))$. Using the physical element concept, this is called the **elemental method of constants**.

Chapter 8.

REFINEMENTS FOR INTEGRAL MODELING

8.1 A Very General Approach.

Hurd and Loeb [1985] construct a very general integral concept. For example, consider any hyperfinite $\{x_1, \dots, x_\Gamma\} \subset X$, let B be any set of internal hyperreal-valued functions defined on X , and $\{a_1, \dots, a_\Gamma\}$ a fixed set of hyperreal nonnegative numbers. They then consider the hyperfinite sum operator $\sum_\Gamma$ defined on each $f \in B$ by $\sum_\Gamma f = \sum_1^\Gamma a_i f(x_i)$. This and other examples are generalized and an entire theory of integration is developed that incorporates various classical generalizations of the Riemann integral, especially the Lebesgue. They apply their theory to stochastic processes such as the Poisson process and Brownian motion. Since the background necessary to study their generalization is beyond the scope of these manuals, their theory will not be presented. Indeed, except for concept of hyperfinite summation, their general approach is probably unsuited for elementary modeling. Our goal in this last chapter on integral modeling is to examine more closely the specific contents of the hyperfinite sum as defined in definition 5.1.1 relative to modifications of its geometric or physical meaning.

Recall definition 5.1.1. Let $f: R \rightarrow \mathbb{R}$ be bounded and $\mathcal{P}$ the set of simple partitions of R . Then f is said to be integrable if there exists some $r \in \mathbb{R}$ and a simple fine partition, $P \in {}^*\mathcal{P}$ such that for each of its internal intermediate partitions $Q = \{\vec{v}_q\}$, where $1 \leq q \leq \Gamma \in \mathbb{N}_\infty$, it follows that $\sum_{k=1}^\Gamma {}^*f(\vec{v}_q) {}^*v(R_q) \in \mu(r)$.

Obviously, modifications can be made in the concept of the basic partition P or the intermediate partition Q . Such modifications are discussed at the conclusion of this chapter. It is important to stress at this point something that is not apparent about this definition. It is immediate from examination of the proofs in appendix 5, that, basically, there are two reasons why this definition works. First, ${}^*v(R_q) \in \mu(0)$. And, secondly, due to behavior of the measure v , the integral value r is independent of the fine partition chosen. This must be taken into consideration if the partition concepts are not to be altered. For the basic modifications that follow, we extend the partitioning requirement to any arbitrary fine partition and any arbitrary intermediate partition and modify the terms of the hyperfinite sum with a view towards applications as they appear in elementary calculus courses.

8.2. The Line Integral.

Let bounded $\phi: [a, b] \rightarrow \mathbb{R}$. Our first consideration is to modify the value of the *-measure ${}^*v([t_{i-1}, t_i])$, where $[t_{i-1}, t_i] \subset {}^*[a, b]$. We know the importance of the “increment” in the calculus, hence, our first modification is to replace ${}^*v([t_{i-1}, t_i])$ with the standard extension of the increment operator. In place of ${}^*v([t_{i-1}, t_i])$ write ${}^*\phi(t_i) - {}^*\phi(t_{i-1})$. What happens if ϕ is continuous? Well, in that case, if $t_i - t_{i-1} \in \mu(0)$, then ${}^*\phi(t_i) - {}^*\phi(t_{i-1}) \in \mu(0)$ and one of the most basic requirements for that factor of the term of the hyperfinite sum is met.

Definition 8.2.1. (Riemann - Stieltjes Integral). Let bounded $f: [a, b] \rightarrow \mathbb{R}$ and bounded $\phi: [a, b] \rightarrow \mathbb{R}$. Then f is **RIEMANN STIELTJES Integrable with respect to ϕ** if there exists a real r such that for any fine partition $P = \{a = t_0, \dots, t_\Gamma = b\}$ of $[a, b]$ and any internal intermediate

partition $Q = \{t'_1, \dots, t'_\Gamma\}$ of P it follows that

$$\sum_{i=1}^{\Gamma} {}^*f(t'_i) ({}^*\phi(t_i) - {}^*\phi(t_{i-1})) \in \mu(r).$$

Rather than investigate the Riemann-Stieltjes integral as defined in 8.2.1, I pass directly to the line integral of elementary calculus. Referring to example 4.4.1A, let bounded $c: [a, b] \rightarrow \mathbb{R}^n$, $c(t) = (c_1(t), \dots, c_n(t))$ be considered a curve with graph C . Assume that $E \subset \mathbb{R}^n$, $C \subset E$ and bounded $F: E \rightarrow \mathbb{R}^n$. Write F as $F(\vec{x}) = (f_1(\vec{x}), \dots, f_n(\vec{x}))$. Consider the composite function $(Fc): [a, b] \rightarrow \mathbb{R}^n$ defined by $(Fc)(t) = (f_1(c(t)), \dots, f_n(c(t)))$, $t \in [a, b]$. Notice that if $t'_j \in [t_{j-1}, t_j] \subset {}^*[a, b]$, then, letting $\bullet$ denote the “dot” (inner) product,

$${}^*(Fc)(t'_j) \bullet {}^*\vec{v}_j = \sum_{i=1}^n {}^*f_i({}^*c(t'_j)) ({}^*c_i(t_{j-1}) - {}^*c_i(t_j)),$$

where ${}^*\vec{v}_j = {}^*c(t_j) - {}^*c(t_{j-1})$ represents the directed hyperline segment portion of some hyperpolygonal curve representation for C . Now taking a hyperfinite sum yields

$$\begin{aligned} \sum_{j=1}^{\Gamma} {}^*(Fc)(t'_j) \bullet {}^*\vec{v}_j &= \sum_{j=1}^{\Gamma} \left(\sum_{n=1}^n {}^*f_i({}^*c(t'_j)) ({}^*c_i(t_{j-1}) - {}^*c_i(t_j)) \right) = \\ &= \sum_{i=1}^n \left(\sum_{j=1}^{\Gamma} {}^*f_i({}^*c(t'_j)) ({}^*c_i(t_{j-1}) - {}^*c_i(t_j)) \right). \end{aligned}$$

Hence, if for each $i = 1, \dots, n$, $\sum_{j=1}^{\Gamma} {}^*f_i({}^*c(t'_j)) ({}^*c_i(t_{j-1}) - {}^*c_i(t_j)) \in \mu(r_i)$, then $\sum_{j=1}^{\Gamma} {}^*(Fc)(t'_j) \bullet {}^*\vec{v}_j \in \mu(r_1 + \dots + r_n)$. On the other hand, if $\sum_{j=1}^{\Gamma} {}^*(Fc)(t'_j) \bullet {}^*\vec{v}_j \in \mu(r)$ and for each $i = 1, \dots, n$, $\sum_{j=1}^{\Gamma} {}^*f_i({}^*c(t'_j)) ({}^*c_i(t_{j-1}) - {}^*c_i(t_j)) \in \mathcal{O}$, then $\text{st}(\sum_{j=1}^{\Gamma} {}^*f_i({}^*c(t'_j)) ({}^*c_i(t_{j-1}) - {}^*c_i(t_j))) = r_i$ implies that $r_1 + \dots + r_n = r$. This leads to the notion of the line integral as an extension of the Riemann-Stieltjes integral.

Definition 8.2.2. (Line Integral). Let bounded $c: [a, b] \rightarrow \mathbb{R}^n$, $c(t) = (c_1(t), \dots, c_n(t))$ be considered a curve with graph C . Assume that bounded $F: C \rightarrow \mathbb{R}^n$, where $F(\vec{x}) = (f_1(\vec{x}), \dots, f_n(\vec{x}))$. Consider the composite function $(Fc): [a, b] \rightarrow \mathbb{R}^n$ defined by $(Fc)(t) = (f_1(c(t)), \dots, f_n(c(t)))$, $t \in [a, b]$. Then F is **LINE INTEGRABLE with respect to $\mathbf{C}$** if there exists some real r such that for any fine partition $P = \{a = t_0, \dots, t_\Gamma = b\}$ of $[a, b]$ and any internal intermediate partition $Q = \{t'_1, \dots, t'_\Gamma\}$ of P it follows that

$$\sum_{j=1}^{\Gamma} {}^*(Fc)(t'_j) \bullet {}^*\vec{v}_j \in \mu(r).$$

In which case we write

$$r = \text{st} \left(\sum_{j=1}^{\Gamma} {}^*(Fc)(t'_j) \bullet {}^*\vec{v}_j \right) = \int_C F \bullet d\vec{R}.$$

↓ IMPORTANT ↓

⇒ In the next derivation, the significant method of the maximum and minimum is applied to physical elements. This approach is different than that used in integral rule IR3. Moreover, a special relationship between the work done by a force field over a hyperline segment and points on a rectifiable curve is advanced. This relationship aids in our comprehension of energy related NSP-world properties. ⇐

Application 8.2.1. *Energy expended within a force field while moving along a curve.*

Let $c: [a, b] \rightarrow \mathbb{R}^n$ be a continuous differentiable curve with graph C . Assume that continuous $F: E \rightarrow \mathbb{R}^n$, open $E \supset C$. The work done in moving through the force field on the path C is

$$W(C) = \int_C F \bullet d\vec{R} = \int_a^b \left(\sum_{i=1}^n f_i(c_1(t), \dots, c_n(t)) c'_i(t) \right) dt.$$

Derivation. In experimental physics, the concept of “work” (energy expended) is introduced. **All one needs to do is to establish its properties for a polygonal curve.** Suppose we have continuous force field $F: E \rightarrow \mathbb{R}^n$. Let $\mathcal{P}_k \subset \mathbb{R}^n$ be any finite polygonal curve, ℓ_j one of the line segment portions of $\mathcal{P}_k$ with $\vec{v}_j = (c_1(t_j) - c_1(t_{j-1}), \dots, c_n(t_j) - c_n(t_{j-1}))$, denoting this line segment considered as a directed line segment in the direction of motion through the field F . If F is constant on ℓ_j , then the work done moving along ℓ_j is defined as $W(\ell_j) = F \bullet (\vec{v}_j / \|\vec{v}_j\|) \|\vec{v}_j\|$, where length of $\ell_j = \|\vec{v}_j\|$. What if the force field is not constant? Consider $\mathcal{P}_k$ as represented by a continuous $\ell: [a, b] \rightarrow \mathbb{R}^n$ and assume that F is defined on ℓ . Then for a given $\ell_j = \{(x_1(t), \dots, x_n(t)) \mid t \in [t_{j-1}, t_j]\}$ there exists some t_m, t_M such that $W_m(\ell_j) = F(\ell(t_m)) \bullet \vec{v}_j \leq W(\ell_j) = F(\ell(t)) \bullet \vec{v}_j = W_M(\ell_j) = F(\ell(t_M)) \bullet \vec{v}_j$ for each $t \in [t_{j-1}, t_j]$. Let's make the one assumption that the actual amount of work expended moving along the line segment ℓ_j is $W(\ell_j)$ and that $W_m(\ell_j) \leq W(\ell_j) \leq W_M(\ell_j)$. Then from continuity there exists some $h'_j \in [t_{j-1}, t_j]$ such that $W(\ell_j) = F(\ell(h'_j)) \bullet \vec{v}_j$. The idea of the nonconstant force field over a line segment is embedded into the NSP-world by *-transfer assuming that what has been established above holds for all such polygonal curves. Hence, let $\mathcal{P}_\Omega$ be a hyperpolygonal representation for the curve generated by a fine partition, ℓ_j an hyperline segment in $\mathcal{P}_\Omega$. Since F is continuous on E then $*F$ is defined on $\mathcal{P}_\Omega$. It follows that $*W(\ell_j) = *F(\ell_j(h'_j)) \bullet *\vec{v}_j$. For polygonal curves, in general, the work done is an additive function. Thus for the hyperpolygonal curve $\mathcal{P}_\Omega$

$$*W(\mathcal{P}_\Omega) = \sum_{j=1}^{\Omega} *F(\ell_j(h'_j)) \bullet *\vec{v}_j. \quad (1)$$

We now show that there exists a real number $W(C)$ such that for any fine partition $\mathcal{P}_\Gamma$ and any intermediate partition $Q = \{t'_1, \dots, t'_\Gamma\}$, $\mathbf{st}(*W(\mathcal{P}_\Gamma)) = W(C)$.

Consider

$$*F(*c(t'_j)) \bullet *\vec{v}_j = \sum_{i=1}^n *f_i(*c_1(t'_j), \dots, *c_n(t'_j)) (*c_i(t_j) - *c_i(t_{j-1})). \quad (2)$$

The curve c being continuously differentiable on $[a, b]$ implies that $*c_i(t_j) - *c_i(t_{j-1}) = (*c'_i(t_j) + \delta_{ij})(t_j - t_{j-1})$, where $\delta_{ij} \in \mu(0)$. Hence, $*F(*c(t'_j)) \bullet *\vec{v}_j =$

$$\sum_{i=1}^n *f_i(*c_1(t'_j), \dots, *c_n(t'_j)) *c'_i(t_j) (t_j - t_{j-1}) + \left(\sum_{i=1}^n \delta_{ij} \right) (t_j - t_{j-1}). \quad (3)$$

Consequently, $*F(*c(t'_j)) \bullet *v_j =$

$$\sum_{i=1}^n *f_i(*c_1(t'_j), \dots, *c_n(t'_j)) *c'_i(t_j) (t_j - t_{j-1}) + \delta_j (t_j - t_{j-1}). \quad (4)$$

Uniform continuity of the f_i yields that $*f_i(*c_1(t'_j), \dots, *c_n(t'_j)) = *f_i(*c_1(t_j), \dots, *c_n(t_j)) + \lambda_{ij}$, where $\lambda_{ij} \in \mu(0)$. Once again this yields $*F(*c(t'_j)) \bullet *v_j =$

$$\begin{aligned} & \sum_{i=1}^n *f_i(*c_1(t_j), \dots, *c_n(t_j)) *c'_i(t_j) (t_j - t_{j-1}) + \\ & \lambda_j (t_j - t_{j-1}) + \delta_j (t_j - t_{j-1}) \end{aligned} \quad (5)$$

Now continuing the basic **elemental derivation process** leads to $\sum_{j=1}^{\Gamma} (*F(*c(t'_j)) \bullet *v_j) =$

$$\sum_{j=1}^{\Gamma} \left(\sum_{i=1}^n *f_i(*c_1(t_j), \dots, *c_n(t_j)) *c'_i(t_j) (t_j - t_{j-1}) \right) + \delta, \quad \delta \in \mu(0). \quad (6)$$

Therefore, from Theorem 5.1.2

$$\begin{aligned} W(C) &= \mathbf{st} \left(\sum_{j=1}^{\Gamma} (*F(*c(t'_j)) \bullet *v_j) \right) = \\ & \int_a^b \left(\sum_{i=1}^n f_i(c_1(t), \dots, c_n(t)) c'_i(t) \right) dt. \end{aligned} \quad (7)$$

The fact that this is a line integral follows from Definition 8.2.1. Finally, it will almost always be the case that the special energy property $*F(\ell_j(t'_j)) \bullet *v_j = *F(*c(t'_j)) \bullet *v_j + \epsilon_j \|*v_j\|$, $\epsilon_j \in \mu(0)$, $t'_j = h'_j$ holds. [See note [2] on page 148.] Since C is rectifiable then $\sum_{j=1}^{\Gamma} *F(\ell_j(t'_j)) \bullet *v_j \approx \sum_{j=1}^{\Gamma} *F(*c(t'_j)) \bullet *v_j$. We are using the hyperpolygonal representations for C as the basic NSP-world entity to determine the N-world physical effects. It is clear that the appropriate measure for the work expended moving along the path C should be the unique value $\mathbf{st}(\sum_{j=1}^{\Gamma} *F(\ell_j(t'_j)) \bullet *v_j)$ obtained in (7). This completes the derivation.

Obviously, derivation 8.2.1 also establishes the elementary method for the calculation of a line integral.

8.3. Order Ideals and Approximations.

One often reads in the literature that such and such an expression is a “first- order approximation” or some such phrase. These vague approximation concepts can be discussed from the infinitesimal viewpoint and, indeed, lead to the notions of the “microconstruction” and “microeffects.”

Theorem 8.3.1. *For each $\epsilon \in \mu(0)$ the set $o(\epsilon) = \{\epsilon h \mid h \in \mu(0)\}$ is an ideal in $\mu(0)$.*

Definition 8.3.1. (Order Ideals). For a given $\epsilon \in \mu(0)$ the set $o(\epsilon)$ is called an **ORDER IDEAL** (of infinitesimals).

We briefly investigate some of the basic properties of the order ideals. (I note that these order ideal properties appear for the first time in this manual.) First, it is obvious that $o(\epsilon) = o(-\epsilon)$. In all that follows, let $\mu(0)^+ = \{x \mid x \in \mu(0) \wedge x \geq 0\}$ be the set of all **nonnegative infinitesimals**. The next theorem seems to be one of the more significant ones relative to order ideals.

Theorem 8.3.2. *Let $\epsilon \in \mu(0)^+$. Suppose that $w \in {}^*\mathbb{R}$ and $0 \leq w \leq \epsilon h \in o(\epsilon)$. Then $w \in o(\epsilon)$.*

Theorem 8.3.3. *Let $\epsilon, \delta \in \mu(0)^+$. If $0 \leq \delta \leq \epsilon$, then $o(\delta) \subset o(\epsilon)$ and $o(\delta)$ is a ideal in $o(\epsilon)$.*

Are there order ideals such that $o(\delta) \subset o(\epsilon)$ and $o(\delta) \neq o(\epsilon)$?

(1) Let $0 < \delta \leq \epsilon$, $\delta, \epsilon \in \mu(0)^+$. Then $o(\delta\epsilon) \subset o(\epsilon)$ and $o(\delta\epsilon) \neq o(\epsilon)$.

(2) Let $n \in \mathbb{N}^+ = \mathbb{N} - \{0\}$, $\epsilon \in \mu(0)^+$. Then $o(\epsilon^n) \subset o(\epsilon^{n-1}) \subset \dots \subset o(\epsilon)$ and $o(\epsilon^i) \neq o(\epsilon^j)$; $1 \leq i, j \leq n$; $i \neq j$.

Suppose that you have the set $\{\epsilon_1, \dots, \epsilon_n\}$. Then consider the chain $C_1 : o(\epsilon_1 \times \dots \times \epsilon_n) \subset o(\epsilon_1 \times \dots \times \epsilon_{n-1}) \subset \dots \subset o(\epsilon_1)$. The chain C_1 is just one possible chain of order ideals leading from $o(\epsilon_1 \times \dots \times \epsilon_n)$ to $o(\epsilon_1)$. Such chains are used for comparison purposes and, in this case, the ideal $o(\epsilon_1 \times \dots \times \epsilon_n)$ is called an **n'th order ideal** where, in general, such ideals as $o(\epsilon)$ are called **first-order ideals**. Part of this chapter will deal with the relation between n'th order ideals and the infinitesimal concept of n'th order approximations. It is useful to consider other operational methods that might generate different order ideals rather than simply restricting their generation to products of infinitesimals.

Let $\{\epsilon_1, \dots, \epsilon_k\} \subset \mu(0)$. Define $o(\epsilon_1, \dots, \epsilon_k) = \{\epsilon_1 h_1 + \dots + \epsilon_k h_k \mid h_i \in \mu(0) \wedge 1 \leq i \leq k\}$.

Theorem 8.3.4. *Let $\epsilon = \max \{|\epsilon_1|, \dots, |\epsilon_k|\}$. Then $o(\epsilon_1, \dots, \epsilon_k) = o(\epsilon)$.*

Theorem 8.3.5. *Let $\epsilon = \max \{|\epsilon_1|, \dots, |\epsilon_k|\}$. Then*

$$o(\epsilon_1, \dots, \epsilon_k) = o(\sqrt{\epsilon_1^2 + \dots + \epsilon_k^2}) = o(\epsilon).$$

Thus neither the difference of infinitesimals nor the *-Euclidean norm is a useful process for the generation of higher order ideals. As will be illustrated through out the remainder of this manual, the n'th order ideals are related to the notion of the n'th order approximation. Refer back to all of our derivations where the basic elemental derivation process is used. In each case, a single term of the required hyperfinite sum of elemental measures, say ${}^*M(\cdot)$, is investigated for an arbitrary simple partition of infinitesimal volume dX . The derivation shows that this term is equal to the value of a standard extension, say ${}^*F(\vec{x}_i)$, plus $h dX$, where $h \in \mu(0)$. Consequently,

$${}^*M(\cdot) - {}^*F(\vec{x}_i) \in o(dX).$$

This expression also yields an equivalence relation which is often denoted by the algebraic notation

$${}^*M(\cdot) \approx {}^*F(\vec{x}_i) \pmod{o(dX)}$$

and is expressed by stating that ${}^*M(\cdot)$ and ${}^*F(\vec{x}_i)$ are **infinitely close of order dX** . Hence, to apply this derivation process the values ${}^*M(\cdot)$ are not just infinitely close to ${}^*F(\vec{x}_i)$, in a general sense; but,

they are infinitely close of order dX . Noting that for an n -dimensional integral $dX = dx_1 \times \cdots \times x_n$ it follows that, from a comparative viewpoint, ${}^*M(\cdot)$ and ${}^*F(\vec{x}_i)$ can be considered as **infinitely close of order n** . In the literature, you will also find the less descriptive expression “ ${}^*M(\cdot)$ is infinitely close to ${}^*F(\vec{x}_i)$ compared to dX ” as a synonym for “infinitely close of order dX .” The reason it appears necessary that such quantities need to be infinitely close of order dX is that the steps in the derivation process that proceed from this step require a hyperfinite sum to be extracted and the results must remain infinitely close.

One little observation about the order ideals that are created by products. If $0 \neq r \in {}^*[-1, 1] - \mu(0)$, then for any $\epsilon \in \mu(0)$, it follows that $o(\epsilon r) = o(\epsilon)$. To see this, Theorem 8.3.3 yields that $o(|\epsilon r|) = o(\epsilon r) \subset o(|\epsilon|) = o(\epsilon)$. Let $\epsilon h \in o(\epsilon)$. Since $h/r \in \mu(0)$, then $(\epsilon r)(h/r) \in o(\epsilon r)$ implies that $\epsilon h \in o(\epsilon r)$. Further, note that if $0 \neq \delta \in \mu(0)$, then $\epsilon r \in o(\epsilon) - o(\delta \epsilon)$.

8.4. n th Order Increments.

In the next section, we investigate exactly what one means by a tangent line to a curve c at a point p , where c is differentiable at p . First, however, it is useful for this and the future sections on modeling by means of the derivative or differential to formally consider the NSP-world view of the n th order increment (nth difference). For $n \in \mathbb{N}^+$ and bounded $f: [a, nb] \rightarrow \mathbb{R}$, recall that the **n th order increment**, $\Delta^n f(x, b)$, is defined by induction, where for $x \in [a, nb]$, by $\Delta f(x, b) = f(x + b) - f(x)$. This leads to the general expression

$$\Delta^n f(x, b) = \sum_{k=0}^n (-1)^k \binom{n}{k} f(x + (n - k)b) = \sum_{k=0}^n (-1)^k \binom{n}{k} f(x + kb).$$

Notice that as an operator the * - n th order increment, ${}^*(\Delta^n) {}^*f(x, b) = \Delta^n {}^*f(x, b)$.

Theorem 8.4.1. *Let $1 \leq n \in \mathbb{N}$. Suppose that $f^{(n-1)}: [a, nb] \rightarrow \mathbb{R}$ and that $f^{(n)}: (a, nb) \rightarrow \mathbb{R}$, where $f^{(k)}$ denotes the k th derivative of f . Then there exists some $t \in (a, nb)$ such that $\Delta^n f(a, b) = f^{(n)}(t) b^n$.*

Corollary 8.4.1.1 *Let $1 \leq n \in \mathbb{N}$. Suppose that $f^{(n-1)}: [a, b] \rightarrow \mathbb{R}$ and that $f^{(n)}: (a, b) \rightarrow \mathbb{R}$, then for each $dx \in \mu(0)^+$ and $c \in {}^*[a, b)$, there exists some $t \in (c, c + ndx)$ such that $\Delta^n {}^*f(c, c + dx) = {}^*f^{(n)}(t) (dx)^n$.*

Theorem 8.4.1 holds if the hypotheses are appropriately altered to $f^{(n-1)}: [a - na, b] \rightarrow \mathbb{R}$ and that $f^{(n)}: (a - na, b) \rightarrow \mathbb{R}$. In this case, Corollary 8.4.1.1 may be altered to $dx \in \mu(0)$, $dx < 0$, $c \in {}^*(a, b]$, $t \in (c + ndx, c)$ and $\Delta^n {}^*f(c + dx, c) = {}^*f^{(n)}(t) (dx)^n$.

Theorem 8.4.2. *Let $1 \leq n \in \mathbb{N}$. Suppose that $f^{(n-1)}: [a, b] \rightarrow \mathbb{R}$ and that $f^{(n)}: (a, b) \rightarrow \mathbb{R}$. If $c \in (a, b)$, then for each $dx \in \mu(0)$, $dx \geq 0$ [resp. $dx < 0$]*

$$f^n(c) (dx)^n \approx \Delta^n {}^*f(c, c + dx), [\text{resp. } f(c + dx, c)] \pmod{o((dx)^n)}.$$

One important aspect of Theorem 8.4.2 is that the quantities $f^n(c) (dx)^n$ and $\Delta^n {}^*f(c, c + dx)$ are not simply infinitely close; but, rather, are infinitely closed of order n . For an infinitesimal, dx , Robinson and those that founded the infinitesimal calculus consider $\Delta^n {}^*f(c, c + dx)$ to be the

n th order differential of f at c . However, most other authors still retain the notion that $d^n f(c) = f^{(n)}(c) (dx)^n = f^{(n)}(c) dx^n$; which I shall retain as well. Under the hypotheses of Theorem 8.4.2 it follows for nonzero positive [resp. negative] infinitesimal, dx , that $d^n f(c)/dx^n \approx \Delta^n *f(c, c + dx)$, [resp. $f(c + dx, c)$] = $*f^{(n)}(t)$ and, hence, $\mathbf{st}(d^n f(c)/dx^n) = \mathbf{st}(\Delta^n *f(c, c + dx)$, [resp. $f(c + dx, c)$]) = $f^{(n)}(c)$.

Finally, all that has been said about n th order ideals is extended to m -dimensional objects of the form $o^n(\epsilon_1, \dots, \epsilon_m) = o(\epsilon_1) \times \dots \times o(\epsilon_m)$.

8.5. Microgeometry - Tangents to Curves.

A tangent to a curve $c: [a, b] \rightarrow \mathbb{R}^p$ is usually defined as the intuitive limit of a set of secants. How might this be viewed within the NSP-world? In order to analytically answer this question, let c be differentiable at $t \in (a, b)$. Next, let any $r \in *[-1, 1]$. Fix $\epsilon \in \mu(0)^+$ and let f_i be the coordinate functions of c . The Fundamental Theorem of differential calculus states, in infinitesimal form, that there exists some $h_i \in \mu(0)$ such that

$$*f_i(t + r\epsilon) = f_i(t) + f'_i(t)(r\epsilon) + h_i(r\epsilon). \quad (1)$$

For any $r_1 \in *[-1, 1] - \mu(0)$, equation (1) then yields, since $o(r_1\epsilon) = o(\epsilon)$

$$*c(t + r_1\epsilon) = c(t) + c'(t)(r_1\epsilon) + (h_1, \dots, h_p)(r\epsilon), \quad (2)$$

$$*c(t + r_1\epsilon) \approx c(t) + c'(t)(r_1\epsilon) \pmod{o^n(\epsilon)}. \quad (3)$$

Each component of $o^n(\epsilon)$ is a first-order ideal and what follows next is a direct result of this fact and the concept of the **resolving power of a microscope**. Expression (1) is first transformed into an external relation determined by

$$g_i(t + r\epsilon) = f_i(t) + f'_i(t)(r\epsilon), \quad (4)$$

$$\vec{g}(t + r\epsilon) = c(t) + c'(t)(r\epsilon). \quad (5)$$

Assume that we are “looking at” the geometric NSP-world situation with an infinite powered microscope with “first-order ϵ -resolving power.” Physically, this corresponds to the idea that if the distance between two objects is a member of a first-order ideal $o(\epsilon)$, then the objects cannot be resolved (i.e. cannot be distinguished one from the other.) A simple proof shows that if $r \in *[-1, 1]$, then the best we can say is that

$$\| *c(t + r\epsilon) - \vec{g}(t + r\epsilon) \| \in o(\epsilon). \quad (6)$$

Equation (6) does not mean that for selected $r\epsilon$ the value $\|c(t + r\epsilon) - \vec{g}(t + r\epsilon)\|$ may not be in an n th order ideal. Indeed, if $r = \epsilon^{n-1}$, then this would be the case. Thus (6) means that, in general, $o(\epsilon)$ is the “smallest” order ideal that can be guaranteed to contain this value in all possible cases. For an “ ϵ -infinitesimal microscope” (abbr: ϵ -IM) with first-order ϵ -resolving power, (6) implies that within the field of view

$$*c(t + r\epsilon) = \vec{g}(t + r\epsilon) = c(t) + c'(t)(r\epsilon). \quad (7)$$

A microscope is suppose to magnify, however. How is this feat accomplished. The point, $c(t)$, is being considered as the center of view of the ϵ -IM and this point is translated to the origin.

This gives us the expression ${}^*c_1(t + r\epsilon) = {}^*c(t + r\epsilon) - c(t) = c'(t)(r\epsilon)$. The magnification and resolving power of ordinary microscopes are related. Hence, for consistency, consider for any $x_j \in {}^*\mathbb{R}$ and $\delta_j \in \mu(0)$ the general **infinite magnification operator** $m(x_1\delta_1, \dots, x_p\delta_p) = (x_1, \dots, x_p)$, which is also assumed to be linear. If you wish to specify a specific shaped field of view, say a p -dimensional closed sphere, you can also restrict this magnification to $\sum_{j=1}^p x_j^2 \leq 2$. Obviously, both the translation and magnification can be combined into one operator. Letting, as before, $r \in {}^*[-1, 1]$ the final view in the ϵ -IM only shows the hyperline segment $\{m(f'_i(t)r\epsilon, \dots, f'_p(t)r\epsilon) | r \in {}^*[-1, 1]\} = \{(f'_1(t)r, \dots, f'_p(t)r) | r \in {}^*[-1, 1]\} = \{c'(t)r | r \in {}^*[-1, 1]\}$.

8.6 Microgeometry - Surface Elements.

The well-known difficulties of determining a single Euclidean configuration as an appropriate approximation for the surface of a 3-dimensional object will not be discussed in this manual. [See **Ce-sari** [1956]] Instead, I pass directly to analytical considerations with the appropriate constraints. Let open $G \subset \mathbb{R}^2$. Assume that $\vec{r}: G \rightarrow \mathbb{R}^3$ and that for $(u, v) \in G$ the continuous partial derivatives $\vec{r}_u(u, v)$, $\vec{r}_v(u, v)$ exist. Let $\vec{r}_u(u_0, v_0) = \vec{a}$, $\vec{r}_v(u_0, v_0) = \vec{b}$ and $|\vec{a} \times \vec{b}| \neq 0$. In order to be consistent with requirement IR2, define, in matrix notation on column vectors, the linear transformation $L: \mathbb{R}^3 \rightarrow \mathbb{R}^3$ as follows:

$$L = \begin{pmatrix} r_1(u_0, v_0) & a_1 & b_1 \\ r_2(u_0, v_0) & a_2 & b_2 \\ r_3(u_0, v_0) & a_3 & b_3 \end{pmatrix}. \quad (1)$$

In the previous section, a hyperline segment portion of the tangent line was viewed within an IM. The same procedures are now applied to the tangent plane to the surface generated by $\vec{r}$. Let $(s, t) \in {}^*[-1, 1] \times {}^*[-1, 1] = I$; $\delta, \epsilon \in \mu(0)^+$. Then the set $R_S = \{(u, v) \mid (u, v) = (u_0 + s\delta, v_0 + t\epsilon) \wedge (s, t) \in I\}$ is an internal infinitesimal rectangle and $R_S \subset \mu((u_0, v_0)) \in {}^*G$. The linear transformation ${}^*L$ transforms the internal set of vectors $\{(1, u - u_0, v - v_0)^T \mid (u, v) \in R_S\} = D$ onto a configuration ${}^*L(D)$ which is a * -Euclidean hyperparallelogram containing $\vec{r}(u_0, v_0)$. When considered restricted to D , this linear transformation can be expressed by

$$\vec{k}(u, v) = \vec{r}(u_0, v_0) + (u - u_0)\vec{a} + (v - v_0)\vec{b}. \quad (2)$$

Using the * -Euclidean concepts, this hyperparallelogram is a NSP-world portion of the standard tangent plane to the surface at $\vec{r}(u_0, v_0)$. Two adjacent sides of this hyperparallelogram have end points $\vec{k}(A) = \vec{r}(u_0, v_0) + (-\delta)\vec{a} + (-\epsilon)\vec{b}$; $\vec{k}(B) = \vec{r}(u_0, v_0) + (-\delta)\vec{a} + (\epsilon)\vec{b}$; $\vec{k}(C) = \vec{r}(u_0, v_0) + (\delta)\vec{a} + (-\epsilon)\vec{b}$. This yields two hyperline segments of hyperlength $|\vec{AB}| = \|2\delta\vec{a}\|$ and $|\vec{AC}| = \|2\epsilon\vec{b}\|$, with the hyperarea being $\|(4\delta\epsilon)\vec{a} \times \vec{b}\| = 4\delta\epsilon\|\vec{a} \times \vec{b}\| = \|\vec{a} \times \vec{b}\| dX$.

Now to analyze the order ideal relationship between members of this hyperparallelogram and the surface itself, assume that $\vec{r}_u$ and $\vec{r}_v$ are continuous at (u_0, v_0) . The Fundamental Theorem of Differential Calculus in infinitesimal form, implies that for any $s\delta$ and any $t\epsilon$, $(s, t) \in I$ there exists $\vec{\eta} \in \mu^3(\vec{0})$ such that

$${}^*\vec{r}(u_0 + s\delta, v_0 + t\epsilon) = \vec{r}(u_0, v_0) + s\delta\vec{a} + t\epsilon\vec{b} + \|(s\delta, t\epsilon)\| \vec{\eta}. \quad (3)$$

Considering any $(s, t) \in {}^*[-1, 1] \times {}^*[-1, 1]$ a simple proof yields that

$$\|{}^*\vec{r}(u_0 + s\delta, v_0 + t\epsilon) - {}^*\vec{k}(u_0 + s\delta, v_0 + t\epsilon)\| \in o(\max\{\delta, \epsilon\}) = o(\lambda). \quad (4)$$

The magnification operator for our (δ, ϵ) -IM is the mapping $m(x_1\delta + y_1\epsilon, \dots, x_3\delta + y_3\epsilon) = m(x_1 + y_1, \dots, x_3 + y_3)$, where the x 's and y 's are hyperreal numbers. Translating and magnifying

$*k(u_0 + s\delta, v_0 + t\epsilon)$ yields as the final view in the (δ, ϵ) -IM the hyperparallelogram $\{s\vec{a} + t\vec{b} \mid (s, t) \in I.\}$ **BUT**, these results are more significant than a simple exercise in analyzing the IM view of the tangent plane.

Thus far we have decided upon three types of geometric elements.

(A) Rectifiable curves $\Leftrightarrow$ **hyperline segments** and **hyperpolygonal curves**.

(B) For 1 - 3 dimensional geometric measures by means of the 1-dimensional integral $\Leftrightarrow$ **hypertrapezoids**.

(C) For n-dimensional integrals $\Leftrightarrow$ **infinitesimal rectangles**.

These elements are intuitive in character and only by means of a restrictive analytical description are physical quantities relative to them, such as mass, infinitesimal energy and the other applications given in this manual, actually calculable by means of the integral. However, as evident from all that has preceded, **most individuals consider the integral as but a hyperfinite sum of entities that are intuitively defined** and do not, generally, concerned themselves with the difficulties in calculation. I have been slightly restrictive in some of the basic definitions by requiring that functions that generate hyperfinite sums, at least, be bounded. Of course, in certain cases this restriction might be relaxed. Indeed, in the older literature, geometers utilized intuitive infinitesimal geometry and these notions were not expressed originally in terms of any such analytical constraints.

If $\mathcal{S}$ denotes the geometric point-set called a surface, then $^*\mathcal{S}$ is the hypersurface. If, intuitively, $T \subset \mathcal{S}$ is the set of surface points at which tangent planes $\mathcal{T}$ to $\mathcal{S}$ exist, then $^*T$ is the set of points in $^*\mathcal{S}$ at which the hypertangent planes $^*\mathcal{T}$ exist. For the surface integral the geometric element - **the surface element**, σ - is an infinitesimal parallelogram containing a point from the hypersurface. This surface element is considered to be contained in a hypertangent plane. As such this element has an infinitesimal area $d\sigma$. **From the viewpoint of infinitesimal modeling this description of the geometric surface element is adequate.** From the view point of surface integral calculation, since there are infinitely many geometric surface elements of different area, such a description is not sufficient.

The only question that remains is which collection of surface elements should be required for an analytical definition? For this elementary manual, the * -Euclidean area notion will be maintained along with a fixed set of vectors normal to the hypertangent planes. Let $E \subset \mathbb{R}^2$ and bounded $\vec{r}: E \rightarrow \mathbb{R}^3$. Suppose that $\mathcal{S} = \{\vec{r}(u, v) \mid (u, v) \in E\}$ and that for nonempty $T' \subset E$ the set of points $\vec{r}[T'] = T$ is called a set of **tangent points to the surface $\mathcal{S}$** . Further, there exists a mapping $\vec{\nu}: T \rightarrow (\mathbb{R}^3 - \{\vec{0}\})$, where each $\nu(\vec{t})$ is called a **normal vector** to the surface $\mathcal{S}$. Then each $\vec{t} \in T$ defines a unique **tangent plane** $\mathcal{T}(\vec{t}, \vec{\nu}(\vec{t})) = \{(x_1, \dots, x_3) \mid (x_1, \dots, x_3) \in \mathbb{R}^3 \wedge \vec{\nu}(\vec{t}) \bullet ((x_1, \dots, x_3) - \vec{t}) = 0\} \subset \mathbb{R}^3$. Assume that $E \subset R \subset \mathbb{R}^2$. If P' is any partition of R , then there exists a nonempty finite set of subrectangles $R'_i \subset R$ such that $T' \cap R'_i \neq \emptyset$, $1 \leq i \leq k$ and for each such R'_i there exist the intermediate partitions $Q' = \{t'_1, \dots, t'_k\}$ such that $t'_i \in R'_i \cap T'$. Hence, for any fine partition P of $^*R$ there exists a hyperfinite set of infinitesimal subrectangles $\tau(P) = \{S \mid S \cap ^*T' \neq \emptyset \wedge S \in P\}$ and a corresponding set of internal intermediate partitions $\eta(P)$. These ideas and notations are used in the next definition.

Definition 8.6.1. (Surface Integral.) Let the surface with its tangent planes be defined as in the above paragraph and assume that bounded $F: \mathcal{S} \rightarrow \mathbb{R}$. Let $E \subset R \subset \mathbb{R}^2$. Then F is said to be **SURFACE INTEGRABLE with respect to $\mathcal{S}$** if there exists some simple fine partition P of R and some $r \in \mathbb{R}$ such that for each intermediate partition $Q \in \eta(P)$

$$\sum_{S \in \tau(P), t' \in S \cap Q} {}^*F({}^*\vec{r}(t')) \| {}^*\vec{v}({}^*\vec{r}(t')) \| {}^*v(S) \in \mu(r).$$

[**Remark.** In the definition of the line integral, it is required that all fine partitions be considered. This was done so that consideration could be given to rectifiable curves that need not be continuously differentiable. It is obvious that Definition 8.6.1. is styled solely for the integral as defined by 5.1.1.]

Since it is clear from Theorem 7.2.2 and Definition 5.1.1 that for F to be integrable with respect to $\mathcal{S}$ the values ${}^*F({}^*\vec{r}(t')) \| {}^*\vec{v}({}^*\vec{r}(t')) \|$ must be obtained from an integrable function defined on an appropriate Jordan-measurable $J \subset R \subset \mathbb{R}^2$, I see no need to state the various well-known functions that lead to this conclusion - with one exception. Let compact $J \subset R \subset \mathbb{R}^2$ and bounded $\vec{r}: J \rightarrow \mathbb{R}^3$ generate a surface $\mathcal{S}$. How do we generalize the concept of Euclidean area to such a surface?

Suppose that $\vec{r}(u, v) = \vec{p} + u\vec{a} + v\vec{b}$, where $\vec{p}, \vec{a} \neq \vec{0}, \vec{b} \neq \vec{0}$ are fixed members of $\mathbb{R}^3$ and $(u, v) \in [a, b] \times [c, d] = J$. Then $\vec{r}[J]$ is a parallelogram with area equal to $(d - c)(b - a) \|\vec{a} \times \vec{b}\|$. But, $\vec{r}_u = \vec{a}$ and $\vec{r}_v = \vec{b}$. Assume that $K = \text{int}(J) - W$, where W is a set of Jordan-content zero. Then taking other simple Euclidean surfaces generated by corresponding simple sets of surface equations leads to the conclusion that it should be required for functions $\vec{r}: J \rightarrow \mathbb{R}^3$ that $\vec{r}_u(u, v)$ and $\vec{r}_v(u, v)$ exist and $\|\vec{r}_u(u, v) \times \vec{r}_v(u, v)\| \neq 0$ for $(u, v) \in K$, if area preserving surface integration is desired. This is precisely the starting point for elementary analysis. Notice that the linear transformations that correspond to $\mathbb{R}^2$ are pointwise defined and each is considered to generate only the specific hyperparallelogram associated with a specific term in the hyperfinite sums used in Definition 8.6.1. I leave it to the reader to combine the elemental method of constants on surface elements with Definition 8.6.1 to obtain the concept of flux of a vector field across a surface.

8.7 Microgeometry - Other Stuff.

What about the geometry of the coordinate transformations? In general, it appears necessary to consider stronger constraints than for surface integration. If a map $f: R \rightarrow \mathbb{R}^m$, where $R \subset \mathbb{R}^m$, is to be considered a coordinate transformation, then it needs to be considered locally infinitely close to a linear transformation determined by a nonzero Jacobian. What this signifies is that each m -dimensional infinitesimal subrectangle is mapped onto an infinitesimal parallelepiped - a hyperparallelepiped. Thus the infinitesimally partitioned space is distorted into a space of hyperparallelepipeds. All the details as to why this appears necessary can be found in **Stroyan and Luxemburg** beginning in section 5.8 [110].

In the above mentioned section of the book by Stroyan and Luxemburg, you will also find in section 5.8 the infinitesimal approach to orientated partitions via differential forms. In section 5.9, the authors investigate the infinitesimal calculus on manifolds. In all of these sections, the authors extend the integral concept to internal functions that usually preserve monads. Thus their integral concept, although it is the same as has been presented here for nonstandard extensions of standard functions, is actually defined on a wider class of objects.

8.8 Gauge Integrals

Recently, additional emphases has been given to a generalization of the Riemann sum called the **gauge integral** [Swartz and Thomson [1988]]. This integral concept is also called the Riemann-complete integral, [Henstock [1961]]; or S-integral, [Mawhin [1985]]; among other names. The gauge integral can also be generalized to the Jarnik, Kurzweil, Schwabik [1983] integral (the **M-integral**, Mawhin [1985].) Indeed, a Riemann sum styled generalization leads to the classical Lebesgue integral [McShane [1973]]. From the infinitesimal viewpoint, all of these generalizations are very similar.

For the S and M-integrals, a positive real valued function, δ , called a **gauge** is utilized to select not only a special partition but a special intermediate partition as well.

Within the NSP-world a set of gauges determines an internal mapping $\hat{\delta}$ from ${}^*R \subset {}^*\mathbb{R}^m$ into $\mu(0)$ called a **microguage**. However, associated with every gauge, δ , is an object $(Q_\delta, P_\delta) = \{(t_1, S_1), \dots, (t_k, S_k)\}$ where $P_\delta = \{S_1, \dots, S_k\}$ is a subdivision of R by nonoverlapping subsets and $Q_\delta = \{t_1, \dots, t_k\}$ is a partial sequence where each $t_i \in R$. Such an object (Q_δ, P_δ) called a **δ -fine partition**. Thus for a microguage $\hat{\delta}$ there exists a hyperfinite subdivision $P_{\hat{\delta}} = \{S_1, \dots, S_\Gamma\}$ of ${}^*R$ and an hyperfinite intermediate partition $Q_{\hat{\delta}} = \{t_1, \dots, t_\Gamma\}$. The pair $(Q_{\hat{\delta}}, P_{\hat{\delta}})$ is called a **micropartition**. I point out that each member of $P_{\hat{\delta}}$ is an infinitesimal subrectangle in this case. For a function $f: R \rightarrow \mathbb{R}^k$, Mawhin [1986] shows that f is *S-integrable* if and only if there exists an $\vec{r} \in \mathbb{R}^k$ such that for each micropartition $(Q_{\hat{\delta}}, P_{\hat{\delta}})$ of ${}^*R$

$$\sum_{i=1}^{\Gamma} {}^*f(t_i) {}^*v(S_i) \in \mu(\vec{r}).$$

Mawhin gives a completely similar characterization for the M-integral but replaces the general micropartition with the more specialized regular micropartition.

Looking at another type gauge, which I shall call the **L-gauge**, λ , and a corresponding partition pair (Q_λ, P_λ) , where it is not assumed that if $t_i \in Q_\lambda$, then $t_i \in S_i \in P_\lambda$, McShane [1973] defines his integral by the same Riemann sum technique. The **McShane integral** is equivalent to the Lebesgue integral. Using the same technique employed by Mawhin, the following is established in appendix 8.

Theorem 8.8.1. *A bounded function $f: R \rightarrow \mathbb{R}$ is Lebesgue integrable with value $r \in \mathbb{R}$ if and only if for each L-micropartition $(Q_{\hat{\lambda}}, P_{\hat{\lambda}})$ of ${}^*R$*

$$\sum_{i=1}^{\Gamma} {}^*f(t_i) {}^*v(S_i) \in \mu(r).$$

Thus from the infinitesimal viewpoint there is no basic difference between any of these integral concepts discussed in this section. **They are all hyperfinite sums, where each term is the scalar product of a limited object (the value of a bounded function) by the volume of an infinitesimal subrectangle.**

Chapter 9.

THE BEGINNINGS OF DIFFERENTIAL EQUATION MODELING

9.1. Brief Discussion.

In the previous sections, we presented the standard rules, nonstandard rules and the highly successful method of elements in order to establish integral models. The strengths of the derivations depended first, upon the acceptance of an appropriate correspondence between the terms within a specific discipline and objects within the mathematical structure, and then upon three specific methods. The presence of the Infinite Sum Theorem and the two Self-Evident Theorems led to what appear to be strong derivation arguments. As far as the very popular “method of elements” is concerned, for the geometric case the nonuniqueness of the chosen elements introduced a definite weakness into the method. For the physical case, the point charge method of application 7.2.5 and the elements of energy used in application 8.2.1 may not be considered particularly realistic in character. Interestingly enough, however, the **elemental method of constants** which allows function evaluation at “convenient points” is often an easily accepted facet of infinitesimal modeling. The reason for this is that experience has shown that, for the physical applications, quantities that are infinitely close within the NSP-world produce the same natural world effects.

Differential equation models are somewhat more difficult to justify for numerous reasons. Primarily, this is due to the often vague “rules of correspondence” which must be introduced. These rules, stated in a secondary metalanguage, create a correspondence, Φ , between a list of terms or relations selected from a discipline theory and a list of terms or relations selected from a mathematical structure. For differential equation modeling, the relation Φ is defined in a more piecemeal fashion than is the case for integral modeling. Whereas in integral modeling, the hyperfinite sum concept, the Self-Evident Theorems, and the like require that Φ maps certain terms and corresponding relations to specific mathematical objects, this is not the process dictated for differential equation modeling.

Within a specific discipline what constitutes simplistic or idealized behavior is not just determined by nonmathematical experience and intuition. Using a back-and-forth correspondence technique, specific simplistic behavior that approximates the more complex is selected in concert with an available mathematical structure. This simplistic behavior is then imposed, by intuitive *-transfer, upon an infinitesimal portion of the NSP-world.

The historical development of the differential calculus does not follow the customary patterns incorporated into our modern textbooks. Indeed, the originators of this subject relied heavily upon visualization and the experiential aspects of objective reality, not only for their basic modeling techniques, but also for their methods of algebraic manipulation. The scholastic training of the 1600’s emphasized certain acceptable approaches that did not clash with philosophical dictum. Their methods became controversial when abstractions were made that led away from what some considered to be perceivable reality.

What are the intuitive notions that the originators of the infinitesimal concept attempted to model and can we learn the foundations of such modeling from their writings? First, it was assumed that geometry was the servant of natural philosophy. In the 1686 preface of the first edition of Newton’s “Mathematical Principles of Natural Philosophy” he writes “*Therefore geometry is founded in mechanical practice, and is nothing but that part of universal mechanics which accurately proposes and demonstrates the art of measuring.*” [Newton [1686]] Berkeley tells us that “*Lines are supposed*

to be generated by the motion of points, planes by the motion of lines, and solids by the motion of plans.” [Berkeley [1734]] Consequently, during the period of its formation the infinitesimal calculus and its application to natural philosophy was based upon point-motion that produced geometric configurations. Newton took the concept of the locus, made additional observations and, with respect to natural philosophy, claimed that such motion was caused by “forces.” These forces coupled with experiential evidence lead to other motions that produced other geometric configurations that can predict other motions and observable patterns. Thus physical observation of moving objects came first; followed by the concept of motion; which leads to geometry; which, coupled with force relations, leads to other geometric patterns; which were interpreted as paths of motion. Finally, these paths of motion predicted the location of observable moving objects. As is seen in the next section, it is the coalescing of motion, distance traveled and absolute time that produced the infinitesimal calculus and is the fundamental methodology that leads to geometric differential modeling.

9.2. The Limit.

For elementary limit theory all that one needs is a simple alteration of Definition 4.4.1. For $\vec{p} \in \mathbb{R}^n$, let the **deleted monad** be $\mu'(\vec{p}) = \mu(\vec{p}) - \{\vec{p}\}$.

Definition 9.1.1. (Limit of a Function). For any nonzero $n, m \in \mathbb{N}$ and any nonempty $A \subset \mathbb{R}^n$ a function $f: A \rightarrow \mathbb{R}^m$ has the **LIMIT** = $\vec{L}$ as $\vec{x} \rightarrow \vec{p} \in \mathbb{R}^n$ if $\mu'(\vec{p}) \cap *A \neq \emptyset$ and $*f[\mu'(\vec{p}) \cap *A] \subset \mu(\vec{L})$.

The following theorem is obtained in the same manner as the corresponding one for continuity.

Theorem 9.1.1. *Definition 9.1.1 for the limit of a function is equivalent to the classical $\delta - \epsilon$ definition.*

The basic properties of the limit operator follow from those of the standard part operator. And, for this reason, the next theorem, which follows from Chapter 10, section 1, corresponds for extended standard functions to the classical definition of the derivative.

Theorem 9.1.2. *Suppose that nonempty open $A \subset \mathbb{R}^n$ and $\vec{a} \in A$. Then $f: A \rightarrow \mathbb{R}^m$ is differentiable at $\vec{a}$ if and only if there exists a unique linear mapping $T_a: \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that for each $\vec{\epsilon} \in \mu(\vec{0})$,*

$$*f(\vec{a} + \vec{\epsilon}) - f(\vec{a}) = *T_a(\vec{\epsilon}) + (\|\vec{\epsilon}\|)\lambda(\vec{\epsilon}),$$

where $\lambda(\vec{\epsilon}) \in \mu(\vec{0})$.

9.3. Fluxions and Dynamic Geometry.

In the middle 1600’s Newton utilized a purely dynamic method to arrive at his geometry. He introduced a new type of dynamics that for some natural philosophers was not related to the material world and would need to be rejected. I will not go into this controversy in this manual. However, Newton’s dynamical (loci) methods, restated from the view point of Robinson’s theory, may be a better approach to the foundations of infinitesimal modeling than presently presented

within the standard calculus course. Unfortunately, some of Newton's actual demonstrations of the more complex geometric concepts, such as curvature, are not valid from the viewpoint of Robinson's theory and require slight alterations. When these alterations are conjoined with Robinson's theory, then elementary demonstrations are easily obtained and comprehended.

In Newton's paper [Summer 1665], an algorithm is given that yields the relations between the "fluxions" p , q associated with the variables x , y . These variables are related by an algebraic expression that is assumed to generated a geometric configuration. In Newton [Oct. 1665 - May 1666: 383] the algorithm is specifically described. How Newton, by observation, arrived at this algorithm and what exactly p , q represent is discussed later in his paper. A better explanation of how he formulated his algorithm and the meaning of the term fluxion appears in his Oct. 1666 tract [Newton, 1666]. Under proposition 7 [Newton 1666:402], he explains his algorithm, step by step. After some examples, he discusses how he arrived at this algorithm and what fluxions signify [Newton 1666:414]. First, he considers two "bodies A , B moving uniformly." He lets an algebraic expression $f(x, y) = 0$ represent a relation between the distance traveled by these two bodies. Then Newton introduces the concept of the distance traveled by a body having uniform velocity p , usually, over a "moment" of "infinitely small" time o . Newton represents the distance each body travels by the sum of line segment lengths. Body A first travels along $\overline{ac}$ and at the same time body B travels along $\overline{bg}$. Now in an "infinitely small" period of time, o , body A travels along $\overline{cd}$ and during the same time interval body B travels along the segment $\overline{gh}$. He states that the motion is not, in general uniform, but it is "*as if the body A with its velocity p describe the infinitely little line $\overline{cd} = p \times o$ in one moment, in that moment the body B with the velocity q will describe the line $\overline{gh} = q \times o$. So that if the described lines be $\overline{ac} = x$, and $\overline{bg} = y$, in one moment, they will be $\overline{ad} = x + po$, and $\overline{bh} = y + qo$ in the next.*"

Newton claims that the fluxions p and q are a type of velocity (what type?) and he proceeds to demonstrate how relations between these fluxions, in particular the relations relative to the quotient q/p , are obtained. It's within this demonstration that contradictions occur. Newton writes "*Now if the equation expressing the relation between the lines x and y be $x^3 - abx + a^3 - dy^2 = 0$. I may substitute $x + po$ and $y + qo$ into the place of x and y ; because (by the above) they as well as x and y do signify the lines described by the bodies A and B . [Of course, this statement would only be true if fluxions or the motion of the bodies is uniform over a standard time interval, o , and the ordinary Galilean physics is applied.] By doing so there results*

$$\begin{aligned} & x^3 + 3pox^2 + 3p^2o^2x + p^3o^3 - dy^2 - 2dpoy - dq^2o^2 \\ & \quad - abx - abpo \\ & \quad + a^3 = 0. \end{aligned} \tag{1}$$

But $x^3 - abx + a^3 - dy^2 = 0$ (by supp). Therefore there remains only

$$3pox^2 + 3p^2o^2x + p^3o^3 - 2dqoy - dq^2o^2 - abpo = 0. \tag{2}$$

Or dividing it by o it is

$$3px^2 + 3p^2ox + p^3o^2 - 2dqy - dq^2o - abp = 0. \tag{3}$$

[Thus for the algebraic processes of the 1600's o is a nonzero real number. Newton goes on to write:] *Also those terms are infinitely little in which o is. Therefore omitting them there results*

$$3px^2 - abp - 2dqy = 0. \tag{4}$$

The like may be done in all other equations. Newton would then continue and express his important ratio

$$\frac{q}{p} = \frac{3x^2 - ab}{2dy}. \quad (5)$$

Obviously step (4) is not justified and to some, such as Berkeley, contradicts the nature of the infinitely small nonzero quantity o . After this, and in other demonstrations, Newton indicates that he guessed at portions of his fluxion creating algorithm by applying this steps (1) - (5) to numerous algebraic expressions and making certain observations as to the physical appearance of such equations (4) and (5). This algorithm, it is claimed, simply eliminated the need to apply continually the above, often criticized, process. Newton repeats similar derivations in his Winter 1670 - 1671 tract as well as suggesting that the delineated process may be applied to relations between three or more variables.

Assuming that the paths of two objects can be so linearly displayed, then modern infinitesimal analysis seems to reveal Newton's intentions and correct his vague logic. First, note that Newton assumes without justification that the q and p that appear in equations (1) - (3) are the same as those that appear in (4). Since this should not be so assumed then in all cases suppose that when the process arrives at an expression such as (4) that the q is replaced by Q and the p by P . Newton seems to describe his notion of the "infinitely small" in his Scholium following Lemma XI of Book one of his 1686 treatise. In particular, he discusses the "ultimate velocities" Q and P and his ratios Q/P that he calls "ultimate ratios." *Perhaps it may be objected, that there is no ultimate proportion of evanescent quantities; because the proportion, before the quantities have vanished, is not ultimate, and when they are vanished is none. But by the same argument it may be alleged that a body arriving at a certain place, and there stopping, has no ultimate velocity; because the velocity, before the body comes to the place, is not its ultimate velocity; when it has arrived, there is none. But the answer is easy; for by the ultimate velocity is meant that with which the body is moved, neither before it arrives at its last place and the motion ceases, nor after, but at the very instant it arrives; that is, the velocity with which the body arrives at its last place, and with which the motion ceases. And in like manner, by the ultimate ratio of evanescent quantities is to be understood the ratio of the quantities not before they vanish, nor afterwards, but with which they vanish....For those ultimate ratios with which quantities vanish are not truly the ratios of ultimate quantities, but limits towards which the ratios of quantities decreasing without limit do always converge; and to which they approach nearer than by any given difference, but never go beyond, nor in effect attain to, till the quantities are diminished in infinitum.* What has been described by Newton is very close to the classical limit notion. However, from his applications and arguments this does not seem to be what Newton truly believed but only a popular exposition that would not offend the geometers of his day. As is well-known Newton was very fearful of criticism and, even though he would use his fluxion methods in private to model physical behavior, he did not perform fluxion computations directly within this all important research document.

In equations (1) - (3) suppose that the o is a standard real number and that it is claimed that (3) holds in $\mathcal{M}$ for any such o that is an "infinitely small" real number. Berkeley [Berkeley [1734]] indicates that the Leibniz school would include quantities that are "smaller than" those real numbers previously termed as infinitely small. By *-transfer of Berkeley's description it follows that expressions such as (3) would hold for any infinitesimal ϵ . Further, Newton's notion of motion and velocities can be embedded into an appropriate ${}^*\mathbb{R}^n$. All other objects that appear in such an equation as (3) are considered fixed standard constants or relations in o . In order to determine what the quantities p or q are measuring, let's simplify our investigation and assume that the relationship

between the distances traveled by body A and B is implicitly functional. Indeed, let $f: [a, b] \rightarrow \mathbb{R}$ and, as with Newton, denote $y = f(x)$, where fixed $x \in (a, b)$. Replicating Newton's process down to step (3) obtains

$$q = \frac{f(x + p o) - f(x)}{o}. \quad (6)$$

Equation (6) gives a relation between q and p and the **arbitrary** infinitely small real number o . Newton requires q and p to be some type of velocity (fluent) and it is not unreasonable that whatever kind of velocity these quantities might be measuring that they be bounded over his set of infinitely small times. Further, from the above descriptions the “infinitely small”, we suppose that if o_1 is infinitely small, then any quantity o such that $|o| < |o_1|$ is also infinitely small. Thus even if we consider o as a real number, then $*$ -transfer yields

$$*q = \frac{*f(x + *p \epsilon) - f(x)}{\epsilon}, \quad (7)$$

where $\epsilon \in \mu(0)$.

If $*p = 0$, then $*q = 0$. Suppose, therefore, that $*p \neq 0$. Then

$$*q = *p \left(\frac{f(x + *p \epsilon) - f(x)}{*p \epsilon} \right). \quad (8)$$

Assuming that f is differentiable at x then application of definition 9.1.1 implies that

$$Q = \text{st}(*q) = \text{st}(*p) \text{st} \left(\frac{f(x + *p \epsilon) - f(x)}{*p \epsilon} \right) = P f'(x). \quad (9)$$

Can we be certain that Newton's ratio q/p may be formally written as $Q/P = f'$? In his 1666 tract, Newton claims to obtain the fundamental ratio q/p by first substituting for x the expression $x + o$ and letting $y = y + (q/p) o$, where o is an arbitrary infinitely small real number. Developing the above formal derivation leads to

$$\frac{q}{p} = \frac{f(x + o) - f(x)}{o}. \quad (10)$$

$$*\left(\frac{q}{p}\right) = \frac{*f(x + \epsilon) - f(x)}{\epsilon}, \quad (11)$$

$$\text{st}*\left(\frac{q}{p}\right) = \text{st} \left(\frac{*f(x + \epsilon) - f(x)}{\epsilon} \right) = f'(x) = \frac{Q}{P}. \quad (12)$$

As Newton did in his applications, where q and p were characterized merely as velocities, he would also arrive at an expression for q only, by assuming that since q and p are introduced as arbitrary parameters then one could simply let $p = 1$. The final result he would obtain, prior to omitting the higher order o 's, would be as expressed by equation (10). Thus his two methods are not quite equivalent; but, rather, equation (12) should probably be expressed as

$$\text{st}(*q) = \text{st} \left(\frac{*f(x + \epsilon) - f(x)}{\epsilon} \right) = f'(x) = Q. \quad (13)$$

Equation (13) is significant for, at least, two reasons in his applications. First, for *unit constant* velocity p , the velocity measured by q is the *average velocity* and as such q would behave in the

infinitesimal world in accordance Galilean physics, where infinitesimal time ϵ is considered as a parameter. Secondly, Newton often used the expression

$$\frac{q}{p} = \frac{f(x+o) - f(x)}{o}, p = 1, \quad (14)$$

to obtain what we now know to be the second derivation of f at x . This he did by writing $z = q$ and then expressing the relation between the velocity r (in our notation not his) of z as

$$z + Ro = \frac{f((x+o)+o) - f(x+o)}{o}. \quad (15)$$

This obviously leads to

$$r = \frac{f(x+2o) - 2f(x+o) + f(x)}{o^2} \Rightarrow \quad (16)$$

$$R = f^{(2)}(x), \quad (17)$$

assuming that the second derivative of f exists at x and applying Theorem 8.4.2. In his writings Newton used his famous “dot” notations which, unfortunately, do not correlate formally to their meaning in terms of our modern concepts. For example, in the above displayed equations he would often write $p = \dot{x}$, $q = \dot{y}$ and $r = \dot{z}$. This is not equal to the dot notation employed in elementary calculus, where $\dot{x} = \text{st}(*q)$.

9.4. Fluxions and Higher Order Infinitesimals.

I have often wondered while reading physical arguments that purport to derive a particular expression relating measurable quantities what rules, if any, govern the elimination or omission of “higher order” increments. Is it only experience or a deep seated intuition that leads to the assumption that the omitted quantities will not, indeed, essentially influence the outcome or are their other unexpressed factors that have entered into such derivations?

The actual method used by Newton to establish the majority of the physical principles discussed in his Mathematical Principles of Natural Philosophy [Newton [1686]] is the theory of fluxions and a specialized process of omitting higher order increments. Newton would expand expressions into finite or infinite series intending to remove eventually all expressions involving his infinitely small o through a division process. This required him to omit various expressions involving products with o . However, whether or not he omitted expressions involving the higher order terms (i.e. terms containing the factors o^n , where $n \geq 2$), or not depended upon the final proposition being sought.

Technically, the process was not fixed; but, was often altered in such a way that by omitting certain, but not necessarily all, higher order terms from his expressions the final results could be obtained through a division process that eliminated all of the remaining infinitely small factors. Moreover, if, due to such technical uncertainties, one attempted derivation failed to verify a result, then many others were tried. This method of derivation is, indeed, vague and forced. But, there is a basic philosophy being expressed by such forced procedures. This modeling philosophy is closely resembles the concept of physical completeness for a mathematical model. *One believes that the descriptive physical content of a theory is absolutely correct. Then it is believed that the standard mathematical structure correlates exactly to the physical theory. Thus if parameters are introduced, such as infinitely small numbers, and any “reasonable” method is used to eliminate these parameters, then, since the final expressions involve only terms that correlate to the original physical theory, the final expressions “must” give an accurate model for physical reality.* This is the same reason

why the original creators of differential equation models were convinced that their equations had solutions, since in their philosophy the equation obtained must predict physical behavior and the physical behavior does seem to occur.

Obviously, Newton's methods were very successful. The time parameter that Newton introduced into his dynamic geometry need not, of course, be an actually parameter. Thus when some of the infinitely small time increments could not be removed by a division process the final expressions could retain the increments expressed in terms of o with the understanding that the result was approximate - the approximation being relative to the "smallest" of o . The Newtonian methods of arbitrarily eliminating higher order increments continues today. Our concern, however, is with rigorous differential equation modeling where such arbitrariness may be partially eliminated.

9.5. What is a tangent?

Our modern analogue for the fluxion concept is captured by the vector notion. Today, core mathematics courses utilize the theory of free vectors at an early stage in the calculus curriculum. The interpretation of such vectors as represented by directed line segments "attached" to points in $\mathbb{R}^n$ yields the associated geometry. In all that follows, it is assumed that all vectors are limited (i.e. members of $\mathcal{O}^n$.) Referring back to Example 4.4.1A or the first page of Appendix 6, a continuous $c: [0, 1] \rightarrow \mathbb{R}^n$ is called a curve and is assumed also to be generated by n continuous coordinate functions $f_i: [0, 1] \rightarrow \mathbb{R}$, where for each $t \in [0, 1]$; $c(t) = (f_1(t), \dots, f_n(t))$. In order to visualize the geometry, one may assume that $n = 3$ even though all of the results in this section hold for any positive $n \in \mathbb{N}$.

From our NSP-world analytical viewpoint, a curve is represented entirely by a collection of hyperpolygonal curves. Can we formulate a geometric definition for a possible tangent vector to the curve c at some $c(t)$, where $t \in (0, 1)$ which appropriately generalizes the Euclidean concept? Obviously, requiring a tangent line to be unique would eliminate immediately the basic plane geometry definition that such a line would intersect the curve in at most one point. Since, however, the properties of the curve should be entailed by the collection of hyperpolygonal representations $\{\mathcal{P}_\Gamma\}$, then a seemingly natural definition for the existence of a tangent line might involve the collection of all hyperline segments that have standard $c(t)$ as one end point and constitutes one of the hyperline segments (the **elements**) in some member of $\{\mathcal{P}_\Gamma\}$. More specifically, this means that a hypervector that would represent such a hyperline segment would be of the form $*c(t + dx) - c(t)$, where dx is an nonzero infinitesimal. Intuitively, the tangent line and these hyperline segments should be infinitesimally near to each other. This corresponds to the requirement that the measure of the plane angle between a tangent vector and each of these hyperline segments be infinitesimal or infinitesimal close to π . The next definition models this easily grasped geometric picture since the cosine of the measure of such an angle is infinitely close to ± 1 .

Definition 9.5.1. (Tangent Vector to a Curve). Let continuous $c: [0, 1] \rightarrow \mathbb{R}^n$, and $t \in (0, 1)$. A standard unit vector $\vec{T}$ is a **TANGENT** at $c(t)$ if for each $dx \in \mu'(0)$,

$$\vec{T} \bullet \frac{*c(t + dx) - c(t)}{\|*c(t + dx) - c(t)\|} \approx \pm 1.$$

Definition 9.5.1 is meaningless if there exists some $dx \in \mu'(0)$ such that $\|*c(t + dx) - c(t)\| = 0$. Now writing Theorem 9.1.2 in the form $(*c(t + dx) - c(t))/|dx| =$

$(^*T_t)(dx/|dx|) + \lambda(dx)$ it is obvious that if c is differentiable at t and $c'(t) \neq 0$, then $^*c(t+dx) - c(t) \neq 0$ for any $dx \in \mu'(0)$. Definition 9.5.1 points directly to derivative of c as being a primary candidate for tangent consideration, as is well-known. In some textbooks on this subject a tangent vector is simply defined as the derivative. This approach does not show, conclusively, how the tangent vector corresponds to either the geometric notions or to the elemental method. In this section applications to geometry are being stressed. For this reason, the following theorems are established in their entirety within this section rather than hiding them in an appendix. The dot product, $\bullet$, is the Euclidean inner product.

Theorem 9.5.1. *Let continuous $c: [0, 1] \rightarrow \mathbb{R}^n$ be differentiable at $t \in (0, 1)$ and $c'(t) \neq \vec{0}$. Then $c'(t)/\|c'(t)\|$ is a unit tangent vector at $c(t)$.*

Proof. From the above discussion we know that $^*c(t+dx) - c(t) \neq 0$ for any $dx \in \mu'(0)$. Referring back to the basic elemental derivation process of example 4.4.1A in appendix 6 or as used in application 8.2.1 it follows that for each $dx \in \mu'(0)$, $\| ^*c(t+dx) - c(t) \| = \|c'(t)\| |dx| + |dx|\delta$, where $\delta \in \mu(0)$. Thus let $dx \in \mu'(0)$. Then

$$\frac{c'(t)}{\|c'(t)\|} \bullet \frac{^*c(t+dx) - c(t)}{\| ^*c(t+dx) - c(t) \|} = \frac{c'(t)}{\|c'(t)\|} \bullet \frac{\frac{^*c(t+dx) - c(t)}{|dx|}}{\|c'(t)\| + \delta} = \quad (1)$$

$$\frac{c'(t)}{\|c'(t)\|} \bullet \frac{(\pm c'(t)) + \lambda}{\|c'(t)\| + \delta} = w, \quad (2)$$

where $\lambda \in \mu(\vec{0})$. Distributing obtains

$$w = \frac{\pm c'(t) \bullet c'(t) + c'(t) \bullet \lambda}{\|c'(t)\|(\|c'(t)\| + \delta)} \approx \quad (3)$$

$$\pm \frac{c'(t) \bullet c'(t)}{\|c'(t)\|(\|c'(t)\| + \delta)} \approx \pm \frac{c'(t) \bullet c'(t)}{\|c'(t)\|\|c'(t)\|} = \pm 1. \quad (4)$$

Theorem 9.5.2. *Let continuous $c: [0, 1] \rightarrow \mathbb{R}^n$ be differentiable at $t \in (0, 1)$ and $c'(t) \neq \vec{0}$. If $\vec{T}$ is a unit tangent vector at $c(t)$, then $\vec{T} = \pm(c'(t)/\|c'(t)\|)$.*

Proof. Let $\vec{T} = (a_1, \dots, a_n)$ and $dx \in \mu'(0)$. Then

$$1 \approx \left| \vec{T} \bullet \left(\frac{^*c(t+dx) - c(t)}{\| ^*c(t+dx) - c(t) \|} \right) \right| \approx \left| \vec{T} \bullet \left(\frac{c'(t)}{\|c'(t)\|} \right) \right| \quad (5)$$

Thus

$$\frac{|\sum_{i=1}^n a_i f'_i(t)|}{\sqrt{\sum_{i=1}^n f'_i(t)^2}} = 1 \Rightarrow \quad (6)$$

$$\left| \sum_{i=1}^n a_i f'_i(t) \right| = \sqrt{\sum_{i=1}^n f'_i(t)^2} = \left(\sqrt{\sum_{i=1}^n a_i^2} \right) \left(\sqrt{\sum_{i=1}^n f'_i(t)^2} \right). \quad (7)$$

Since neither vector is a zero vector then the Schwarz Inequality states that there exists some nonzero d such that $\vec{T} = d(c'(t)/\|c'(t)\|)$. Thus $\|\vec{T}\| = |d| \|c'(t)/\|c'(t)\|\| \Rightarrow |d| = 1$ and the proof is complete.

9.6. What is an (k-surface) Osculating Plane?

One of the early concepts used by proponents of differential geometry is the notion of “consecutive” points in a curve or surface. Modern textbooks, such as **Struik [1961]**, emphasize this intuitive approach, even to the point that some of their derivations may be considered less than fully justified. The definitions for those significant geometric objects associated with the tangents are often given without adequate discussion as to the geometric content of the definition and why various constraints are necessary. Modern infinitesimal analysis eliminates this vagueness.

Before extending our results to the k -surface case we look at the 3-dimensional prototype. Let $t_0, t_1, t_2 \in (0, 1)$ and assume that the vectors $\vec{c}_1 = c(t_1) - c(t_0)$, $\vec{c}_2 = c(t_2) - c(t_0)$ are linearly independent. The unique plane containing the two line segment representations for $\vec{c}_1$, $\vec{c}_2$ attached to $c(t_0)$ is $\Pi(c(t_0), \vec{c}_1, \vec{c}_2) = \{\vec{x} \mid \vec{x} \in \mathbb{R}^3 \wedge (\vec{x} = c(t_0) + \lambda \vec{c}_1 + \nu \vec{c}_2) \wedge (\lambda, \nu \in \mathbb{R})\}$. Elementary linear algebra may also be used to find $\Pi(c(t_0), \vec{c}_1, \vec{c}_2)$. Consider the matrix

$$A = A(\vec{x}, c(t_0), \vec{c}_1, \vec{c}_2) = \begin{pmatrix} \vec{x} - c(t_0) \\ \vec{c}_1 \\ \vec{c}_2 \end{pmatrix}. \quad (1)$$

Let nonzero $\vec{b}$ be a member of the 1-dimensional null space of $A(c(t_0), c(t_0), \vec{c}_1, \vec{c}_2)$. Then $\Pi(c(t_0), \vec{c}_1, \vec{c}_2) = \Pi(\vec{b}, c(t_0)) = \{\vec{x} \mid \vec{x} \in \mathbb{R}^3 \wedge (\vec{x} - c(t_0)) \bullet \vec{b} = 0\} = \{\vec{x} \mid \vec{x} \in \mathbb{R}^3 \wedge \det(A) = 0\}$. Once $\vec{b}$ is selected then the fact that $c(t_0), c(t_1), c(t_2)$ are members of this plane can be characterized by the equation

$$f(t) = (c(t) \bullet \vec{b}) - (c(t_0) \bullet \vec{b}). \quad (2)$$

Now let us assume that the second derivation $c^{(2)}$ is continuous on $(0, 1)$ and that $t_0 < t_1 < t_2$. Since, in this case, $f(t_0) = f(t_1) = f(t_2) = 0$ and $[t_0, t_2] \subset (0, 1)$, then Rolle’s theorem tells us that there exists v_1, v_2 such that $t_0 \leq v_1 \leq t_1 \leq v_2 \leq t_2$ and such that

$$\begin{aligned} f^{(1)}(v_1) &= c^{(1)}(v_1) \bullet \vec{b} = 0 \\ f^{(1)}(v_2) &= c^{(1)}(v_2) \bullet \vec{b} = 0. \end{aligned} \quad (3)$$

Moreover, there exists some v_3 such that $v_1 \leq v_3 \leq v_2$ and

$$f^{(2)}(v_3) = c^{(2)}(v_3) \bullet \vec{b} = 0. \quad (4)$$

Embedding the above remarks into the NSP-world then it follows that for three $t_0, t_1, t_2 \in {}^*(0, 1)$ and $t_0 \approx t_1 \approx t_2 \approx t' \in (0, 1)$ (ordered as previously) such that ${}^*c(t_1) - {}^*c(t_0) = {}^*c_1$, ${}^*c(t_2) - {}^*c(t_0) = {}^*c_2$ are linearly independent in ${}^*\mathbb{R}^3$ and for any nonzero $\vec{b} \in \mathcal{O}^3$ in the 1-dimensional null space for ${}^*A({}^*c(t_0), {}^*c(t_0), {}^*c_1, {}^*c_2)$ there exist v_1, v_3 such that $v_1 \approx v_3 \approx t_0 \approx t_1 \approx t_3$ and

$$({}^*c(t_0) \bullet \vec{b}) - ({}^*c(t_0) \bullet \vec{b}) = {}^*c^{(1)}(v_1) \bullet \vec{b} = {}^*c^{(2)}(v_3) \bullet \vec{b} = 0. \quad (5)$$

But, we also know that $\{v_1, v_3, t_0, t_1, t_2\} \subset \mu(t')$. Continuity of the derivatives yields that

$$c^{(1)}(t') \bullet \vec{b} \approx 0, \quad c^{(2)}(t') \bullet \vec{b} \approx 0. \quad (6)$$

and thus

$$c^{(1)}(t') \bullet \mathbf{st}(\vec{b}) = 0, \quad c^{(2)}(t') \bullet \mathbf{st}(\vec{b}) = 0. \quad (7)$$

Now what would happen if $c^{(1)}(t'), c^{(2)}(t')$ are linearly independent? Let $B = A(\vec{x}, c(t'), c^{(1)}(t'), c^{(2)}(t'))$ and consider the plane $\Pi(\text{st}(\vec{b}), c(t'))$. Obviously, if $\vec{b}$ is considered a limited normal to the object ${}^*\Pi(\vec{b}, {}^*c(t_0))$, then we have normals to the two planes that are infinitely close. More significantly, however, is the fact that these planes do satisfy an “infinitely close” type property. Extending the above discussion to n -dimensional spaces we have

Theorem 9.6.1.

- (i) Let $c: [0, 1] \rightarrow \mathbb{R}^n$, $n > 1$ and assume that for some k such that $1 \leq k \leq n-1$ the k -derivative, $c^{(k)}$, is continuous on $(0, 1)$.
- (ii) Let $\{t_0, \dots, t_k\} \subset {}^*(0, 1)$ have the properties that $\{ {}^*c_i \mid {}^*c_i = {}^*c(t_i) - {}^*c(t_0) \wedge 1 \leq i \leq k \}$ is linearly independent in ${}^*\mathbb{R}^n$ and $t_i \approx t_j \approx t' \in (0, 1)$, $0 \leq i, j \leq k$.
- (iii) Let $\{c^{(1)}(t'), \dots, c^{(k)}(t')\}$ be linearly independent.

Then for every limited $\vec{a} \in \Pi_1 = \{ \vec{x} \mid \vec{x} \in {}^*\mathbb{R}^n \wedge (\vec{x} = {}^*c(t_0) + \lambda_1 {}^*c_1 + \dots + \lambda_k {}^*c_k) \wedge (\lambda_1, \dots, \lambda_k \in {}^*\mathbb{R}) \}$ there exists $\vec{c} \in \Pi = \{ \vec{x} \mid \vec{x} \in \mathbb{R}^n \wedge (\vec{x} = c(t') + \lambda_1 c^{(1)}(t') + \dots + \lambda_k c^{(k)}(t')) \wedge (\lambda_1, \dots, \lambda_k \in \mathbb{R}) \}$ such that $\vec{a} \approx \vec{c}$.

Proof. As in the illustrated case, consider the matrix

$$A(\vec{x}, c(t_0), \vec{c}_1, \dots, \vec{c}_k) = \begin{pmatrix} \vec{x} - c(t_0) \\ \vec{c}_1 \\ \vdots \\ \vec{c}_k \end{pmatrix} \quad (9)$$

Let W be the k -dimensional subset of $\mathbb{R}^n$ spanned by $\{\vec{c}_1, \dots, \vec{c}_k\}$ and $W^\perp$ the $(n-k)$ -dimensional orthogonal complement with respect to the Euclidean inner product. Let $\{\vec{b}_1, \dots, \vec{b}_{n-k}\}$ be an orthonormal basis for $W^\perp$. Now let B be the $n \times (n-k)$ matrix formed by considering the vectors $\vec{b}_i$, $1 \leq i \leq n-k$ as column vectors. Then consider the (affine) k -plane $\Pi(B, c(t_0)) = \{ \vec{x} \mid \vec{x} \in \mathbb{R}^n \wedge (\vec{x} - c(t_0)) \cdot B = 0 \} = \{ \vec{x} \mid \vec{x} \in \mathbb{R}^n \wedge \exists z (z \in W \wedge \vec{x} = z + c(t_0)) \} = \{ \vec{x} \mid \vec{x} \in \mathbb{R}^n \wedge (\vec{x} = c(t_0) + \lambda_1 \vec{c}_1 + \dots + \lambda_k \vec{c}_k) \wedge (\lambda_1, \dots, \lambda_k \in \mathbb{R}) \}$, where $t_0 \in (0, 1)$. Let

$$f_i(t) = (c(t) - c(t_0)) \bullet \vec{b}_i, \quad 1 \leq i \leq n-k, \quad (10)$$

where $t \in (0, 1)$. Applying the same argument as in the above illustration we obtain the hyperplane $\Pi_1 = {}^*\Pi(B, {}^*c(t_0))$, where B is an $n \times (n-k)$ matrix having limited column vectors. Notice that the standard part of B is the standard part of each column vector and since the column vectors form an * -orthonormal set in ${}^*\mathbb{R}^n$ then the set of standard parts forms an orthonormal, hence independent, set of vectors in $\mathbb{R}^n$. Now equations (7) becomes

$$c^{(1)}(t') \cdot \text{st}(B) = 0, \dots, c^{(k)}(t') \cdot \text{st}(B) = 0. \quad (11)$$

Hence, $\Pi = \{ \vec{x} \mid \vec{x} \in \mathbb{R}^n \wedge (\vec{x} - c(t')) \cdot \text{st}(B) = 0 \}$. Consequently, if limited $\vec{a} \in \Pi_1$, then $(\vec{a} - {}^*c(t_0)) \cdot B = 0 \Rightarrow (\text{st}(\vec{a}) - \text{st}({}^*c(t_0))) \cdot \text{st}(B) = (\text{st}(\vec{a}) - c(t')) \cdot \text{st}(B) = 0 \Rightarrow \vec{c} = \text{st}(\vec{a}) \in \Pi$ and that $\vec{a} \approx \vec{c}$.

Any set of $(k+1)$ points that satisfies part (ii) of Theorem 9.6.1 is called a set of $(k+1)$ -consecutive points in the curve c . If all of the hypotheses of Theorem 9.6.1. are satisfied, then

Π is called the **osculating k -plane** and this object is infinitely close, in the above sense, to any of the other objects Π_1 . Observe that if $k = 1$, then Π is but that tangent line to c at $c(t')$.

9.7. Curvature.

Continue to assume that for the curve $c: [0, 1] \rightarrow \mathbb{R}^n$ the k -derivative, $c^{(k)}$, is continuous on $(0, 1)$ and for our discussion let $k = 2$ ($n > 2$), even though the following results also appear to hold for $k = 3$. As is customary for technical purposes, consider c re-expressed in terms of the arc length parameter, s . Therefore, assume that $c: [0, L] \rightarrow \mathbb{R}^n$ has continuous k -derivatives on $(0, L)$, where L is the length of the arc. For $s' \in (0, L)$, let $\{dc(s')/ds, d^2c(s')/ds^2\}$ be linearly independent.

Since $dc(s')/ds \neq 0$ then Theorem 9.5.1 implies that $dc(s')/ds = T$ is a unit tangent vector at $c(s')$. It is an easy exercise to show that $dT(s')/ds = d^2c(s')/ds^2$ is orthogonal to T and is a linear combination of $\{c^{(1)}(t'), c^{(2)}(t')\}$ for the corresponding $t' \in (0, 1)$. Thus, $dT(s')/ds$ is a member of the osculating k -plane at $c(t')$ and for this useful reason $dT(s')/ds$ is selected as a (principle) normal vector to the curve in that it is orthogonal to T . Normalizing, we obtain $\vec{k} = \kappa \vec{N}$, where $\vec{N}$ is a **unit normal vector to the curve**. What might κ represent, where we have that $\kappa > 0$?

As in section 9.6, but in terms of the arc length parameter, let $s_0, s_1, s_2 \in (0, L)$ and assume that the vectors $\vec{c}_1 = c(s_1) - c(s_0)$, $\vec{c}_2 = c(s_2) - c(s_0)$ are linearly independent. The unique k -plane containing the two line segment representations for $\vec{c}_1$, $\vec{c}_2$ attached to $c(s_0)$ is $\Pi(c(s_0), \vec{c}_1, \vec{c}_2) = \{\vec{x} \mid \vec{x} \in \mathbb{R}^n \wedge (\vec{x} = c(s_0) + \lambda \vec{c}_1 + \nu \vec{c}_2) \wedge (\lambda, \nu \in \mathbb{R})\}$. Consider a k -sphere of radius r with center $\vec{c} \in \Pi(c(s_0), \vec{c}_1, \vec{c}_2)$. A general equation for such a k -sphere may be written as

$$f(s) = (\vec{x} - \vec{c}) \bullet (\vec{x} - \vec{c}) - r^2 = 0, \quad (1)$$

where we have assumed, as with equation (2) in section 9.6, that (1) is a function in s and that it is satisfied by the points s_0, s_1, s_2 . Following the exact same process that led from equation (2) through equation (5) of section 9.6 yields, for $s_0, s_1, s_2 \in {}^*(0, L)$, $s_0 \approx s_1 \approx s_2 \approx s' \in (0, L)$, $\vec{a} \in \Pi_1$ and $r_1 \in {}^*\mathbb{R}$, the hyperreal numbers $v_1 \approx v_3 \approx s'$ such that

$$\begin{aligned} ({}^*c(s_0) - \vec{a}) \bullet ({}^*c(s_0) - \vec{a}) - r_1^2 &= 0 \\ ({}^*c(s_0) - \vec{a}) \bullet {}^*(dc/ds)(v_1) &= 0 \\ ({}^*c(s_0) - \vec{a}) \bullet {}^*(d^2c/ds^2)(v_3) + {}^*(dc/ds)(v_3) \bullet {}^*(dc/ds)(v_3) &= 0. \end{aligned} \quad (2)$$

The third equation in (2) implies that $\vec{a} \in \mathcal{O}^n$. The first equation shows that $r_1 \in \mathcal{O}$ if and only if $\vec{a} \in \mathcal{O}^n$. The standard part operator obtains

$$\begin{aligned} (c(s') - \text{st}(\vec{a})) \bullet (c(s') - \text{st}(\vec{a})) - (\text{st}(r_1))^2 &= 0 \\ (c(s') - \text{st}(\vec{a})) \bullet (dc/ds)(s') &= 0 \\ (c(s') - \text{st}(\vec{a})) \bullet (d^2c/ds^2)(s') + (dc/ds)(s') \bullet (dc/ds)(s') &= \\ (c(s') - \text{st}(\vec{a})) \bullet (d^2c/ds^2)(s') + 1 &= 0. \end{aligned} \quad (3)$$

Now let $\vec{c} = \text{st}(\vec{a})$ and $r = \text{st}(r_1)$. By Theorem 9.6.1, $\vec{c}$ is a member of the osculating k -plane and thus $c(s') - \vec{c} = \lambda(dc(s')/ds) + \nu(d^2c(s')/ds^2)$. Employing equations (3) we find that $\lambda = 0$ and $-1 = \nu(d^2c(s')/ds^2) \bullet d^2c(s')/ds^2 = \nu\kappa^2$. Consequently,

$$c(s') - \vec{c} = (1/\kappa)\vec{N}, \quad (4)$$

and the radius of the k -sphere or the k -circle formed by the intersection of the osculating k -plane and this k -sphere is $1/\kappa$. As is well-known this k -circle is called the **osculating k -circle** and gives a measure to the curvature of c at $c(s')$.

Summarizing these concepts from the NSP-world view point, we have hypercircles in the hyperplane Π_1 of Theorem 9.6.1 determined by the three consecutive points ${}^*c(s_0)$, ${}^*c(s_1)$, ${}^*c(s_2)$ that are infinitely close to $c(s')$ with the center $\vec{a} \in \Pi_1$ and radius $r_1 \in {}^*\mathbb{R}$. The continuity of the various derivatives involved implies that the center of any such hypercircle is limited and infinitely close to the center of a unique circle in Π with coordinates $c = c(s') + (1/\kappa)\vec{N}$ and having radius $1/\kappa$. Notice that if $\kappa < 0$ the same result holds except that the radius of this unique circle is $|1/\kappa|$. The 2-dimensional case is obtained by considering the 3-dimensional case but restricting the curve to a coordinate plane. Originally, the concept of the curvature, as we know it today, was not the approach used by Newton. His derivation was for the radius of curvature in the plane. He obtained the correct expression but his proof was filled with serious logical contradictions. Intuitively, differential geometers describe the osculating circle as the unique circle containing $c(s')$ in the osculating plane and “containing three consecutive points” from the curve. The nonstandard approach has eliminated these vague concepts with their often forced derivations and replaces them with comprehensible and rigorous ideas. It would be significant to undergraduate education if the entire subject matter of elementary differential geometry, with its wide area of physical application, was rewritten and presented in this more easily visualized NSP-world approach. Once the basic notions are mastered, then the intuition developed is an indispensable asset when they are extended to differentiable manifolds.

Chapter 10.

**THE DIFFERENTIAL
AND PHYSICAL MODELING**

10.1. Basic Properties.

There is one principle that is utilized continually in order to pass from the monadic environment to the standard world without implication of the standard part operator. The principle is called the **Cauchy Principle** and is the critical step required to establish the major results in this section.

Theorem 10.1.1. (Cauchy Principle) *Let $P(x)$ be a bounded formal first-order expression in one free variable and employing internal constants (i.e. a bounded formal internal property as discussed in Appendix 4, within the proof of Theorem 4.2.2.) If $P(\epsilon)$ holds for each $\epsilon \in \mu^n(\vec{0})$, then there exists an open ball B about $\vec{0}$ such that $P(\vec{x})$ holds for each $\vec{x} \in {}^*B$.*

In the last section, it was assumed that we had established the concept of the differential formally. Obviously, this was not done and the situation needs to be rectified prior to our brief applications to physical problems. First, a slight notational change is beneficial. Let $\vec{v} \in \mathbb{R}^n$. Then to prevent confusion in this chapter, we denote $\mu(\vec{v})$ by $\mu^n(\vec{v})$. I point out that what follows can be easily extended to normed linear spaces over the real or complex numbers.

Definition 10.1.1 (Local Map). Let $A \subset {}^*\mathbb{R}^n$ and internal $f: A \rightarrow {}^*\mathbb{R}^m$. Then f is a **LOCAL MAP** if $f[\mu^n(\vec{0})] \subset \mu^m(\vec{0})$. This means that f **preserves infinitesimals**.

Definition 10.1.2. (Equivalent Maps). Let f and g be local maps. If for each nonzero $\vec{x} \in \mu^n(\vec{0})$

$$\frac{f(\vec{x}) - g(\vec{x})}{\|\vec{x}\|} \in \mu^m(\vec{0}),$$

then we write $f \sim g$ and say that the two maps are **EQUIVALENT**.

Theorem 10.1.2. *Two internal maps f and g are equivalent if and only if there is a local map α such that for each nonzero $\vec{x} \in \mu^n(\vec{0})$*

$$f(\vec{x}) = g(\vec{x}) + (\alpha(\vec{x}))\|\vec{x}\|.$$

Notice that since $\mu^n(\vec{0})$ and $\mu^m(\vec{0})$ are external sets and the domain, D_f , and range, $f[D_f]$, of an internal map, f , are internal sets then $\mu^n(\vec{0})$ and $\mu^m(\vec{0})$ are, respectively, proper subsets of D_f and $f[D_f]$. Probably the simplest type of local map that one can associate with a linear space would be one that preserves linearity.

Definition 10.1.3. (Locally Linear) A local map f is said to be **LOCALLY LINEAR** (i) if for each $\vec{x}, \vec{y} \in \mu^n(\vec{0})$, then $f(\vec{x} + \vec{y}) = f(\vec{x}) + f(\vec{y})$ and (ii) for each $r \in \mathcal{O}$ and each $\vec{x} \in \mu^n(\vec{0})$ it follows that $f(r\vec{x}) = rf(\vec{x})$.

The interesting thing about locally linear maps is that even though they are restricted to linearity on infinitesimals they are uniquely associated with an internal linear map defined on the entire space ${}^*\mathbb{R}^n$.

Theorem 10.1.3. *Suppose that f is any locally linear map. Then there exists a unique internal linear map $T_f: {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}^m$ such that $\|T\| \in \mathcal{O}$ and there exists an open $E \subset \mathbb{R}^n$, where $\vec{0} \in E$, such that $f(\vec{x}) = T_f(\vec{x})$ for each $\vec{x} \in {}^*E$.*

For the present, we are not particularly concerned with the domain nor range of a locally linear map only that it satisfies the infinitesimal requirements. What the next few theorems indicate may be slightly surprising.

Theorem 10.1.4. *Let f, g be locally linear maps. Then $f \sim g$ if and only if $T_f(\vec{x}) \approx T_g(\vec{x})$ for each $\vec{x} \in {}^*\mathbb{R}^n$ such that $\|\vec{x}\| = 1$.*

Previously, the differential was defined for $f: [a, b] \rightarrow \mathbb{R}$ at a point $c \in (a, b)$ where f' exists to be $f'(c) dx$ for each $dx \in \mu(0)$. This can be viewed as the restriction of the nonstandard extension of the standard function $D: [a, b] \rightarrow \mathbb{R}$ defined by $D(x) = f'(c)x$.

Definition 10.1.4 (Differential). A local linear map f is a **DIFFERENTIAL** if there exists a (standard) bounded linear transformation $F: \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that ${}^*F(\vec{x}) = f(\vec{x})$ for each $\vec{x} \in \mu^n(\vec{0})$.

Theorem 10.1.5. *If f is a differential, then there exists a unique bounded linear transformation $F: \mathbb{R}^n \rightarrow \mathbb{R}^m$ and an open set $E \subset \mathbb{R}^n$ such that $\vec{0} \in E$ and ${}^*F(\vec{x}) = f(\vec{x})$ for all $\vec{x} \in {}^*E$ and, in particular, $F(\vec{x}) = f(\vec{x})$ for each $\vec{x} \in E$.*

What Theorem 10.1.5 does is to lift the internal differential f up to the status of a standard linear transformation on some standard nonempty set E . This may be a slight surprise since it is only assumed that this is the case for the infinitesimals. For applications, the behavior of a local map with respect to the infinitesimals is of paramount importance. Further, differentials are essentially unique with respect to equivalence.

Theorem 10.1.6. *If f and g are differentials and $f \sim g$, then there exists some open $E \subset \mathbb{R}^n$ such that $\vec{0} \in E$ and $f(\vec{x}) = g(\vec{x})$ for each $\vec{x} \in {}^*E$.*

Corollary 10.1.6.1. *If f and g are differentials, $f \sim g$, and F, G are, respectively, the corresponding (standard) linear transformation representations for f, g , then $F = G$.*

We now establish the basic relation between differentials and the derivative.

Definition 10.1.5. (Local Increment). Let nonempty open $G \subset \mathbb{R}^n$, and $\vec{c} \in G$. For any $f: G \rightarrow \mathbb{R}^m$ the **LOCAL INCREMENT** is the map $\Delta^*f_{\vec{c}}$ defined on $\mu^n(\vec{0})$ by letting

$$\Delta^*f_{\vec{c}}(\vec{b}) = {}^*f(\vec{c} + \vec{b}) - f(\vec{c}),$$

where $\vec{b} \in \mu^n(\vec{0})$.

Theorem 10.1.7. Let nonempty open $G \subset \mathbb{R}^n$, and $\vec{c} \in G$. A function $f: G \rightarrow \mathbb{R}^m$ is continuous at $c \in G$ if and only if the local increment map $\Delta^*f_{\vec{c}}$ is a local map.

Theorems 10.1.5 and 10.1.6 show that there is a standard open set E about $\vec{0}$ such that a differential is unique on E up to equivalence and is exactly equal to some linear transformation restricted to E . However, the behavior of a differential was only characterized with respect to infinitesimals. These results coupled with Theorem 10.1.7 suggest the follow definition.

Definition 10.1.6. (Differentiable). Let nonempty open $G \subset \mathbb{R}^n$, and $\vec{c} \in G$. A function $f: G \rightarrow \mathbb{R}^m$ is **DIFFERENTIABLE AT $\vec{c}$** if there exists a differential, denoted by $d_{\vec{c}}f$, that is equivalent to $\Delta^*f_{\vec{c}}$.

Obviously, from the above results if $d_{\vec{c}}f$ exists then it is representable by a linear transformation L on some standard open neighborhood of $\vec{0}$. To complete our basic results, all that is needed is to determine the appearance of an $m \times n$ -matrix representation A for such a transformation. However, this is immediate from the fact that if f , of Definition 10.1.5, is differentiable at c , then for each nonzero $\vec{b} \in \mu^n(\vec{0})$

$$\Delta^*f_{\vec{c}}(\vec{b}) = {}^*A(\vec{b}) + (\alpha(\vec{b}))\|\vec{b}\|. \quad (1)$$

Writing this in coordinate function and column vector form yields

$$({}^*f_1(\vec{c} + \vec{b}) - {}^*f_1(\vec{c}), \dots, {}^*f_m(\vec{c} + \vec{b}) - {}^*f_m(\vec{c}))^T = ({}^*A)\vec{b}^T + (\alpha(\vec{b}))\|\vec{b}\|, \quad (2)$$

where nonzero $\vec{b} \in \mu^n(\vec{0})$. Now considering any

$$\vec{b} = \overbrace{(0, \dots, \epsilon, \dots, 0)}^j, \quad \epsilon \in \mu'(0), \quad (3)$$

then ${}^*f_i(\vec{c} + \vec{b}) - {}^*f_i(\vec{c}) = a_i\epsilon + (\alpha(\vec{b}))|\epsilon|$ implies that $a_i \approx ({}^*f_i(\vec{c} + \vec{b}) - {}^*f_i(\vec{c}))/\epsilon$. Thus Definition 9.1.1 and the fact that each a_i is standard yields $a_i = \partial f_i(\vec{c})/\partial x_j$. The elementary propositions about differentiable functions follow readily from Definition 10.1.6 and equation (1).

With respect to applications equation (1) or

$$\Delta^*f_{\vec{c}}(\vec{b}) \approx {}^*A(\vec{b}) \quad (4)$$

is the most useful. Further, $\Delta^*f_{\vec{c}}(\vec{b})$ and ${}^*A(\vec{b})$ are related by first-order ideals in the sense that

$$\|\Delta^*f_{\vec{c}}(\vec{b}) - {}^*A(\vec{b})\| = \|\alpha(\vec{b})\| \|\vec{b}\|. \quad (5)$$

implies that

$$\|\Delta *f_{\vec{c}}(\vec{b}) - *A(\vec{b})\| \in o(\|\alpha(\vec{b})\|) = o(\epsilon), \quad \epsilon \in \mu(0), \quad (6)$$

from Theorem 8.3.5.

As far as the notion of the n-dimensional derivative is concerned, simply consider equation (1) written for each nonzero $\vec{b} \in \mu^n(\vec{0})$ as

$$\frac{\Delta *f_{\vec{c}}(\vec{b})}{\|\vec{b}\|} = *A\left(\frac{\vec{b}}{\|\vec{b}\|}\right) + \alpha(\vec{b}). \quad (7)$$

For the directional derivative, assuming that f is differentiable at $\vec{c} \in G$, start with a fixed standard unit vector $\hat{u}$, let $\epsilon \in \mu'(0)$ and write (1) as

$$\Delta *f_{\vec{c}}(\epsilon\hat{u}) = *A(\epsilon\hat{u}) + (\alpha(\epsilon\hat{u}))\|\epsilon\hat{u}\|. \quad (8)$$

This leads to

$$\Delta *f_{\vec{c}}(\epsilon\hat{u}) = *A(\epsilon\hat{u}) + (\alpha(\epsilon\hat{u}))|\epsilon|; \quad (9)$$

which may be written in the more familiar form

$$\frac{\Delta *f_{\vec{c}}(\epsilon\hat{u})}{\epsilon} = *A(\hat{u}) \pm \alpha(\epsilon\hat{u}) \Rightarrow \quad (10)$$

$$\text{st}\left(\frac{\Delta *f_{\vec{c}}(\epsilon\hat{u})}{\epsilon}\right) = A(\hat{u}). \quad (11)$$

Note that if (8) holds for all $\hat{u} \in {}^*\mathbb{R}^n$ and $\epsilon \in \mu(0)$, then (1) holds.

There is one other useful concept relative to applications and vectors. Two nonzero $\vec{v}, \vec{w} \in {}^*\mathbb{R}^n$ are said to be **almost parallel** if $\vec{u} = \vec{v}/\|\vec{v}\| \approx \pm \vec{w}/\|\vec{w}\|$. In an appropriate infinitesimal microscope, two almost parallel vectors appear to be parallel and, for applications, yield the same effects as do parallel vectors. The hypotheses of the next theorem are often realized in applied problems.

Theorem 10.1.8. *Let noninfinitesimal $\vec{a} \in \mathcal{O}^n$ and suppose that $\vec{a} - \vec{b} \in \mu^n(\vec{0})$. Then $\vec{a}$ is almost parallel to $\vec{b}$.*

10.2. Some General Observations

In this basic manual, it is only possible to give a few cursory illustrations of how to apply the derivative to physical problems since there are but a few general procedures that can be followed. Moreover, most such applications require specific knowledge relative to the refined behavior patterns one associates with the development of a natural system. It is more appropriate to concentrate upon the modeling of such phenomena within the confines of future more specific manuals rather than in this introductory one.

It is instructive to return to the thoughts of Newton. When Newton modeled his concept of “instantaneous velocity” he allowed the scalar velocity v to behavior over “infinitely small” time periods as if it was constant. Thus, if $x(t)$ represents the scalar distance and the derivative exists at t , then for each $\epsilon \in \mu'(0)$ it would follow from Newton’s viewpoint that $x(t+\epsilon) = x(t) + v\epsilon$. Or that $v = (x(t+\epsilon) - x(t))/\epsilon$ - the Galilean average velocity. Therefore, $\text{st}(v) = \text{st}(x(t+\epsilon) - x(t))/\epsilon = x'(t) = v$. But, it also holds that if $v \approx (x(t+\epsilon) - x(t))/\epsilon$, then $v(t) = x'(t)$. The same analysis applies to the partial derivative, the directional derivative and expression (2) of section 10.1 since

$*A = A$, the standard part operator distributes over matrix multiplication, and each such $\vec{b}/\|\vec{b}\|$ is limited. Also, observe that if the velocity is considered to be a vector quantity, $\vec{v}$, that is nonzero and continuous at t , then $\vec{v}(t + \epsilon)$ is almost parallel to $\vec{v}(t)$. **Almost parallel vectors have the same behavior in the NSP-world as parallel vectors have in the N-world.** For our basic illustrations, our functions are restricted to maps from $\mathbb{R}^n$ into $\mathbb{R}$.

(i) One of the primary principles employed when determining simplistic behavior within the monadic environment is to consider, in the natural world, how constant rates of change (i.e. linearly varying quantities) or other constant quantities are related, and then to pass such relations over to the infinitesimal world. When this is properly done one may discover that the end result is simply the replacing of $=$ with $\approx$, parallel vector properties replaced by almost parallel, etc.

(ii) The **general** physical laws for such simplistic behavior may be passed over directly to the NSP-world by $*$ -transfer. In certain cases, $*$ -transfer will not be sufficient since the basic NSP-world law may need to be written in terms of external notation.

Step (ii) may seem slightly vague; but, this is necessary since we cannot experiment within the NSP-world monadic environment and determine what the actual relation might be. This knowledge can only be indirectly obtained after the differential equation is derived, solved and used as a predictor of natural system behavior.

(iii) Due to the operational restrictions of infinitesimals and limited numbers to ring properties, it is often necessary to represent a transferred natural law in a manner different from the customary form.

In the next section, an attempt is made to use these three general observations to derive the partial differential equation for a vibrating membrane, with the obvious extension, and the equation for 3-dimensional heat transfer. The reader can determine whether or not these derivations are more rigorous and more convincing than those that appear in our present day texts.

10.3. Vibrating Membrane

Elementary nonrelativistic dynamical problems are closely associated with Newton's second law of motion. Little concern is given to whether or not this law is expressed by such an equation as $\vec{F} = m \vec{a}$ or by $\vec{F}/m = \vec{a}$. However, when this law is passed over to the monadic environment it may be necessary, due to the ring nature of the infinitesimals and limited numbers, to be more selective as to the specific expression utilized. [This is discussed more fully in Section 10.5.]

Definition 10.3.1. (Monadic Second Law of Motion) Let internal $\vec{v}: A \rightarrow *\mathbb{R}^m$, where $A \subset *\mathbb{R}^{n+1}$, $(\vec{v}(x_1, \dots, x_n, t) = \vec{v}(t))$, denote an internal function representing an internal velocity vector. Consider another internal $\vec{F}: A \rightarrow *\mathbb{R}^m$, $(\vec{F}(x_1, \dots, x_n, t) = \vec{F}(t))$, $t' \in *\mathbb{R}$, $dt \in \mu'(0)$, $0 \neq m \in *\mathbb{R}$, and $t' + dt \in A$. Then

$$\frac{\vec{F}(t')}{m} \approx \frac{\vec{v}(t' + dt) - \vec{v}(t')}{dt}. \quad (1)$$

The usual constraints placed upon a vibrating membrane such as (i) it is perfectly elastic, (ii) it is attached along its entire boundary to a plane, (iii) has a “very small” deflection (compared to its size) and only in a direction perpendicular to the plane, and (iv) it is vibrating about an equilibrium position, can be successfully idealized in the following manner.

At a moment of time, let A, B, C, D be four points in the interior of the membrane such that the configuration A, B, C, D forms a rectangle. Select the Cartesian coordinate system such that

$\overline{AD}$ is parallel to the x -axis, $\overline{AB}$ is parallel to the y -axis and the rectangle lies in the xy -plane. The line segments $\overline{AB}$, $\overline{BC}$, $\overline{DC}$, $\overline{AD}$ are contained in secants to membrane surface curves. The tensions that yield membrane motion are measured in terms of forces per unit length and separated into four nonzero constant vectors, $\vec{F}_1$, $\vec{F}_2$, $\vec{F}_3$, $\vec{F}_4$. These tensions are assumed to be exterior in nature, $\vec{F}_1$ is applied to $\overline{AD}$ in the $-z$ direction, $\vec{F}_2$ is applied to $\overline{BC}$ in the z direction, $\vec{F}_3$ is applied to $\overline{AB}$ in the $-z$ direction, and $\vec{F}_4$ is applied to $\overline{DC}$ in the z direction. Since perfect nontearing vibration occurs then two forces have been selected to counteract the effect of the remaining two.

Deformable Body Rule

One of the basic concepts used to model infinitesimally the behavior of a deformable body is that in the monadic environment its physical elements exhibit the same dynamic behavior as that of a rigid body.

The scalar force F that will accelerate the parallelogram in the $\hat{k}$ direction, when it is above the equilibrium position, is

$$F = -\|\vec{F}_1\|\|\overline{AD}\| + \|\vec{F}_2\|\|\overline{AD}\| - \|\vec{F}_3\|\|\overline{AB}\| + \|\vec{F}_4\|\|\overline{AB}\|. \quad (2)$$

⇓ IMPORTANT ⇓

⇒ The following derivations of some significant partial differential equation models for natural system behavior are stated in a protracted mode. Within these derivations, I have presented certain general modeling concepts, gleaned from indirect evidence associated with simplistic NSP-world behavior. The physical behavior of the NSP-world relative to each particular N-world event must be individually investigated and is usually characterized as the intuitive *-transfer of the simplest N-world behavior. In classical applications to continuum models, this simplistic behavior is often conceived of as intuitively discontinuous and piecemeal in character. When this is the case, the entire complex N-world effect is but a filtered composition of an infinite replication of what occurs within a single monadic environment. It is intuitively disjoint in character since the composition can be characterized as a “jumping” from one monad to another rather than some type of continuous joining of the effects. On the other hand, as recently shown [Herrmann [1989]], if we assume that certain N-world behavior is fractal or even finitely discontinuous in nature, then the NSP-world behavior that would yield such effects may be viewed as ultrasmooth and ultracontinuous in character. A representation for this ultrasmooth NSP-world behavior would necessary need to satisfy a differential equation expressed in terms of the hyperreals. However, by considering a nonstandard model of a nonstandard model it might be possible to show that such NSP-world differential equation models are once again a composition of simplistic NSP-world behavior but on a much deeper level.⇐

Application 10.3.1. A differential equation model for a vibrating membrane.

All functions are assumed to be defined on an open neighborhood, G , of (a, b, c, t_0) where, it appears to be necessary, to assume that $G \cap \mathbb{R}^3$ contains the closed membrane and, at the least, the functions are continuous where defined. Let $\vec{T} = \vec{T}(x, y, z)$ represent the internal restoring tension

per unit length. Let $z = z(x, y, t)$ represent the configuration of the membrane, as a surface, and assume that $\partial^2 z / \partial t^2$, $\partial(\|\vec{T}\|(\partial z / \partial x)) / \partial x$, $\partial(\|\vec{T}\|(\partial z / \partial y)) / \partial y$ exist and that z_x , z_y are continuous at (a, b, t_0) . Let $\rho = \rho(x, y, t)$ represent the mass per unit area. Finally, let $h = h(x, y, z, t)$ represent the per unit area lumped load function that yields the $\hat{k}$ direction external forces that may be applied to the membrane. For any such $(x, y, z, t) = (a, b, c, t_0)$, a partial differential equation model for the vibrating membrane configuration z is

$$(\rho(x, y)) \left(\frac{\partial^2 z}{\partial t^2} \right) = \frac{\partial(\|\vec{T}\|(\partial z / \partial x))}{\partial x} + \frac{\partial(\|\vec{T}\|(\partial z / \partial y))}{\partial y} + h(x, y, z, t). \quad (3)$$

Derivation. We now infinitesimalize the behavior of an infinitesimal portion of the membrane assuming that it will have the same effect in the N-world as an infinitesimal vibrating parallelogram. The key to the basic modeling technique lies in modeling intuitive statements (iii) and (iv). **(A) Expressions that involve the physical notion that something is “small” compared with something else tend to be the basis for differential equation modeling.** First, since (iv) states that this “small” deflection (i.e. the value of z at (a, b, t_0)) takes place at equilibrium then this should entail that the tension that produces such a deflection is also “small” in the $\hat{k}$ direction. Now select the plane of attachment for the membrane as the xy -plane and let the plane represent the equilibrium position. By physical observation, as the deflection is made “smaller” and “smaller” then the $ABCD$ rectangle discussed above would have its normal more nearly parallel to $\hat{k}$. The concept of “small” deflection is now related to the time and the tension. Again by physical observation, it can be assumed that to obtain a “small” deflection near the xy -plane all one needs to consider is that the tension itself is “small” in the $\hat{k}$ direction and is applied for but a “small” period of time Δt .

The above concept of “smallness” is now embedded into the monadic world. The infinitesimal parallelogram representation for the membrane surface is assumed to move parallel to the z -axis. Thus let $A = (a, b, c)$, $B = (a, b + dy, c)$, $C = (a + dx, b + dy, c)$, $D = (a + dx, b, c)$, where positive dx , $dy \in \mu(0)$. Considering only the $\hat{k}$ component of the tension $\vec{T}$, we write this component as $\|\vec{T}\| \cos \gamma$, where γ is a function continuous on G_1 . By continuity $\|\vec{T}(x, y, z, t)\| \cos \gamma(x, y, z, t) \approx \|\vec{T}(a, b, c, t)\| \cos \gamma(a, b, c, t)$ for each $(x, y, z, t) \in \mu^n((a, b, c, t))$, $n = 3$.

For the $\hat{k}$ component, what infinitesimal property will characterize the standard modeling notion that the deflection must be “small” compared to the size of the membrane? **(B) Certainly, the smallest possible N-world deflection would occur if there were NO apparent standard deflection at all. But, to analyze the problem nontrivially it would need to be assumed that deflection did take place in the monadic world.**

Monadic Restrictions of Physical Processes

The idea that a physical process may be restricted to a monadic environment in such a way that it does not appear to occur in the standard world but does occur in the monadic environment is yet another significant infinitesimal modeling technique.

In order to apply the monadic restriction technique to this problem and without actually evaluating the functions, simply let

$$\text{st}(\|\vec{T}(a, b, c)\| \cos \gamma(a, b, c)) = 0, \quad \vec{T}(a, b, c) \neq \vec{0}. \quad (4)$$

(C) Notice the important modeling technique that from the standard world point of view the infinitesimal rectangle $ABCD$ appears as if it is the single point (a, b, c) . From

N-world physical observation, and indeed the definition of the equilibrium position, a single point will vibrate only if it is displaced either above or below the equilibrium position. By *-transferring this observation to the infinitesimal rectangle $ABCD$, it follows that in order to have a “small” deflection take place “near” the equilibrium position $ABCD$ should be placed either above or below the hyper- xy -plane and all members of $ABCD$ should be infinitely close to the hyper- xy -plane. Let $c = z(a, b, t_0)$.

The hyperline segments $\overline{AB}$ and $\overline{AD}$ are geometric elements for membrane (surface) curves. This fact and our definition (9.5.1) for tangents to space curves leads to the conclusion that the hyperslope of $\overline{AD} \approx z_x(a, b, t_0)$ and hyperslope of $\overline{AB} \approx z_y(a, b, t_0)$. Since every point in $ABCD$ is infinitely close to the hyper- xy -plane, then slope $\overline{AB}$ and $\overline{AD}$ are both infinitely close to 0. Observe that since by continuity there exists some open ball about (a, b, c) such that $\vec{T} \neq \vec{0}$ for any member of this ball then ${}^*\vec{T}(x, y, z,) \neq \vec{0}$ for any $(x, y, z,) \in \mu^n((a, b, c))$. Expression (4) now implies that

$${}^*\cos {}^*\gamma(a, b, c) \approx {}^*\cos {}^*\gamma(a + dx, b, c) \approx {}^*\cos {}^*\gamma(a, b + dy, c) \approx 0. \quad (5)$$

Further, using the continuity of z_x, z_y we obtain

$$z_x(a, b, t_0) \approx z_x(a + dx, b, t_0) \approx z_y(a, b, t_0) \approx z_y(a, b + dy, t_0) \approx 0. \quad (6)$$

The problem we now face is how to select the proper combination of entities from (5) and (6). The following expressions (7), (8), (9) and (10) are obtained by means of the observation that, **(D) mathematically, the coordinate spaces are independent one from the other and, physically, tensions parallel to the z -axis along one boundary of the hyperrectangle do not effect substantially the adjacent boundaries.** Since ${}^*\vec{T}$ is a limited vector for each $(x, y, z,) \in \mu^n((a, b, c))$ then

$$\| {}^*\vec{T}(a, b, c) \| {}^*\cos {}^*\gamma(a, b, c) \approx \| {}^*\vec{T}(a, b, c) \| z_x(a, b, t_0), \quad (7)$$

$$\| {}^*\vec{T}(a, b, c) \| {}^*\cos {}^*\gamma(a, b, c) \approx \| {}^*\vec{T}(a, b, c) \| z_y(a, b, t_0), \quad (8)$$

$$\begin{aligned} & \| {}^*\vec{T}(a + dx, b, c) \| {}^*\cos {}^*\gamma(a + dx, b, c) \approx \\ & \| {}^*\vec{T}(a + dx, b, c) \| z_x(a + dx, b, t_0), \end{aligned} \quad (9)$$

$$\begin{aligned} & \| {}^*\vec{T}(a, b + dy, c) \| {}^*\cos {}^*\gamma(a, b + dy, c) \approx \\ & \| {}^*\vec{T}(a, b + dy, c) \| z_y(a, b + dy, t_0). \end{aligned} \quad (10)$$

Expression (2) must now be infinitesimalized. The major method used for such infinitesimalizing is call **(E) the (Differential Equation) Method of Maximum and Minimum. In this case, it leads to the conclusion that forces may be considered as attached to the A, B, D vertices of this physical (geometric) element.** Before proceeding, might there be a purely physical NSP-world argument for selecting the values for T at the points A, B, D - an argument that is substantially different from the usual one that such a selection is done since it “works”? Recently Simhony [1987] has advanced a theory that claims that all of the material universe is composed of combinations of the configuration he has termed as the electron-positron lattice. This is a simple lattice structure with electrons and positrons located at the lattice nodes. For such a theory, these objects would be the building blocks of the material universe and all forces could be considered as attached to the lattice nodes. Our infinitesimal representation $ABCD$ is uniquely determined by

the vertices A , B , and D . Even though Simhony's investigations of the possible lattice nature of the material universe have interesting conclusions that could by *-transfer lead to an effective lattice structure within the NSP-world, such a structure can also evolve from mathematical considerations.

The function $\|\vec{T}(x, y, z)\| \cos \gamma(x, y, z)$ is continuous on every standard rectangle determined by vertices $A = (a, b, c)$, $B_1 = (a, b + b_1, c)$, $C_1 = (a + a_1, b + b_1, c)$, $D_1 = (a + a_1, b, c)$, where $0 < a_1 \leq r_1$, $0 < b_1 \leq r_2$ for some $r_1, r_2 \in \mathbb{R}$. Thus, since the line segment $\overline{AB_1}$ is compact, then there exists points (a, b_m, c) , $(a, b_M, c) \in \overline{AB_1}$ such that

$$\begin{aligned} \|\vec{T}(a, b_m, c)\| \cos \gamma(a, b_m, c) &\leq \\ \|\vec{T}(a, y, c)\| \cos \gamma(a, y, c) &\leq \|\vec{T}(a, b_M, c)\| \cos \gamma(a, b_M, c) \end{aligned} \quad (11)$$

for each $(a, y, c) \in \overline{AB_1}$. Hence,

$$\begin{aligned} \|\vec{T}(a, b_m, c)\| \cos \gamma(a, b_m, c) |\overline{AB_1}| &\leq \\ \|\vec{T}(a, y, c)\| \cos \gamma(a, y, c) |\overline{AB_1}| &\leq \\ \|\vec{T}(a, b_M, c)\| \cos \gamma(a, b_M, c) |\overline{AB_1}|, \end{aligned} \quad (12)$$

for each $(a, y, c) \in \overline{AB_1}$. Whether it be obtained by observation, consideration of the integral or by axiomation, method (E) states, for this application, that there exists a net scalar tension F' such that

$$\begin{aligned} \|\vec{T}(a, b_m, c)\| \cos \gamma(a, b_m, c) |\overline{AB_1}| &\leq \\ F' &\leq \|\vec{T}(a, b_M, c)\| \cos \gamma(a, b_M, c) |\overline{AB_1}| \end{aligned} \quad (13)$$

and F' may be considered the force that accelerates the rigid body $\overline{AB_1}$. Assuming continuity on connected $\overline{AB_1}$ then there exists some $(a, \bar{b}, c)$ such that

$$\|\vec{T}(a, \bar{b}, c)\| \cos \gamma(a, \bar{b}, c) |\overline{AB_1}| = F', \quad (a, \bar{b}, c) \in \overline{AB_1} \quad (14)$$

and in like manner for the other three tensions. Following (2) we have that the total force of motion may be considered the sum of these four scalar $\pm \hat{k}$ directed tensions. The results in this paragraph are now *-transferred to the NSP-world, in which case (14) becomes

$$\|{}^*\vec{T}(a, \bar{b}, c)\| {}^*\cos {}^*\gamma(a, \bar{b}, c) dy = F', \quad (15)$$

where $b \leq \bar{b} \leq b + dy$.

Continuity at (a, b, c) implies that $\|{}^*\vec{T}(a, \bar{b}, c)\| {}^*\cos {}^*\gamma(a, \bar{b}, c) \approx \|{}^*\vec{T}(x, y, c)\| {}^*\cos {}^*\gamma(x, y, c)$, where $(x, y, c) \in \{(a, b, c), (a + dx, b, c), (a, b + dy, c), (a + dx, b + dy, c)\} = A$. Now selection of the appropriate members of A comes from repeating the above argument and *-transfer of the behavior. Applying (15) and (2) this implies that the effective scalar tension, ${}^*F_T$, that moves the hyperrectangle $ABCD$ has the property that

$$\begin{aligned} {}^*F_T &\approx \|{}^*\vec{T}(a + dx, b, c)\| {}^*\cos {}^*\gamma(a + dx, b, c) dy - \\ &\quad \|{}^*\vec{T}(a, b, c)\| {}^*\cos {}^*\gamma(a, b, c) dy + \\ &\quad \|{}^*\vec{T}(a, b + dy, c)\| {}^*\cos {}^*\gamma(a, b + dy, c) dx - \\ &\quad \|{}^*\vec{T}(a, b, c)\| {}^*\cos {}^*\gamma(a, b, c) dx. \end{aligned} \quad (16)$$

Applying (7), (8), (9) and (10) yields

$$\begin{aligned}
 {}^*F_T &\approx \|\vec{T}(a+dx, b, c)\| {}^*z_x(a+dx, b, t_0) dy - \\
 &\quad \|\vec{T}(a, b, c)\| {}^*z_x(a, b, t_0) dy + \\
 &\quad \|\vec{T}(a, b+dy, c)\| {}^*z_y(a, b+dy, t_0) dx - \|\vec{T}(a, b, c)\| {}^*z_y(a, b, t_0) dx.
 \end{aligned} \tag{17}$$

Applying (E) again and by *-transfer, we note that there exists points $(\bar{a}, \bar{b})$, (c, d, f, g) such that the mass, m , of the hyperrectangle is ${}^*\rho(\bar{a}, \bar{b}) dx dy$ and the entire lumped load L can be represented by ${}^*h(c, d, f, g) dx dy$, where $(\bar{a}, \bar{b}) \approx (a, b)$ and $(c, d, f, g) \approx (a, b, c, t_0)$. By continuity, ${}^*\rho(\bar{a}, \bar{b}) dx dy \approx \rho(a, b) dx dy$; ${}^*h(c, d, f, g) dx dy = {}^*h dx dy \approx h(a, b, c, t_0) dx dy$. Writing expression (1) in a scalar form for motion parallel to the z -axis obtains ${}^*F(t_0)/m \approx ({}^*v(t_0 + dt) - v(t_0))/dt$ and the total scalar force producing a change in the instantaneous velocity is ${}^*F_T + {}^*h(c, d, f, g)$. Hence,

$$\begin{aligned}
 &\frac{{}^*v(t_0 + dt) - v(t_0)}{dt} \approx \\
 &\frac{\|\vec{T}(a+dx, b, c)\| {}^*z_x(a+dx, b, t_0) dy - \|\vec{T}(a, b, c)\| {}^*z_x(a, b, t_0) dy}{{}^*\rho(\bar{a}, \bar{b}) dx dy} + \\
 &\quad \frac{\|\vec{T}(a, b+dy, c)\| {}^*z_y(a, b+dy, t_0) dx}{{}^*\rho(\bar{a}, \bar{b}) dx dy} - \\
 &\quad \frac{\|\vec{T}(a, b, c)\| {}^*z_y(a, b, t_0) dx + {}^*h dx dy}{{}^*\rho(\bar{a}, \bar{b}) dx dy} = \\
 &\frac{\|\vec{T}(a+dx, b, c)\| {}^*z_x(a+dx, b, t_0) - \|\vec{T}(a, b, c)\| {}^*z_x(a, b, t_0)}{{}^*\rho(\bar{a}, \bar{b}) dx} + \\
 &\frac{\|\vec{T}(a, b+dy, c)\| {}^*z_y(a, b+dy, t_0) - \|\vec{T}(a, b, c)\| {}^*z_y(a, b, t_0)}{{}^*\rho(\bar{a}, \bar{b}) dy} + \\
 &\quad \frac{{}^*h}{{}^*\rho(\bar{a}, \bar{b})} \approx \frac{{}^*z_t(t_0 + dt) - z_t(t_0)}{dt}.
 \end{aligned} \tag{18}$$

The derivation is completed by applying the standard part operator to expression (18) and employing the hypothesized requirement that the various partials exist.

The following are some useful observations relative to the above derivation. If one assumes that the tension $\vec{T}$ and the density ρ are constant along with $h \equiv 0$, then the usual elementary 2-dimensional wave equation follows from (3). More to the point, however, is the, not necessarily obvious, fact that the derivation was not obtained by attempting to comprehend second order rates of change. Berkeley was correct in his criticism that such things are difficult to mentally perceive. **(F) The general overall approach in the derivation the vibrating membrane equation was to consider first-order rates of change of other rates of change that had already been investigated and found to be representable by infinitely close (locally linear) functions expressed in partial derivative form.** This general overall approach is almost always the best and least confusing.

10.4. Internal Heat Transfer

Consider a vertical (rectangular) slab of material that uniformly conducts heat where “heat flow” is perpendicular to the left and right-hand surfaces and “flows” to the right. We quote from **Sears and Zemansky [1952]**: “The figure represents a slab of material of cross section A and thickness L . Let the whole of the left face of the slab be kept at a temperature U_2 , and the whole of the right face at a lower temperature U_1 . The direction of heat current is then from left to right through the slab.

After the faces of the slab have been kept at the temperatures U_1 and U_2 for a sufficient length of time, the temperature at points within the slab is found to decrease uniformly [linearly] with distance from the hot to the cold face. At each point, however, the temperature remains constant with time. The slab is said to be in a “linear steady state.”

It is found by experiment that the rate of flow of heat through the slab in the steady state is proportional to the area A , proportional to the temperature difference $(U_2 - U_1)$, and inversely proportional to the thickness L .”

Let Q be the usual measure of the quantity of heat, $0 < \Delta t \in \mathbb{R}$ the time, $\neq 0 < K \in \mathbb{R}$ the constant of proportionality (the *thermal conductivity*) and $0 < A \in \mathbb{R}$ the area. Then for this very special and simplistic case, it follows that

$$Q = \frac{KA(\Delta t)(U_2 - U_1)}{L}. \quad (1)$$

Notice that since this is a rectangular solid and the distance L is measured perpendicular to the faces then the convention of heat flow is taken normal to the faces.

Before the modern theory of heat conduction, heat was thought of as a invisible weightless fluid called *caloric* which was produced when a substance burned and which could be transmitted by conduction from one body to another. The analogy of “flowing heat” has persisted and due to the above steady state law this direction is taken normal to this section. For the moment, assume that the temperature $U(a, y, z, t_0)$ is only dependent upon time and thickness, a , of the slab. Now alter the position of the coordinate system relative to the rectangular slab and let $a + L$, $L > 0$, represent the thickness of the of the slab and $\Delta t = t - t_0$. Our first natural law is now expressed as

$$Q = \frac{KA(\Delta t)(U(a, y, z, t_0) - U(a + L, y, z, t_0))}{L}. \quad (2)$$

Let the specific heat, σ , and the density, ρ , for a given “small” rectangular solid, R , of volume, V , be constants. At time t_0 let the temperature of the entire solid be a constant $U(t_0)$. After a time change Δt and if there is but a “small” positive temperature change $U(x, y, z, t_0 + \Delta t)$ which appears also to be constant throughout R , then a second physical law appears to emerge. This law states that the amount of heat, H , necessary to achieve such a temperature change is

$$H = \rho \sigma V (U(x, y, z, t_0 + \Delta t) - U(x, y, z, t_0)), \quad (3)$$

where, for this simplistic case, U is not dependent upon position.

We need two more requirements prior to our derivation of the heat equation. First, recall that Count Rumford suggested that heat was really energy in another form and Sir James Joule experimentally verified this natural law. In what follows, we use the concepts of the conservation and additivity of heat energy. The second requirement is Theorem 10.4.1. For nonempty open $G \subset \mathbb{R}^n$, let $f: G \rightarrow \mathbb{R}$. Denote any $\vec{v} \in G$ by $\vec{v} = (x_1, \dots, x_n)$. Let $y = x_i$, $1 \leq i \leq n$. Recall

that a set C_y is said to be **convex in the direction y** if for any two points $\vec{p} = (p_1, \dots, p_n)$, $\vec{q} = (p_1, \dots, p_i + h_i, \dots, p_n) \in C_y$ and any $k \in \mathbb{R}$ such that $\vec{w} = (p_1, \dots, p_i + k, \dots, p_n)$ is a member of the line segment with end point $\vec{p}$, $\vec{q}$, then $\vec{w} \in C_y$.

Theorem 10.4.1. *Let $f: G \rightarrow \mathbb{R}$, where nonempty open $G \subset \mathbb{R}^n$ and standard $\vec{v} \in G$. Suppose that $y = x_i$, $1 \leq i \leq n$, f_y is defined on G and continuous at $\vec{v}$. Let $C_y \subset {}^*G$, where internal C_y is $*$ -convex in the direction y . If for $h \in \mu(0)$, such that $\vec{p} = (p_1, \dots, p_i, \dots, p_n)$, $\vec{q} = (p_1, \dots, p_i + h, \dots, p_n) \in C_y$ and $\vec{v} \approx \vec{p}$, then there exists $\epsilon \in \mu(0)$ such that*

$${}^*f(\vec{q}) - {}^*f(\vec{p}) = f_y(\vec{v})h + h\epsilon.$$

Application 10.4.1 *A differential equation model for 3-dimensional heat transfer.*

All functions are assumed to be defined on a open neighborhood, G , of (a, b, c, t_0) where, it appears necessary, to assume that $G \cap \mathbb{R}^3$ contains the closed solid and, at the least, the functions are continuous where defined. Let $U(x, y, z, t)$ represent the temperature, never zero $\rho(x, y, z,)$ the density of the material, never zero $\sigma(x, y, z,)$ the specific heat and a constant thermal conductivity K . It is assumed that σ, ρ do not vary with temperature. This constraint implies that our differential equation model is only valid for small temperature changes. Even though no sources or sink are assumed for this application, they can be easily adjoined to the derivation in a manner analogous to the lumped load of application 10.3.1. Let U_x, U_y, U_z , be continuous on G and U_t be defined on G and continuous at (a, b, c, t_0) . Let U_{xx}, U_{yy}, U_{zz} , exist at (a, b, c, t_0) . For any such $(x, y, z, t) = (a, b, c, t_0)$ a differential equation model for internal heat transfer is

$$\rho(a, b, c) \sigma(a, b, c) \frac{\partial U}{\partial t} = K \left(\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} + \frac{\partial^2 U}{\partial z^2} \right).$$

Derivation. Equation (2) assumes that we are dealing with a uniform temperature change in that we have a steady state condition. Still retaining the idea that the temperature is constant on a section, (2) must be generalized to the case where the temperature is not uniformly decreasing with the thickness of the slab. We make a strong appeal to the flow analogy and, thus, consider the temperature evaluation to take place along a flow line parallel to $\hat{i}$. With this in mind, $*$ -transfer yields

$${}^*Q = \frac{K {}^*A(\Delta t)(U(a, b, c, t_0) - {}^*U(a + \epsilon, b, c, t_0))}{\epsilon}, \quad (4)$$

where $0 < L = \epsilon \in \mu(0)$. Can we consider the thermal conductivity as a nonconstant function? It appears that if K were not constant, then the differential equation for heat transfer would not be derivable. Fortunately, in most physical cases, the constancy of K can be verified. Assume that $\partial U / \partial x$ exists at (a, b, c, t_0) . Then (4) leads to the nonsteady state expression

$$\begin{aligned} Q = \text{st}({}^*Q) &= \text{st} \left(\frac{K {}^*A(\Delta t)(U(a, b, c, t_0) - {}^*U(a + \epsilon, b, c, t_0))}{\epsilon} \right) = \\ &= -KA(\Delta t) \left(\frac{\partial U(a, b, c, t_0)}{\partial x} \right). \end{aligned} \quad (5)$$

Equation (5) is conceived of as representing the quantity of heat after the time period Δt that “flows” through a rectangular shaped section cut from a conducting solid; but, we are still assuming

that the temperature is constant on each section. [Remark: *It is claimed by some authors that equation (5) expressed in words with this flow convention has been experimentally obtained. They then use (5), along with integral concepts and the Divergence Theorem to formulate the heat equation. I shall only take (2) as fundamental.*]

Once again applying the directed flow analogy, the external view of the net amount of heat that would accumulate in the interior of the slab from this $\hat{i}$ direction flow would be

$$H_{\hat{i}} = -KA(\Delta t) \left(\frac{\partial U(a, b, c, t_0)}{\partial x} \right) - \left(-KA(\Delta t) \left(\frac{\partial U(a + L, b, c, t_0)}{\partial x} \right) \right). \quad (6)$$

Now letting V be the volume of the slab notice that

$$H_{\hat{i}}/(V(\Delta t)) = K(U_x(a + L, b, c, t_0) - U_x(a, b, c, t_0))/L. \quad (7)$$

Unfortunately, (7) still refers to a temperature that is constant on the parallel faces of the slab. To generalize to the nonconstant case, first infinitesimalize the slab itself. By *-transfer, (7) holds for an infinitesimal rectangular solid R having determining vertices $A = (a, b, c)$, $B = (a + dx, b, c)$, $C = (a, b + dy, c)$, $D = (a, b, c + dz)$, $0 < dx, dy, dz \in \mu(0)$ and for an infinitesimal time period $0 < dt \in \mu(0)$. Hence, (7) becomes

$$^*H_{\hat{i}}/(dx dy dz)(dt) = K(^*U_x(a + dx, b, c, t_0) - U_x(a, b, c, t_0))/dx. \quad (8)$$

Next we correct for the specialized constant temperature case by assuming that heat flow takes place only along parallel flow lines. Concentrating upon (7), where the temperature is not assumed to be constant over the sections, we apply the maximum and minimum method (E) from the last section and *-transfer the result to the faces of R . Thus, there exists some $(b_1, c_1) \in \mu((b, c))$ such that the total heat per unit volume-time accumulated by the $\hat{i}$ direction flow is

$$^*H_{\hat{i}}/(dx dy dz)(dt) = K(^*U_x(a + dx, b_1, c_1, t_0) - ^*U_x(a, b_1, c_1, t_0))/dx. \quad (9)$$

Applying Theorem 10.4.1 yields

$$^*H_{\hat{i}}/(dx dy dz)(dt) = KU_{xx}(a, b, c, t_0) + \epsilon_1, (\epsilon_1 \in \mu(0)), \quad (10)$$

and repeating the argument for the $\hat{j}$ and $\hat{k}$ directions (and applying the *-transfer of the additivity of energy etc.) yields that the total heat accumulated per volume- time for R , as viewed externally, is

$$^*H/(dx dy dz)(dt) \approx K(U_{xx}(a, b, c, t_0) + U_{yy}(a, b, c, t_0) + U_{zz}(a, b, c, t_0)). \quad (11)$$

Now consider the simplistic natural law represented by equation (3). For a more general case, where the temperature, density, and even the specific heat vary by position (but not temperature) and are continuous on an open neighborhood G , one can and apply the maximum and minimum method (E). Viewing σ , ρ , U as a functions of position only, where t_0 and Δt are assumed fixed. Then (dependent upon t_0 and Δt) there exists some $\vec{p} \in$ the solid such that

$$H = V\rho(\vec{p})\sigma(\vec{p})(U(\vec{p}, t_0 + \Delta t) - U(\vec{p}, t_0)). \quad (12)$$

Equation (12) can be written for nonzero Δt in the form

$$H/(V\rho(\vec{p})\sigma(\vec{p})\Delta t) = (U(\vec{p}, t_0 + \Delta t) - U(\vec{p}, t_0))/\Delta t. \quad (13)$$

By *-transfer, there exists some $\vec{q} \in R$ such that

$$*H/(dx dy dz * \rho(\vec{q}) * \sigma(\vec{q}) dt) = (*U(\vec{q}, t_0 + dt) - *U(\vec{q}, t_0))/dt. \quad (14)$$

Applying Theorem 10.4.1 and the fact that $*\rho(\vec{q})$ and $*\sigma(\vec{q})$ are limited, we obtain

$$*H/(dx dy dz)(dt) \approx *\rho(\vec{q}) *\sigma(\vec{q}) U_t(a, b, c, t_0). \quad (15)$$

The final step is clear. From expressions (11) and (15) we have

$$\begin{aligned} *\rho(\vec{q}) *\sigma(\vec{q}) U_t(a, b, c, t_0) &\approx K(U_{xx}(a, b, c, t_0) + \\ &U_{yy}(a, b, c, t_0) + U_{zz}(a, b, c, t_0)) \end{aligned} \quad (16)$$

and application of the standard part operator coupled with the continuity of ρ and σ at (a, b, c) implies the result that

$$\rho(a, b, c) \sigma(a, b, c) \frac{\partial U}{\partial t} = K \left(\frac{\partial^2 U}{\partial x^2} + \frac{\partial^2 U}{\partial y^2} + \frac{\partial^2 U}{\partial z^2} \right). \quad (17)$$

10.5. Concluding Remarks

For some derivations, an alternative to the Monadic Second Law of Motion may be necessary due to the ring nature of the infinitesimals and the limited numbers. Let us compare the Newtonian view discussed in section 10.1, Definition 10.3.1 and the fundamental expression (1) of section 10.4. If one lets the scalar velocity $v(t)$ linearly change with respect to a “small” time change Δt (i.e. uniformly changing in the old terminology), then observation does indeed confirm that the constant scalar force that produces such a change is $F(\Delta t) = m((v(t_0 + \Delta t) - v(t))/\Delta t)$. If this last statement is assumed to hold for all nonnegative time changes less than or equal to Δt then, as was done in order to obtain equation (4) of section 10.4, *-transfer yields

$$*F(\epsilon) = *m((v(t_0 + \epsilon) - v(t))/\epsilon), (\epsilon \in \mu(0)). \quad (1)$$

From (1) and application of the standard part operator Newton’s Second Law of Motion follows. If one removes the standard part operator, then all that can be stated is that

$$*F(\epsilon) \approx *m((v(t_0 + \epsilon) - v(t))/\epsilon), (\epsilon \in \mu(0)). \quad (2)$$

Equation (1) is the alternate NSP-world form of Newton’s Second Law of Motion. It may be necessary to utilize this alternate form for those cases where division by infinitesimals and other similar algebraic processes are used - processes that do not preserve $\approx$. Further, the use of (E) and Theorem 10.4.1 are useful in order to retain expressions that do not include the external notion of $\approx$. For the derivation of the heat equation it was necessary to retain equational expressions until steps (15) and (16).

Is it necessary to make the derivation of integral or differential equation models more rigorous or have I wasted you time? Some mathematicians claim that such an exercise is without merit since all

one needs is a good guess; and if the magically obtained equation is solvable and reasonably predicts natural system behavior, then this is all that is required. Not with standing such pronouncements, other very productive mathematicians believe that there are a few obvious advantages to a more rigorous approach.

There is an advantage in education. In the physical sciences, beginning students observe simplistic behavior and follow the well-established path of attempting to comprehend complex behavior as a composition of the simplistic. Extending these intuitively grasped concepts to the NSP-world leads more directly to integral or differential equation models and gives them a stronger incentive to seek solution methods and investigate the mathematical structures. Motivation still remains one of our major educational devices.

Scientific theories are not immutable. New experimental scenarios, new measuring devices and, indeed, new insights into the foundations of natural system behavior have, historically, led to new theories that appear to predict more accurately system development. Many new insights and theories abound, today, within certain scientific journals that specialize in such speculation.

In the early part of this century, two of the great achievements of the human mind were the Special and General Theories of Relativity. Yet, as has been shown [Herrmann [1995]], the major experimentally verified conclusions of the Special Theory are obtainable by simple infinitesimal modeling, without the use of such concepts as frames of reference and the like. I believe that this fact will tend to eliminate some of the controversy that still surrounds the Special Theory and eventually lead to a better understanding of the basic nature of electromagnetic radiation. I have previously mentioned how the recent concept of fractal behavior has been successfully modeled within the NSP-world environment.

Indirectly, we are learning the simplistic NSP-world behavior that may be assumed to be the cause for certain natural world effects that are measured by standard instruments. As is well-known, predicting accurately the behavior of many macroscopic natural systems, even with our present day theories, is often notably unsuccessful. It may be that the causes for such behavior are objectively real in character within a actual substratum world. And, hence, the more proficient we become in analyzing and applying infinitesimal techniques the more likely we are to development new mathematical models that improve upon such predictions.

NOTES

[1] In section 10.1 the concept of two vectors being almost parallel was introduced. The following theorem is sometimes useful.

Theorem N.1. *Two unit vectors, $\vec{v}, \vec{u} \in {}^*\mathbb{R}^n$ have the property that $\vec{v} \approx \pm \vec{u}$ if and only if $\vec{v} \bullet \vec{u} \approx \pm 1$.*

[2] Relative to the statement on page 67 as to the relation between the work done along a hyperline segment and the curve if we add to the hypotheses of Application 8.2.1 the requirement that for each $t \in [a, b]$, $c'(t) \neq \vec{0}$ and that F is uniformly continuous on E , then using the notation in that section it follows that there exists some $\epsilon_j \in \mu(0)$ such that ${}^*F(\ell_j(t'_j)) \bullet {}^*\vec{v}_j = {}^*F({}^*c(t'_j)) \bullet {}^*\vec{v}_j + \epsilon_j \|{}^*\vec{v}_j\|$. From this we also have that $\sum_{j=1}^{\Gamma} {}^*F(\ell_j(t'_j)) \bullet {}^*\vec{v}_j \approx \sum_{j=1}^{\Gamma} {}^*F({}^*c(t'_j)) \bullet {}^*\vec{v}_j$. The proof is at the end of Appendix 10.

Appendix For Chapter 2.

Throughout these proofs lower case Greek letters will always denote infinitesimals.
 [Remark: The complete proofs for many of these fundamental propositions appear for the first time in these appendixes.]

Theorem 2.1.1. *Assumption (II) holds for ${}^*\mathbb{R}$ if and only if ${}^*\mathbb{R}$ is not Archimedean.*

Proof. Assume that (II) holds. Since ${}^*\mathbb{R}$ is a field then $\epsilon^{-1} \in {}^*\mathbb{R}$. Let $0 \neq n \in \mathbb{N}$. Then $0 \neq \epsilon < 1/n$. Thus $n < \epsilon^{-1}$ implies that ${}^*\mathbb{R}$ is not Archimedean.

Conversely, assume that ${}^*\mathbb{R}$ is not Archimedean. Then there exists some $\Gamma \in {}^*\mathbb{R}$ such that for each $n \in \mathbb{N}$, $|\Gamma| \geq n$. Since $|\Gamma| \in {}^*\mathbb{R}$, then it follows that for each $n \in \mathbb{N}$, $|\Gamma| \geq n+1 > n \Rightarrow |\Gamma^{-1}| < 1/n$, $\forall n \in \mathbb{N}$, $n \neq 0$ and the result follows.

Theorem 2.2.1. *The set of limited numbers, $\mathcal{O}$, is a subring of ${}^*\mathbb{R}$, $\mathbb{R} \subset \mathcal{O}$ and $\mathcal{O}$ is not a field.*

Proof. Let $x, y \in \mathcal{O}$. Then there exists some $r_1, r_2 \in \mathbb{R}$ such that $|x| < r_1$, $|y| < r_2$. Thus $|x \pm y| \leq |x| + |y| < r_1 + r_2 \in \mathbb{R} \Rightarrow x \pm y \in \mathcal{O}$, $|xy| = |x||y| < r_1 r_2 \in \mathbb{R} \Rightarrow xy \in \mathcal{O}$. If $0 \neq \epsilon$, then $|\epsilon| < r^{-1}$, $\forall r \in \mathbb{R}^+ \Rightarrow \epsilon^{-1} \notin \mathcal{O}$. The result follows.

Theorem 2.2.2. *The infinitesimals, $\mu(0)$, form a subring of $\mathcal{O}$ and Γ is infinite if and only if there is some nonzero $\epsilon \in \mu(0)$ such that $\Gamma = 1/\epsilon$.*

Proof. Let $x, y \in \mu(0)$. Then $\forall r \in \mathbb{R}^+$, $|x| < r/2$, $|y| < r/2$, $\Rightarrow |x \pm y| < r$. Now we also know that $|x| < \sqrt{r}$, $|y| < \sqrt{r}$. Hence, $|xy| = |x||y| < r$. Since r is positive and arbitrary then the first result follows. The second follows as in the proof of Theorem 2.2.1.

Theorem 2.2.3. *The set of infinitesimals, $\mu(0)$, is an ideal of $\mathcal{O}$.*

Proof. Let $b \in \mathcal{O}$. Then $\exists t \in \mathbb{R}^+$ such that $|b| < t$. Also $\forall r \in \mathbb{R}^+$, $\forall x \in \mu(0)$, $|x| < r/t$. Hence $|xb| < r$ and the result follows.

Theorem 2.2.4. *The binary relation $\approx$ is an equivalent relation on ${}^*\mathbb{R}$.*

Proof. (i) Since $0 \in \mathcal{O}$ then $x - x = 0 \in \mu(0) \Rightarrow x \approx x$.

(ii) Let $x \approx y$. Then $x - y \in \mu(0) \Rightarrow -(x - y) = y - x \in \mu(0) \Rightarrow y \approx x$.

(iii) Let $x \approx t$, $y \approx z$. Then $x - y = \epsilon$, $y - z = \delta$. Then $\epsilon + \delta \in \mu(0) \Rightarrow x - z \in \mu(0) \Rightarrow x \approx z$.

Theorem 2.2.5. *For each $x, y \in \mathbb{R}$,*

(i) $\mu(x) \cap \mu(y) = \emptyset$ if and only if $x \neq y$,

(ii) $\mathcal{O} = \bigcup \{\mu(r) | r \in \mathbb{R}\}$.

Proof. (i) follows since each monad is an equivalence class for the equivalence relation $\approx$.

(ii) As previously noted $\mathbb{R} \subset \mathcal{O}$. Note that $\mu(r) \cap \mu(r+1) = \emptyset$. Let $y \in \mu(r)$, $z \in \mu(r+1)$. Then $y = r + \epsilon$, $z = r+1 + \delta$. Assume that $z \leq y$. Then $r+1 + \delta \leq r + \epsilon \Rightarrow 1 \leq \epsilon - \delta \in \mu(0)$: a contradiction. Thus $y < z$. But $r+1 \in \mu(r+1) \Rightarrow y < r+1$. In like manner, $r-1 < y$. Thus $y \in \mathcal{O} \Rightarrow \mu(r) \subset \mathcal{O} \Rightarrow \bigcup \{\mu(r) | r \in \mathbb{R}\} \subset \mathcal{O}$. Now let arbitrary $x \in \mathcal{O}$. and define $S = \{y | y \in \mathbb{R} \wedge y < x\}$. The set S is bounded above and nonempty since $\exists r \in \mathbb{R}^+$ such that $|x| < r$, and $-r \in S$. By completeness $\sup S \in \mathbb{R}$. For each $p \in \mathbb{R}^+$, $\exists y \in S$ such that $\sup S - p < y < x$. Also $x \leq \sup S + p$,

for otherwise we have that $\sup S + p \in S$, which contradicts the sup definition. Thus $|x - \sup S| \leq p$. Hence $x - \sup S \approx 0 \Rightarrow x \in \mu(\sup S) \Rightarrow \mathcal{O} \subset \bigcup \{\mu(r) | r \in \mathbb{R}\}$ and the result follows.

Corollary 2.2.4.1. *If $x, y \in \mathbb{R}$, $x < y$, $z \in \mu(x)$, $w \in \mu(y)$, then $z < w$.*

Proof. Since $x < y$ then $\exists r \in \mathbb{R}^+$ such that $y = x + r$. Now consider the obvious portion of the above argument applied to the monads $\mu(x)$, $\mu(x + r)$.

Corollary 2.2.4.2. *If $x, y \in \mu(r)$, $z \in {}^*\mathbb{R}$, $x < z < y$, then $z \in \mu(r)$.*

Proof. Since $|y - z| < |y - x| < r$, $\forall r \in \mathbb{R}^+$ then $y - z \in \mu(0)$. Thus $r \approx y$, $y \approx z$, $\Rightarrow r \approx z \Rightarrow z \in \mu(r)$.

Theorem 2.2.6. *Each monad and the set of limited numbers are bounded above [resp. below], but do not possess a least upper bound [resp. greatest lower bound].*

Proof. This is established for the “bounded above” case only. Let $\mu(r)$ be any arbitrary monad. From Corollary 2.2.4.1, the set $\mu(r)$ is bounded above by the real number $r + 1$. Assume that $\sup \mu(r) = s \in {}^*\mathbb{R}$. First, assume that $s \in \mathbb{R}$. In this case, it is necessary that $s \neq r$ for we know $\exists w > 0$ and if $s = r$, then $s + w \in \mu(r)$, $s < s + w$, contradicts the bounding character of s . Thus $r < s \Rightarrow \forall x \in \mu(r)$, $\forall y \in \mu(s)$; $x < y$. But $-w + s \in \mu(s) \Rightarrow \forall x \in \mu(r)$; $x < -w + s < s$. Thus s would not be the $\sup \mu(r)$. We must, therefore, pass to the second possible case that $s \in {}^*\mathbb{R} - \mathbb{R}$. Obviously, $r < s < r + 1$ and thus $s \in \mathcal{O}$. Consequently, $\exists r_1 \in \mathbb{R}$ such that $s \in \mu(r_1)$. Let $r_1 \neq r$. Then $r < r_1$ and $\forall x \in \mu(r)$, $\forall y \in \mu(r_1)$, $x < y$. But again using $-w + s$ we would have that $s \neq \sup \mu(r)$. The final case, requires $s \in \mu(r)$. But if this were so, then $s < s + w \in \mu(r)$ yields that s would not be an upper bound. Thus no such s exists and the proof is complete for monads.

Let $s = \sup \mathcal{O}$. Then $s \notin \mathcal{O}$ since if not, then $s + 1 \in \mathcal{O}$. On the other hand, if s is infinite, then $s > 0 \Rightarrow s - 1$ is infinite and that $s - 1 > 0$. Thus no such s exists in ${}^*\mathbb{R}$ and the proof is complete.

Theorem 2.3.1. *For any $n \in \mathbb{N}$, $n \geq 1$, the set of limited vectors in ${}^*\mathbb{R}^n$ is equal to*

$$\mathcal{O}^n = \overbrace{\mathcal{O} \times \cdots \times \mathcal{O}}^{n \text{ factors}}.$$

Proof. See Theorem 2.3.3

Theorem 2.3.2. *For any $n \in \mathbb{N}$, $n \geq 1$, then monad of the vector $\vec{v} = (x_1, \dots, x_n) \in \mathbb{R}^n$ is equal to*

$$\mu(\vec{v}) = \overbrace{\mu(x_1) \times \cdots \times \mu(x_n)}^{n \text{ factors}}.$$

Proof. See Theorem 2.3.3

Theorem 2.3.3. *For any $n \in \mathbb{N}$, $n \geq 1$, and for each $\vec{v}, \vec{w} \in \mathbb{R}^n$,*

- (i) $\mu(\vec{v}) \cap \mu(\vec{w}) = \emptyset$ if and only if $\vec{v} \neq \vec{w}$,
- (ii) $\mathcal{O}^n = \bigcup \{\mu(\vec{v}) | \vec{v} \in \mathbb{R}^n\}$.

Proof. Theorems 2.3.1 and 2.3.2 follow from the *-transform of the basic Euclidean norm fact that if $\vec{v} = (x_1, \dots, x_n) \in \mathbb{R}^n$, then $\forall i$, $1 \leq i \leq n$, $|x_i| \leq \|\vec{v}\| \leq \sum_{i=1}^n |x_i|$ and the definitions. Theorem 2.3.3 can be established directly or it follows easily from the basic set-theoretic properties for the cross product.

Theorem 2.4.1. *Let $x, y \in \mathcal{O}$. Then*

- (i) $x \approx y$ if and only if $\text{st}(x) = \text{st}(y)$,
- (ii) $x \approx \text{st}(x)$,
- (iii) if $x \in \mathbb{R}$, then $\text{st}(x) = x$,
- (iv) if $x \leq y$, then $\text{st}(x) \leq \text{st}(y)$,
- (v) if $\text{st}(x) \leq \text{st}(y)$, then either $x \leq y$ or $x - y \in \mu(0)$ with unknown order.

Proof. Observe that for the limited hyperreal number x , that $x = \text{st}(x) + \epsilon$.

(i) $x, y \in \mathcal{O} \Rightarrow x = \text{st}(x) + \epsilon; x \approx y \Rightarrow x = y + \delta \Rightarrow y + \delta = \text{st}(x) + \epsilon \Rightarrow y = \text{st}(x) + (\epsilon - \delta) \Rightarrow y \in \mu(\text{st}(x)) \Rightarrow \text{st}(x) = \text{st}(y)$.

If $\text{st}(x) = \text{st}(y)$, then $x = \text{st}(x) + \epsilon, y = \text{st}(y) + \delta \Rightarrow x - \epsilon = y - \delta \Rightarrow x \approx y$.

(ii) $x = \text{st}(x) + \epsilon \Leftrightarrow x \approx \text{st}(x)$.

(iii) For $r \in \mathbb{R}, r \in \mu(r) \Rightarrow r = \text{st}(r)$.

(iv) Let $x \leq y$. Then $x, y \in \mu(0) \Rightarrow \text{st}(x) = \text{st}(y)$. If $x \in \mu(r), y \in \mu(r_1), r \neq r_1$, then $r > r_1 \Rightarrow x > y$. Thus contradiction yields that $\text{st}(x) = r < \text{st}(y) = r_1$.

(v) Assume that $\text{st}(x) = \text{st}(y)$. Then $x - y \in \mu(0)$. If $\text{st}(x) < \text{st}(y)$, then $x \in \mu(\text{st}(x)), y \in \mu(\text{st}(y)) \Rightarrow x < y$.

Theorem 2.4.2. *Let $x, y \in \mathcal{O}$. Then*

- (i) $\text{st}(x \pm y) = \text{st}(x) \pm \text{st}(y)$,
- (ii) $\text{st}(xy) = (\text{st}(x))(\text{st}(y))$.

Proof. Notice that in general, if $r \in \mathbb{R}$, then $\text{st}(r + \epsilon) = r$. Let $r = \text{st}(x), s = \text{st}(y)$.

(i) $\text{st}(x \pm y) = \text{st}((r + \epsilon) \pm (s + \delta)) = \text{st}((r \pm s) + (\epsilon \pm \delta)) = r \pm s$.

(ii) $\text{st}(xy) = \text{st}(rs + (r\delta + s\epsilon + \delta\epsilon)) = rs$.

Corollary 2.4.2.1 *Let $x, y \in \mathcal{O}$. Then*

- (i) if $\text{st}(y) \neq 0$, then $\text{st}(x/y) = \text{st}(x)/\text{st}(y)$.
- (ii) if $y = \sqrt[n]{x}$, then $\text{st}(y) = (\text{st}(x))^{(1/n)}$, where it is always the case that if $x \in \mathcal{O}$, then $\sqrt[n]{x} \in \mathcal{O}$.

Proof. If $\text{st}(y) \neq 0$, then $y \notin \mu(0) \Rightarrow y^{-1} \in \mathcal{O} \Rightarrow (x/y) \in \mathcal{O}$. Then $\text{st}(x) = \text{st}((x/y)y) = (\text{st}(x/y))(\text{st}(y))$.

(ii) Obviously, $x \geq 0$. Also, if $x \geq 0, x \in \mathcal{O}$, then $x < r \in \mathbb{R}^+ \Rightarrow \sqrt[n]{x} < \sqrt[n]{r} \Rightarrow \sqrt[n]{x} \in \mathcal{O}$. Then $y = \sqrt[n]{x} \Rightarrow y^n = x \Rightarrow (\text{st}(y))^n = \text{st}(x)$ and from this the proof is completed.

Theorem 2.4.3. *For $\vec{v} \in \mathcal{O}^n$, let $\text{st}(\vec{v}) = (\text{st}(x_1), \dots, \text{st}(x_n))$. The set $\mathcal{O}^n$ forms a vector space with respect to the ring $\mathcal{O}$ (i.e. a module) and as such the map st distributes over the vector space algebra as well as the component defined dot and for $n = 3$ the cross product operators.*

Proof. This follows immediately from the above standard part operator properties.

Theorem 2.4.3. *The set $\mu(0)$ is a maximal ideal in $\mathcal{O}$ and the quotient ring $\mathcal{O}/\mu(0)$ is isomorphic to $\mathbb{R}$.*

Proof. Let $\mathfrak{S}$ be an ideal such that $\mu(0) \subset \mathfrak{S} \subset \mathcal{O}$ and assume that $\mu(0) \neq \mathfrak{S}$. Let $x \in \mathfrak{S} - \mu(0)$. We know that $x^{-1} \in \mathcal{O}$. Since $\mathfrak{S}$ is an ideal in $\mathcal{O}$ then $x(x^{-1}) = 1 \in \mathfrak{S}$. Thus implies that $\mathfrak{S} = \mathcal{O}$. The standard part operator is obviously the isomorphism and this completes the proof.

Appendix For Chapter 3.

Throughout these proofs lower case Greek letters will always denote infinitesimals, while upper case will denote infinite numbers.

Even though it is possible to restrict our model to objects such as $\mathfrak{R}$ it is useful for future considerations to actually construct a general superstructure as originally envisioned by Robinson and Zakon [1969]. Let $X_0 = \mathbb{R}$ and define by induction $X_{n+1} = \mathcal{P}(\bigcup\{X_j | 0 \leq j \leq n\})$. Recalling that it is assumed that $\mathbb{R}$ is a set of atoms (i.e. they are not sets). Then $\mathcal{H} = \bigcup\{X_n | n \in \mathbb{N}\}$ is the **universe** or **carrier** for a structure $\mathcal{M} = (\mathcal{H}, \epsilon, =)$. In this particular case, this universe is termed a **superstructure on $\mathbb{R}$** . In the literature there are slightly different constructions of such superstructures [Davis 1977]. An object $b \in X_0$ is an individual and otherwise if $b \in X_p$ then b is an entity. Thus b is an individual or entity if and only if $b \in \mathcal{H}$. It will be shown that $\exists p \in \mathbb{N}$ such that $\mathfrak{R} \in X_p$. A superstructure has many different properties. The following is a brief list of some of the more important of these, where the proofs are straightforward and left to the reader. [Note: Later I may change notation and use the usual mathematical practice of considering symbols such as x, y, z in two contexts. One, the a mathematical variable, and the other a formal first-order language variable.]

Proposition 1. *Let $\mathcal{H}$ be a superstructure on $\mathbb{R}$.*

- (i) $X_p \subset X_n$ for $n, p \in \mathbb{N}$, $1 \leq p \leq n$. $X_0 \cap X_n = \emptyset$, $n \geq 1$.
- (ii) $X_0 \cup X_n = X_0 \cup \dots \cup X_n$, $X_{n+1} = \mathcal{P}(X_0 \cup X_n)$, $n \geq 0$.
- (iii) $X_p \in X_{n+1}$, $\forall n \in \mathbb{N}$, $n \geq 1$, $0 \leq p \leq n$.
- (iv) If for $n \geq 1$, $a \in b \in X_n$, then $a \in X_0 \cup X_{n-1}$.
- (v) If $(a_1, \dots, a_n) \in b \in X_p$ ($p \geq 1$), then $a_1, \dots, a_n \in X_0 \cup X_{p-1}$.
- (vi) If $b \in X_n$ and $a \subset b$, then $a \in X_n$.

By use of concepts in abstract model theory, the nonstandard model $^*\mathcal{M} = (^*\mathcal{H}, \epsilon, =)$, $^*\mathcal{H} = \bigcup\{^*X_n | n \in \mathbb{N}\}$ is constructed through application of the notion of internal individuals or entities and, in particular, (iv) of Proposition 1. [For a very clear and concise discussion of this please see page 120 of Robinson and Zakon [1969].] With respect to our notation, this construction also leads to the following significant proposition that gives a general relation between internal objects and elements they contain.

Proposition 2. *Within our set theory*

- (i) If $a \in Q \in ^*X_{n+1}$, then $a \in ^*X_0 \cup ^*X_n$. (Every member of an internal entity is internal. Also note that this is the set-theoretic “or” statement which is a set even if it contains atoms.)
- (ii) The set $^*\mathcal{H}$ is closed under finitely many applications of the basic set operation $\cup$.
- (iii) Any finite collection of members from $^*\mathcal{H}$ is a member of $^*\mathcal{H}$.
- (iv) The set $^*\mathcal{H}$ is closed under n -tuple formation.
- (v) If $(a_1, \dots, a_p) \in Q \in ^*X_{n+1}$, then $a_i \in ^*X_0 \cup ^*X_n$, $1 \leq i \leq p$.
- (vi) The set $^*\mathcal{H}$ is closed under the basic set-theoretic operators $\cap$, $-$, $\times$, among others, for internal sets.

Proof. (i) Sometimes this result is established during construction. However, it follows easily by * -transfer of Proposition 1 (iv).

(ii) Note that for any $n \geq 0$ the set $X_n \cup X_0 \in X_{n+1}$. Hence, by Theorem 3.1.3 (i), $^*(X_n \cup X_0)$ is internal. Now we can use the independently established part (vi) d of Theorem 3.1.3 and obtain

that $*X_0 \cup *X_n = *(X_n \cup X_0)$. Let A and B represent internal entities. For any $n \geq 0$, the sentence

$$\forall x \forall y ((x \in X_{n+1} \wedge (y \in X_{n+1} \rightarrow \exists z ((z \in X_{n+1} \wedge \forall w (w \in (X_n \cup X_0) \rightarrow (w \in z \leftrightarrow (w \in x) \vee (w \in y))))))$$

holds in $\mathcal{M}$; hence, in $*\mathcal{M}$. Now since A and B are internal, there is an n such that $A, B \in *X_{n+1} \cup *X_0 = *(X_{n+1} \cup X_0)$. Thus, there is a $C \in *X_{n+1}$ that contains the same elements of $*X_n \cup *X_0$ as those in “ $A \cup B$.” But by (i), all the elements in A, B, C are members of $*X_n \cup *X_0$. Thus $C = A \cup B \in *\mathcal{H}$. This proof is completed by induction.

(iii) Theorem 3.1.3 (vi)b, shows that this holds for an empty set. The notion of what constitutes the “number” of members of a nonempty finite set is an intuitive metamathematical notion the corresponds this idea to the “number” of $\vee$ symbols. [Note: This is not the only way this can be established. Indeed, we could use finite set generating operators.] For each $n \geq 0$, the sentence

$$\forall x (x \in (X_n \cup X_0) \rightarrow \exists y ((y \in X_{n+1} \wedge \forall z ((z \in (X_n \cup X_0) \wedge z \in y \leftrightarrow z = x))))$$

holds in $\mathcal{M}$; hence, it holds in $*\mathcal{M}$. Now given any internal $a \in *\mathcal{H}$, then $a \in *X_0 \cup *X_n$ for some $n \geq 0$. Thus, for any $a \in *\mathcal{H}$ the set that contains a and only a exists and is an internal. Now to establish this for the notion of a two element with two distinct members, we extend the above sentence. Thus for any $n \geq 0$ the sentence

$$\forall x \forall y (x \in (X_n \cup X_0) \wedge y \in (X_n \cup X_0) \wedge x \neq y \rightarrow \exists y ((y \in X_{n+1} \wedge \forall z ((z \in (X_n \cup X_0) \wedge z \in y \leftrightarrow (z = x) \vee (z = y))))))$$

holds in $\mathcal{H}$; hence, it holds in $*\mathcal{H}$. But, we know that for any two internal a, b there exists some n such that $a \in *X_n \cup *X_0$. Then every set that contains two and only two distinct internal entities is internal by *-transfer. To apply this to any set of finitely many internal entities we use induction or application of (ii).

(iv) Within our set-theoretic definitions for X_0 or $*X_0$ order pair formation is given for specific a, b . We use the abbreviation (a, b) to denote the finite set $\{\{a\}, \{a, b\}\}$. This is then extended by induction to the n-tuple $(a_1, \dots, a_{n-1}, a_n) = ((a_1, \dots, a_{n-1}), a_n)$. Thus any n-tuple is just a finite set, of finite sets, of finites sets, ... constructed in the finite manner from the finite set $\{a_1, a_2, \dots, a_n\}$. For example, consider internal $a, b, \neq b$. Then they are both members of some $*X_n \cup *X_0$. Then $\{a\}$ and $\{a, b\}$ are internal and both members of some $*X_p$. Then the set $\{\{a\}, \{a, b\}\} \in *X_{p+1}$. Apply this idea to an n-tuple and you have that the n-tuple is internal.

(v) This follows from the definition of the n-tuple, repeated applications of (i) and Proposition 1 (i).

(vi) These are established in the same manner as (ii). The $\cap$ is established as an example after Theorem 3.4.1.

Theorem 3.1.1. *It is permissible to assume that*

- (i) if $A \subset \mathbb{R}$, then $A \subset *\mathbb{R}$,
- (ii) if $A \subset \mathbb{R}^n$, then $A \subset *\mathbb{R}^n$,
- (iii) if $A \subset (\mathbb{R}^n) \times (\mathbb{R}^m)$, then $A \subset (*\mathbb{R}^n) \times (*\mathbb{R}^m)$.

Proof.

(i) This follows from the construction of the nonstandard model $^*\mathcal{M} = (^*\mathcal{H}, \epsilon, =)$ by means of the superstructure technique where each member of the carrier $\mathbb{R}$ is treated as a constant sequence [Robinson and Zakon [1969], Stroyan and Luxemburg [1976], Davis [1977], etc.] Basically this follows from the fact that each member of $\mathbb{R}$ is an atom or urelement within our basic set theory.

(ii) Let $(a_1, \dots, a_n) \in A \subset \mathbb{R}^n$. By construction, $\mathbb{R}^n \in \mathcal{H}$. From our basic definition of an n -tuple as a finite set of finite sets etc., then this result is immediately established from independently proved Theorem 3.1.3 parts (i) and (iii) by finitely distributing the $*$ throughout these finite sets and using (i) of this theorem. If $A = \emptyset$, then result follows from Theorem 3.1.3 part (vi).

(iii) This is proved in the same manner as part (ii).

Theorem 3.1.2. *If $A \subset \mathbb{R}$ [resp. $\mathbb{R}^n$, $(\mathbb{R}^n) \times (\mathbb{R}^m)$], then there exists $^*A \subset ^*\mathbb{R}$ [resp. $^*\mathbb{R}^n$, $(^*\mathbb{R}^n) \times (^*\mathbb{R}^m)$] such that*

- (i) $A \subset ^*A$ and
- (ii) $A = ^*A$ if and only if A is finite.

Proof. From the independently established Theorem 3.4.1 it follows by $*$ -transfer that $^*A \subset ^*\mathbb{R}$.

(i) This follows from Theorem 3.1.3 part (i) and, as discussed in Theorem 3.1.1, from the construction of $^*\mathcal{M}$.

(ii) The model $^*\mathcal{M}$ is assumed, at least, to be an enlargement. Thus consider the binary relation $R = \{(x, y) | (x \in A) \wedge (y \in A) \wedge (x \neq y)\}$. Now let $\{(x_1, y_1), \dots, (x_n, y_n)\} \subset R$ and suppose that A is infinite. Then there exists some $b \in A$ such that $x_i \neq b$, $1 \leq i \leq n$. From the concurrency of R , this implies that there exists some internal $c \in ^*A$ such that $\forall a \in {}^\sigma A = A$, $a \neq c$. [The ${}^\sigma$ operator is that defined in Stroyan and Luxemburg, [1976]. In general, it is the set of all constant sequence $\mathcal{U}$ -equivalence classes contained in $^*A$.] The result now follows from Theorem 3.1.3.

Conversely, if $A = \emptyset$ or A is finite, then result follows from Theorem 3.1.3.

[Note: Since $X_{n+1} = \mathcal{P}(X_0 \cup X_n)$, then $(X_0 \cup X_n) \in X_{n+1}$ implies $(X_0 \cup X_n) \in \mathcal{H}$.

Theorem 3.1.3. *Unless otherwise stated all constants represent individuals or entities (i.e. members of $C(\mathcal{H})$ where we do not notationally differentiate between the name of an object and the object itself.)*

- (i) $a \in A$ if and only if $^*a \in ^*A$; $A \neq B$ if and only if $^*A \neq ^*B$.
- (ii) $A \subset B$ if and only if $^*A \subset ^*B$.
- (iii) $^*\{A_1, \dots, A_k\} = \{^*A_1, \dots, ^*A_k\}$.
- (iv) $^*(A_1, \dots, A_k) = (^*A_1, \dots, ^*A_k)$.
- (v) $(A_1, \dots, A_k) \in A$ if and only if $(^*A_1, \dots, ^*A_k) \in ^*A$.
- (vi) Let A, B be sets [resp subsets of $\mathcal{U}$]. Then (d) $^*(A \cup B) = ^*A \cup ^*B$, (c) $^*(A \cap B) = ^*A \cap ^*B$, (a) $^*(A - B) = ^*A - ^*B$, (e) $^*(A \times B) = ^*A \times ^*B$, (b) $^*\emptyset = \emptyset$.
- (vii) If $A \in \mathbb{R}$, then $^*A = A$.

Proof. (i) The sentences

$$a \in A, \neg(A = B)$$

hold in $\mathcal{M}$ if and only if the $*$ -transfers

$$^*a \in ^*A, \neg(^*A = ^*B)$$

hold in $^*\mathcal{M}$.

(vi) [Clearly, by construction $\mathcal{U} \in \mathcal{H}$. Consequently, A, B are entities by Proposition 1.] (a) We know that there exists a $C \in C(\mathcal{H})$ such that $C = A - B$. The sentence

$$\forall x(x \in C \leftrightarrow (x \in A) \wedge (\neg(x \in B)))$$

holds in $\mathcal{M}$ if and only if

$$\forall x(x \in {}^*C \leftrightarrow (x \in {}^*A) \wedge (\neg(x \in {}^*B)))$$

holds in $^*\mathcal{M}$. Then the set ${}^*C = {}^*A - {}^*B$. Now (b) is obtained by letting $\emptyset = \mathbb{R} - \mathbb{R}$. Then from (a), ${}^*\emptyset = {}^*\mathbb{R} - {}^*\mathbb{R} = \emptyset$. (c) We also know that there is a $C \in \mathcal{H}$ such that $C = (A \cap B) = A - (A - B)$. Once again by (a); (c) follows since ${}^*(A \cap B) = {}^*A - ({}^*A - {}^*B) = {}^*A \cap {}^*B$. Then for (d) there is a $A \cup B = C \in \mathcal{H}$. Now simply consider the sentence $\forall x((x \in C) \leftrightarrow (x \in A) \vee (x \in B))$. This result follows by * -transfer. (Another proof for (c) also follows from * -transfer.) (ii) This follows from parts (a) and (b) of (vi) since $A \subset B$ iff $A - B = \emptyset$. (vi) (e) The Cartesian product statement is a little more difficult to establish. In this elementary approach, it is useful to characterize the notion of the singleton and doubleton sets. The set $S(A)$ of all singleton subsets of A and the set $D(A)$ of all doubleton subsets of A are members of X_{n+2} (i.e. $D(A) \subset X_{n+1}$). The following sentences characterize the singleton and doubleton subsets of $A \subset X_n \cup X_0$.

$$\Phi_1 = \forall x(x \in D(A) \rightarrow \exists y \exists w(w \in A \wedge y \in A \wedge y \in x \wedge$$

$$w \in x \wedge \forall z(z \in X_n \cup X_0 \wedge z \in x \rightarrow z = y \vee z = w))).$$

[Note: $P \wedge Q \rightarrow S \equiv P \rightarrow (Q \rightarrow S)$.]

$$\Phi_2 = \forall x(x \in S(A) \rightarrow \exists y(y \in A \wedge y \in x \wedge$$

$$\forall z(z \in X_n \cup X_0 \wedge z \in x \rightarrow z = y))).$$

The * -transfer of these sentences becomes

$${}^*\Phi_1 = \forall x(x \in {}^*D(A) \rightarrow \exists y \exists w(w \in {}^*A \wedge y \in {}^*A \wedge y \in x \wedge$$

$$w \in x \wedge \forall z(z \in {}^*X_n \cup {}^*X_0 \wedge z \in x \rightarrow z = y \vee z = w))).$$

$${}^*\Phi_2 = \forall x(x \in {}^*S(A) \rightarrow \exists y(y \in {}^*A \wedge y \in x \wedge$$

$$\forall z(z \in {}^*X_n \cup {}^*X_0 \wedge z \in x \rightarrow z = y))).$$

[Note: ${}^*D(A) = {}^*(D(A))$, ${}^*S(A) = {}^*(S(A))$.] In what follows, the constants used to name various internal objects are in our extended internal language. What these * -transforms indicate is that the set ${}^*S(A) \subset S({}^*A)$, ${}^*D(A) \subset D({}^*A)$. Indeed, the same would hold true for any finitely numbered subset of A . Observe first that ${}^*D(A)$, ${}^*S(A) \subset {}^*X_{n+1}$ and ${}^*A \subset {}^*X_n \cup {}^*X_0$. For the doubleton case, let $Q \in {}^*D(A)$. Thus each such Q is internal since it is a member of ${}^*X_{n+1}$. There then exists internal $c, d \in {}^*A$ and for any $f \in {}^*X_n \cup {}^*X_0$ and $f \in Q$ it follows that $f = c$ or $f = d$. Now since ${}^*A \subset {}^*X_n \cup {}^*X_0$ then this last statement also includes any $f \in {}^*A$. Proposition 2 (i) states that every $z \in Q$ is internal and a member of ${}^*X_n \cup {}^*X_0$. Thus only $c, d \in Q \Rightarrow Q \in D({}^*A)$. Hence ${}^*D(A) \subset D({}^*A)$.

Now consider the sentences

$$\begin{aligned}\Phi_3 &= \forall x \forall y (x \in A \wedge y \in A \rightarrow \exists z (z \in D(A) \wedge x \in z \wedge y \in z \wedge \\ &\quad \forall w (w \in X_n \cup X_0 \wedge w \in z \rightarrow w = x \vee w = y))) \\ \Phi_4 &= \forall x (x \in A \rightarrow \exists z (z \in S(A) \wedge x \in z \wedge \\ &\quad \forall w (w \in X_n \cup X_0 \wedge w \in z \rightarrow w = x))).\end{aligned}$$

Once again consider the *-transfer

$$\begin{aligned}{}^*\Phi_3 &= \forall x \forall y (x \in {}^*A \wedge y \in {}^*A \rightarrow \exists z (z \in {}^*D(A) \wedge x \in z \wedge y \in z \wedge \\ &\quad \forall w (w \in {}^*X_n \cup {}^*X_0 \wedge w \in z \rightarrow w = x \vee w = y))) \\ {}^*\Phi_4 &= \forall x (x \in {}^*A \rightarrow \exists z (z \in {}^*S(A) \wedge x \in z \wedge \\ &\quad \forall w (w \in {}^*X_n \cup {}^*X_0 \wedge w \in z \rightarrow w = x))).\end{aligned}$$

Now let W, U be objects within our set theory. Let $W \in D({}^*A)$. Then $\exists a, b \in {}^*A$ such that for any $U \in W$, in particular $U \in {}^*X_n \cup {}^*X_0$, $U = a$ or $U = b$. The *-transfer states that there exists some $Q \in {}^*(D(A))$ such that $a, b \in Q$. Thus $W \subset Q$. Postulate 2 states that if any $z \in Q$, then $z \in {}^*X_n \cup {}^*X_0$ and *-transfer yields that $z = a$ or $z = b \Rightarrow W = Q \Rightarrow D({}^*A) \subset {}^*D(A) \Rightarrow {}^*D(A) = D({}^*A)$. In like manner, ${}^*S(A) = S({}^*A)$. (This is not the only way to establish these equalities. I have chosen to use the operators S and D since these and similar operators could prove useful in later investigations.)

In order to establish that ${}^*(A \times B) = {}^*A \times {}^*B$ similar techniques are employed. First, noting that $C = A \times B \in X_n$ for some $n > 1$ we have that the sentence

$$\begin{aligned}\Phi_5 &= \forall z (z \in C \rightarrow \exists x \exists y \exists w \exists u (x \in S(A) \wedge y \in D(A \cup B) \wedge \\ &\quad w \in A \wedge u \in B \wedge w \in x \wedge w \in y \wedge u \in y))\end{aligned}$$

holds in $\mathcal{M}$ if and only if (iff)

$$\begin{aligned}{}^*\Phi_5 &= \forall z (z \in {}^*C \rightarrow \exists x \exists y \exists w \exists u (x \in S({}^*A) \wedge y \in D({}^*A \cup {}^*B) \wedge \\ &\quad w \in {}^*A \wedge u \in {}^*B \wedge w \in x \wedge w \in y \wedge u \in y))\end{aligned}$$

holds in ${}^*\mathcal{M}$. This implies that ${}^*C \subset {}^*A \times {}^*B$. For the converse, it is clear from the result that $S({}^*F) = {}^*S(F)$, $D({}^*F) = {}^*D(F)$ and *-transfer that the sentence

$$\begin{aligned}{}^*\Phi_6 &= \forall x \forall y \forall w \forall z (x \in S({}^*A) \wedge y \in D({}^*A \cup {}^*B) \wedge w \in {}^*A \wedge \\ &\quad z \in {}^*B \wedge w \in x \wedge w \in y \wedge z \in y \rightarrow \exists v (v \in {}^*C \wedge x \in v \wedge y \in v \wedge \\ &\quad \forall u (u \in {}^*X_0 \cup {}^*X_{n-1} \wedge u \in {}^*C \rightarrow u = v))).\end{aligned}$$

holds in ${}^*\mathcal{M}$. Let $W \in {}^*A \times {}^*B$. Then through application of Proposition 2 and arguing as above there is a unique $U \in {}^*(A \times B)$ such that $U = W$. From this the result follows.

Proofs such as part (e) of this theorem using the above method are considered tedious and often do not appear in the literature. In modern nonstandard analysis they are considered trivial examples of the Leibniz Principle and are often left to the reader to obtain. I disagree with such sentiments. It is easy for an author to say these concepts "are expressible by appropriate first-ordered statements

and the result easily follows.” It is, however, not so easy for the untrained to actually do so for certain subtle requirements such as applications of Proposition 2 could easily allude the neophyte.

(iii) This is proved by induction with respect to the set $N_1 = \{n | n \in \mathbb{N} \wedge n \geq 1\}$. Let $n = 1$, $A_1 \in \mathcal{H}$. Then there exists some $p \in \mathbb{N}$, ($p \geq 1$) such that $C = \{A_1\} \in X_p$. Thus $*C \subset *X_0 \cup *X_{p-1}$. The sentence

$$\forall x(x \in X_0 \cup X_{p-1} \rightarrow (x \in C \leftrightarrow x = A_1))$$

holds in $\mathcal{M}$ if and only if

$$\forall x(x \in *X_0 \cup *X_{p-1} \rightarrow (x \in *C \leftrightarrow x = *A_1))$$

holds in $*\mathcal{M}$. Thus $*A \in *C$ and $d \in *C \Rightarrow d \in *X_0 \cup *X_{p-1} \Rightarrow d = *A_1$ yields that $*C = \{A_1\}$. Assume that result holds for some $n \in N_1$ and consider the set $\{A_1, \dots, A_n, A_{n+1}\}$. Note that $\{A_1, \dots, A_n, A_{n+1}\} \in X_p$ for some $p \in \mathbb{N}$ since it is a finite set. Now simply note from the previous case and part (vi) that $\{A_1, \dots, A_n, A_{n+1}\} = \{A_1, \dots, A_n\} \cup \{A_{n+1}\} \Rightarrow *\{A_1, \dots, A_n\} \cup *\{A_{n+1}\} = \{A_1, \dots, A_n\} \cup \{A_{n+1}\} = *\{A_1, \dots, A_n\}$.

(iv) This follows by induction, the definition of n-tuples and part (iii).

(v) This follows by the *-transfer of sentences composed only of constants such as in part (i) and from (iv).

(viii) See proof of part (i) of Theorem 3.1.1. This completes this proof.

Part (vi) (e) of Theorem 3.1.3 has some very useful consequences that allows for a great simplification of our first-order language. First, by induction, one has immediately that $*(A_1 \times \dots \times A_n) = (*A_1 \times \dots \times *A_n)$. Also since any object in $*(A_1 \times \dots \times A_n)$ is an actual set-theoretic n-tuple from the basic definition then for any $a_i \in *A_i$, $1 \leq i \leq n$ **the symbol $(a_1, \dots, a_n)$ may be used as the correct set-theoretic abbreviation for the unique set termed an n-tuple. This simplification is utilized continually throughout the remainder of these manuals.**

[Note: There are numerous equivalent forms that can be used to represent a statement formally in the required “bound” form so that each quantified variable is forced to vary only over members of $\mathcal{H}$ or $*\mathcal{H}$. In this regard, notice that $\mathcal{H}$ [resp. $*\mathcal{H}$] is closed under the basic set-theoretic operations for standard [resp. internal] objects. Thus, for example, if $A \in \mathcal{H}$ and $B \in \mathcal{H}$, then a statement such as $\forall x((x \in A) \wedge (x \in B) \dots)$ satisfies the bound requirement since this is equivalent to $\forall x(x \in A \cup B \dots)$ and $A \cup B \in \mathcal{H}$.]

Theorem 3.1.4. Let $\emptyset \neq R \subset A_1 \times \dots \times A_n$, ($n > 1$),

(i) $*P_i(R) = P_i(*R)$.

(ii) If R is a binary relation (i.e. $n = 2$), then $*R^{-1} = (*R)^{-1}$.

Proof. (i) This is established by induction on the number, m , of Cartesian products. First note that, in general, $R \subset A_1 \times \dots \times A_n \Rightarrow *R \subset *A_1 \times \dots \times *A_n$ and $P_1(R) \subset A_1 \Rightarrow *(P_1(R)) \subset *A_1$. Let $m = 1$. Consider the *-transfer of the following sentence, where the symbol (x, y) is an abbreviation for an obvious but more complex first-order expression.

$$\forall x(x \in A_1 \rightarrow (x \in P_1(R) \leftrightarrow \exists y(y \in A_2 \wedge (x, y) \in R)).$$

Then $q \in *(P_1(R)) \Rightarrow \exists s \in *A_2$ such that $(q, s) \in *R \Rightarrow *(P_1(R)) \subset P_1(*R)$. On the other hand, if $q \in P_1(*R)$, then $q \in *A_1$ and $\exists s \in *A_2$ such that $(q, s) \in *R \Rightarrow q \in *(P_1(R))$. Thus $P_1(*R) = *P(R)$. In like manner for the projection P_2 .

Assume that result holds for an arbitrary $m \in \mathbb{N}_1$, (i.e. for P_i , $1 \leq i \leq m+1$). Consider any $R \subset A_1 \times \cdots \times A_{m+2}$. Now for a little general set theory. From the definition $(A_1 \times \cdots \times A_{m+2}) = (A_1 \times \cdots \times A_{m+1}) \times A_{m+2} = C \times A_{m+2}$ and $R \subset C \times A_{m+2}$. Let $R_C = P_C(R)$. Then for each i , $1 \leq i \leq m+1$, $P_i(R_C) = P_i(R)$. These set-theoretical facts hold for any n -ary relation including $*R$. Thus, from the induction hypothesis, $\forall i$, $1 \leq i \leq m+1$, $*(P_i(R)) = P_i(P_C(*R)) = P_i(*R)$. Then $*(P_{m+2}(R)) = P_{m+2}(*R)$ follows from the first case where $m = 1$.

(ii) This follows from the $*$ -transfer of the sentence

$$\forall x \forall y (x \in A_1 \wedge y \in A_2 \rightarrow ((x, y) \in R \leftrightarrow (y, x) \in R^{-1})).$$

Proposition 3. *If internal $a \in {}^*\mathcal{H}$ and $b \in X_0$, then $a \notin b$.*

Proof. This is often proved directly during the construction of ${}^*\mathcal{M}$ but it also follows by $*$ -transfer of each sentence, where $n \in \mathbb{N}$,

$$\forall x \forall y (x \in X_0 \wedge y \in X_n \rightarrow \neg(y \in x)).$$

Theorem 3.1.5. *Let A be an entity. Then*

- (i) $*\{(x, x) | x \in A\} = \{(x, x) | x \in {}^*A\}$.
- (ii) $*\{(x, y) | x \in y \in A\} = \{(z, w) | z \in w \in {}^*A\}$.

Proof. (i) Let I_A be the identity relation defined on A . Then by $*$ -transfer of the sentence

$$\forall z (z \in I_A \rightarrow \exists x (x \in A \wedge (x, x) = z))$$

it follows that $*(I_A) \subset I({}^*A)$. On the other hand, $*$ -transfer of the sentence

$$\forall x (x \in A \rightarrow (x, x) \in I_A)$$

yields that $I({}^*A) \subset *(I_A)$ and thus $I({}^*A) = *(I_A)$.

(ii) Let $E = \{(x, y) | x \in y \in A\}$, where $A \in X_p$. If $p \leq 1$, then $E = \emptyset$ by Proposition 3 and result is obtained. Hence let $p \geq 2$, $a \in b \in A \in X_p \Rightarrow a \in b \in X_{p-1} \Rightarrow a \in X_0 \cup X_{p-2}$. Now let $K = \{(z, w) | z \in w \in {}^*A\}$. The sentence

$$\forall z (z \in {}^*E \rightarrow \exists x \exists y (x \in {}^*X_0 \cup {}^*X_{p-2} \wedge y \in {}^*X_{p-1} \wedge$$

$$x \in y \wedge y \in {}^*A \wedge z = (x, y)))$$

holds in ${}^*\mathcal{M}$. Thus ${}^*E \subset K$. On the other hand, consider the $*$ -transform

$$\forall x \forall y (x \in {}^*X_0 \cup {}^*X_{p-2} \wedge y \in {}^*X_{p-1} \wedge x \in y \in {}^*A \rightarrow (x, y) \in {}^*E).$$

Now if $(c, d) \in K$, then $c \in d \in {}^*A \Rightarrow c \in d \in {}^*X_{p-1} \Rightarrow a \in {}^*X_0 \cup {}^*X_{p-2}$ by Proposition 2. This implies that $K \subset {}^*E$ and completes the proof.

In the next demonstration we meet for the first time what I call a partial $*$ -transfer. What this means is that for some arbitrary object within a set a sentence is discussed and $*$ -transformed. After this discussion the result is obtained through application of the metalogic axiom of generalization.

Theorem 3.2.1. *Definition 3.2.1 (i.e. $S \rightarrow \vec{r}$) is equivalent to the limit definition for convergence of a sequence (i.e. $\lim_{n \rightarrow \infty} S(n) = \vec{r}$).*

Proof. For the sufficiency, let arbitrary $s \in \mathbb{R}^+$ and assume that $\lim_{n \rightarrow \infty} S(n) = \vec{r}$. Then $\exists N \in \mathbb{N}$ such that $\forall n > N$, $\|S(n) - \vec{r}\| < s$. Hence the sentence

$$\forall x(x \in {}^*\mathbb{N} \wedge x > N \rightarrow \|{}^*S(x) - \vec{r}\| < s)$$

holds in ${}^*\mathcal{M}$. In particular let $\Gamma \in \mathbb{N}_\infty$. Then $\|{}^*S(\Gamma) - \vec{r}\| < s$. But, $\Gamma > N$, $\forall N \in \mathbb{N} \Rightarrow \|{}^*S(\Gamma) - \vec{r}\| < s$, $\forall s \in \mathbb{R}^+$. Hence ${}^*S(\Gamma) \in \mu(\vec{r})$. Since Γ is also an arbitrary member of $\mathbb{N}_\infty$ the sufficiency is obtained.

For the necessity, assume that $\forall \Gamma \in \mathbb{N}_\infty$, ${}^*S(\Gamma) \in \mu(\vec{r})$. Let arbitrary $s \in \mathbb{R}^+$. Since $\mathbb{N}_\infty \neq \emptyset$ then the statement

$$\exists z(z \in {}^*\mathbb{N} \wedge \forall x(x > z \wedge x \in {}^*\mathbb{N} \rightarrow \|{}^*S(x) - \vec{r}\| < s))$$

holds in ${}^*\mathcal{M}$. Thus by removing the * notation

$$\exists z(z \in \mathbb{N} \wedge \forall x(x > z \wedge x \in \mathbb{N} \rightarrow \|S(x) - \vec{r}\| < s))$$

holds in $\mathcal{M}$. But, this is the limit definition for convergence and the proof is complete.

Theorem 3.2.2. *Let the sequence $S: \mathbb{N} \rightarrow \mathbb{R}^n$, ($n \geq 1$). Then $S \rightarrow \vec{r} \in \mathbb{R}^n$ if and only if for each $\Gamma \in \mathbb{N}_\infty$, $\text{st}({}^*S_\Gamma) = \vec{r}$.*

Proof. This follows from the definition of the standard part operator.

Theorem 3.4.1 (The Leibniz Principle). *A sentence Φ holds true for members of $\mathcal{H}$ if and only if the sentence ${}^*\Phi$ holds true for members of ${}^*\mathcal{H}$.*

Proof. The model theory portion of this proof is not elementary and requires additional effort. However, proofs can be found in [Stroyan and Luxemburg \[1976\]](#), [Davis \[1977\]](#).

From this all of our previous theorem involving * -transfer follow. The proofs that the internal objects are closed under the basic set-theoretic operation is very tedious. There are obtained by basic * -transfer and application of the previous results. The method is the same in all cases. Here is one example that shows that if you have a statement like

$$\Phi = \forall x(x \in A \rightarrow \exists y(y \in B \wedge x \subset y)),$$

then it holds in $\mathcal{M}$ if and only if the * -transfer holds in ${}^*\mathcal{M}$, where we do not * the relation symbol $\subset$. However, we must always translate this as stating in the * -transfer mode “internal x a subset of internal y .” This is so since this is an abbreviation for the statement $x \in ({}^*\mathcal{P})(y)$, where the power-set formation is considered as an operator. Consider the binary relation $E = \{(x, y) | x \in A \wedge y \in B \wedge x \subset y\}$. Then $\exists p \in \mathbb{N}$ such that $E \subset X_p$ and this implies also that $E \subset X_p \Leftrightarrow {}^*E \subset {}^*X_p$. By taking the * -transfer of

$$\forall z(z \in X_p \rightarrow (z \in E \leftrightarrow \exists x \exists y(x \in A \wedge y \in B \wedge$$

$$\forall w(w \in X_0 \cup X_{p-2} \wedge w \in x \rightarrow w \in y))),$$

applying Proposition 2 and the usual argument one obtains $*E = \{(x, y) | x \in *A \wedge y \in *B \wedge x \subset y\}$. From this the $*$ -transform may be written as

$$*\Phi = \forall x(x \in *A \rightarrow \exists y(y \in *B \wedge x \subset y)).$$

Thus as long as the $*$ -transfer is applied to sentences where the quantifiers are bounded (or assumed bounded) by standard entities then objects are closed under the basic binary set-theoretic operators. This even includes the more general $\bigcup X$ and $\bigcap X$ where X is an internal entity. However, it does not include all set-theoretic operators. In particular, if A is an infinite entity, then $\mathcal{P}(A)$ is not internal although $*\mathcal{P}(A) = *(\mathcal{P}(A)) = (*\mathcal{P})(*A)$. As seen by the proof of Theorem 3.4.3 every element of $*(\mathcal{P}(A))$ is an entity and $*$ -transfer of the characterizing statement it is also a subset of $*A$. Thus in those set-theoretic expressions where the power set operator is used, it must carry a $*$ notation under $*$ -transfer.

Theorem 3.4.2. *The set of infinite natural numbers $\mathbb{N}_\infty = *\mathbb{N} - \mathbb{N}$.*

Proof. Previously proved in Example 3.4.3.

Theorem 3.4.3. (Leibniz' Principle Restated). *A sentence Φ holds true for members of $\mathfrak{R} = \mathcal{U} \cup \mathcal{P}(\mathcal{U})$ if and only if the sentence $*\Phi$ holds true for the members of $*\mathcal{U}$ or internal subsets of $*\mathcal{U}$.*

Proof. Observe that $*\mathfrak{R} = *\mathcal{U} \cup *\mathcal{P}(\mathcal{U})$. The set $\mathcal{U} \in X_p \Rightarrow \mathcal{U} \subset X_0 \cup X_{p-1} \Rightarrow \mathcal{P}(\mathcal{U}) \subset \mathcal{P}(X_0 \cup X_{p-1}) = X_p \Rightarrow \mathcal{P}(\mathcal{U}) \in X_{p+1}$. Hence, $*(\mathcal{P}(\mathcal{U})) \in *X_{p+1}$. Since, by $*$ -transfer, no internal set is a member of $*X_0$, then by Proposition 2 (i) each member of $*\mathcal{P}(\mathcal{U})$ is an internal subset of $*\mathcal{U}$ and the proof is complete.

Theorem 3.5.1. *Let $S: \mathbb{N} \rightarrow \mathbb{R}$. If S is an increasing [resp. decreasing] sequence and there exists some $\Gamma \in \mathbb{N}_\infty$ such that $*S_\Gamma \in \mathcal{O}$, then $S \rightarrow \text{st}(*S_\Gamma)$.*

Proof. Assume that $S: \mathbb{N} \rightarrow \mathbb{R}$ is increasing. Then by $*$ -transfer $*S: *\mathbb{N} \rightarrow *\mathbb{R}$ is increasing. Hence for $\Gamma \in \mathbb{N}_\infty$ and for $m \in \mathbb{N}$ it follows since $\Gamma > m$ that $S(m) \leq *S(\Gamma)$. Consequently, since $*S(\Gamma) \in \mathcal{O}$ the $\text{st}(S(m)) = S(m) \leq \text{st}(*S(\Gamma)) = r \in \mathbb{R}$. Thus the sentence

$$\forall x(x \in \mathbb{N} \rightarrow S(x) \leq r)$$

holds in $\mathcal{M}$; hence in $*\mathcal{M}$. Thus $\forall \Omega \in \mathbb{N}_\infty$, $*S(\Omega) \leq r \Rightarrow \forall \Omega \in \mathbb{N}_\infty$, $*S(\Omega) \in \mathcal{O}$. Hence $\forall \Omega \in \mathbb{N}_\infty$, $\text{st}(*S(\Omega)) \in \mathbb{R}$. So, let arbitrary $\Omega \in \mathbb{N}_\infty$, $\Omega \neq \Gamma$, $\text{st}(*S(\Omega)) = r_0$. Then the same sentence as above with r_0 replacing r yields that $*S(\Gamma) \leq r_0 \Rightarrow \text{st}(*S(\Gamma)) = r \leq r_0$. Consequently, $\text{st}(*S(\Omega)) = r$. Since Ω is arbitrary then $*S(\Omega) \in \mu(r)$, $\forall \Omega \in \mathbb{N}_\infty \Rightarrow S \rightarrow r$.

Appendix For Chapter 4.

Theorem 4.1.1. *Definition 4.1.1 is equivalent to the limit definition for a sequence $S \rightarrow +\infty$.*

Proof. First, the method of partial transfer is used. Assume that $\lim_{n \rightarrow \infty} S_n = +\infty$. Let arbitrary $B \in \mathbb{R}^+$. Then from the classical definition there exists some $M_B \in \mathbb{N}$ such that the sentence

$$\forall x(x \in \mathbb{N} \wedge x \geq M_B \rightarrow S_x \geq B)$$

holds in $\mathcal{M}$. By $*$ -transfer this particular statement implies since $\forall \Gamma \in \mathbb{N}_\infty$, $\Gamma > M_B$ that $*S_\Gamma \geq B$. No matter what $B \in \mathbb{R}^+$ is chosen the same conclusion for each M_B . Thus in all cases $\forall \Gamma \in \mathbb{N}_\infty$, $*S_\Gamma \geq B$. But B is any arbitrary positive real number. Thus $\forall \Gamma \in \mathbb{N}_\infty$, $*S_\Gamma \in \mathbb{R}_\infty^+$.

For the converse, assume that $\forall \Gamma \in \mathbb{N}_\infty$, $*S_\Gamma \in \mathbb{R}_\infty^+$. Let $B \in \mathbb{R}^+$ and $\Omega \in \mathbb{N}_\infty$ which exists since $\mathbb{N}_\infty \neq \emptyset$. Obviously, if $\Lambda \geq \Omega$, then $\Lambda \in \mathbb{N}_\infty \Rightarrow *S_\Lambda > B$. Hence, the sentence

$$\exists x(x \in \mathbb{N} \wedge \forall y(y \in \mathbb{N} \wedge y \geq x \rightarrow S_y > B))$$

holds in $*\mathcal{M}$ and thus holds in $\mathcal{M}$. But this is the classical definition and the result follows.

Theorem 4.1.2. *Let $r \in *\mathbb{R}^+$, $S: \mathbb{N} \rightarrow \mathbb{R}$ and $S \rightarrow +\infty$. Then for each $\Omega \in \mathbb{N}_\infty$ there exists some $\Delta \in \mathbb{N}_\infty$ such that $*S(\Omega) + r \leq *S(\Delta)$.*

Proof. From Theorem 4.1.1 simply consider the $*$ -transfer of the complete definition for $\lim_{n \rightarrow \infty} S(n) = +\infty$. Take $*S(\Omega) + r$ as the bound. Then $\exists M \in *\mathbb{N}$ such that $\forall \Gamma \in *\mathbb{N}$, $\Gamma \geq M \Rightarrow *S(\Gamma) \geq *S(\Omega) + r$. Obviously, if $M \notin \mathbb{N}_\infty$, then any member of $\mathbb{N}_\infty$ will suffice and this completes the proof.

Theorem 4.2.1. (The Extended Standard Definition Principle)

(i) *A set $A \in *\mathcal{H}$ is an extended standard set (i.e. there exists some $B \in \mathcal{H}$ such that $A = *B$) if and only if there exists some standard set D (i.e. $D \in \mathcal{H}$) and a standard bound formula $\Phi(x)$ in one free variable where each constant in $\Phi(x)$ is a member of $C(\mathcal{H})$ and*

$$A = \{x | x \in *D \wedge *\Phi(x)\}.$$

(ii) *A set A is an extended standard n -ary relation ($n > 1$) if and only if there exist n standard sets $D_1, \dots, D_n$ and a standard bound formula $\Phi(x_1, \dots, x_n)$ in n free variables where each constant in $\Phi(x_1, \dots, x_n)$ is a member of $C(\mathcal{H})$ and*

$$A = \{(x_1, \dots, x_n) | x_1 \in *D_1 \wedge \dots \wedge x_n \in *D_n \wedge *\Phi(x_1, \dots, x_n)\}.$$

Proof. (i) Let $\Phi(x)$ be any bound formula in which all constants are members of $C(\mathcal{H})$. Let $B = \{x | x \in D \wedge \Phi(x)\}$, where D and all constants in $\Phi(x)$ represent members of $\mathcal{H}$. In our general set theory the set A exists. Let $a_1, \dots, a_m$ be the list of all constants in $\Phi(x)$. Then there is a finite set $\{X_i\}$ such that each a_i is a member of some X_i , and some X_p , $p \geq 1$ such that $D \in X_p$ and since $B \subset D$ then from Proposition 1 (vi) $B \in X_p$. Observe that this also implies that the formula $\Phi(x)$ is defined within the superstructure $\mathcal{H}$ (i.e. the vocabulary of $\Phi(x)$ is interpretable in $\mathcal{H}$). Even though once it is understood that $\Phi(x)$ has meaning in $\mathcal{H}$ the following expanded form is not considered necessary, it is presented in preparation for the proof of Theorem 4.2.2. The sentence

$$a_1 \in X_1 \wedge \dots \wedge a_m \in X_m \wedge D \in X_p \wedge B \in X_p \wedge \forall x(x \in B \leftrightarrow (x \in D) \wedge \Phi(x))$$

holds in $\mathcal{M}$ if and only if the $*$ -transform holds in $^*\mathcal{M}$. Consequently, $A = ^*B = \{x | x \in ^*D \wedge ^*\Phi(x)\}$.

Conversely, let $A = ^*B$. Simply consider the formula $\Phi(x) = x \in B$. Then $^*B \in ^*X_p$ and $^*B$ be a set. If $x \in ^*B$, then $x \in ^*X_0 \cup ^*X_{p-1}$. Further, $X_0 \cup X_{p-1} \in X_p \Rightarrow ^*X_0 \cup ^*X_{p-1} \in ^*X_p$. Thus $A = ^*B = \{x | x \in ^*X_0 \cup ^*X_{p-1} \wedge ^*\Phi(x)\}$ from the first part.

(ii) This proof is but a simple modification of part (i). Consider the formula $\Phi(x_1, \dots, x_n)$, $n > 1$ in n free variables and assume as before that the non-quantifier bounding constants represent members of $\mathcal{H}$. Adjoin the fact that each $D_i \in X_{p(i)}$ and that there exists some X_k such that $B \in X_k$, $k \geq 1$ to the other facts. Then the sentence

$$a_1 \in X_1 \wedge \dots \wedge a_m \in X_m \wedge D_1 \in X_{p(1)} \wedge \dots \wedge D_n \in X_{p(n)} \wedge B \in X_k \wedge$$

$$\forall x_1 \dots \forall x_n (x_1 \in D_1 \wedge \dots \wedge x_n \in D_n \rightarrow \forall z (z \in B \leftrightarrow$$

$$z = (x_1, \dots, x_n) \wedge \Phi(x_1, \dots, x_n)))$$

holds in $\mathcal{M}$ if and only if the $*$ -transform holds in $^*\mathcal{M}$. Consequently, $A = ^*B = \{(x_1, \dots, x_n) | x_1 \in ^*D_1 \wedge \dots \wedge x_n \in ^*D_n \wedge ^*\Phi(x_1, \dots, x_n)\}$.

The converse follows in the same manner as in part (i) and the proof is complete.

Corollary 4.2.11.1 *For the Φ in Theorem 4.2.1, if $B = \{x | x \in D \wedge \Phi(x)\}$ [resp. $\{(x_1, \dots, x_n) | x_1 \in D_1 \wedge \dots \wedge x_n \in D_n \wedge \Phi(x_1, \dots, x_n)\}$], then $^*B = \{x | x \in ^*D \wedge ^*\Phi(x)\}$ [resp. $\{(x_1, \dots, x_n) | x_1 \in ^*D_1 \wedge \dots \wedge x_n \in ^*D_n \wedge ^*\Phi(x_1, \dots, x_n)\}$].*

Theorem 4.2.2. (The Internal Definition Principle)

(i) *A set A in our set theory is an internal set if and only if there exists some internal set D and an internal bound formula $\Phi(x)$ in one free variable where each constant in $\Phi(x)$ is a member of $C(^*\mathcal{H})$ and*

$$A = \{x | x \in D \wedge \Phi(x)\}.$$

(ii) *A set A is an internal n -ary relation ($n > 1$) if and only if there exist n internal sets $D_1, \dots, D_n$ and a internal bound formula $\Phi(x_1, \dots, x_n)$ in n free variables where each constant in $\Phi(x_1, \dots, x_n)$ is a member of $C(^*\mathcal{H})$ and*

$$A = \{(x_1, \dots, x_n) | x_1 \in D_1 \wedge \dots \wedge x_n \in D_n \wedge \Phi(x_1, \dots, x_n)\}.$$

Proof. (i) Let $a_1, \dots, a_m$ be the list of all constants in $\Phi(x)$. Then each a_i is a member of some $^*X_i$ and there exists X_p such that $D \in ^*X_p$. Now replace every distinct a_i at each occurrence with a distinct variable y_i yielding the $n+1$ placed formula $\Phi(x, y_1, \dots, y_m)$. The following sentence

$$\forall y_1 \dots \forall y_m \forall z (y_1 \in X_1 \wedge \dots \wedge y_m \in X_m \wedge z \in X_p \rightarrow \exists u (u \in X_p \wedge$$

$$\forall x (x \in X_0 \cup X_{p-1} \rightarrow (x \in u \leftrightarrow x \in z \wedge \Phi(x, y_1, \dots, y_m))))))$$

holds in $\mathcal{M}$ if and only if it holds in $^*\mathcal{M}$. Consequently, by considering the specific internal objects denoted by a_i and by D . Then the corresponding internal u that exists in $^*X_p$ is precisely the set $\{x | x \in D \wedge \Phi(x)\}$.

The converse follows from Proposition 2 in the same manner as in the proof of part (i) of Theorem 4.2.1.

(ii) For this part follow the same procedure as in the second part of the proof of Theorem 4.2.1. Modify the appropriate subformula of the formal sentence that appears in the proof of this theorem part (i) in the same manner as the sentence was modified in the proof of Theorem 4.2.1 part (ii). With this the proof is complete.

Theorem 4.3.1. *Let $n > 1$. Then for each internal $R \subset {}^*A_1 \times \cdots \times {}^*A_n$ and for each i , $1 \leq i \leq n$ it follows that ${}^*P_i(R) = P_i(R)$, where $P_i(R)$ is an internal subset of ${}^*A_i$.*

Proof. Observe that for each i , $1 \leq i \leq n$ the i th projection may be considered a map $P_i: \mathcal{P}(A_1 \times \cdots \times A_n) \rightarrow \mathcal{P}(A_i)$. Thus ${}^*P_i$ is a map from the internal subsets of ${}^*A_1 \times \cdots \times {}^*A_n$ to the internal subsets of ${}^*A_i$. Hence, if internal $Q \subset {}^*A_1 \times \cdots \times {}^*A_n$, then $({}^*P_i)(Q) \subset {}^*A_i$ is internal. Further, in general, there exists some X_p such that if $B \subset A_1 \times \cdots \times A_n$, then $B \in X_p$. The remainder of this proof is by induction on the number of Cartesian products, m , as was done in the proof of Theorem 3.1.4. Let $m = 1$. The following sentence

$$\begin{aligned} & \forall z(z \in \mathcal{P}(A_1 \times A_2) \rightarrow \forall x(x \in A_1 \rightarrow \\ & (x \in P_1(z) \leftrightarrow \exists y(y \in A_2 \wedge (x, y) \in z)))) \end{aligned}$$

holds in $\mathcal{M}$ if and only if it holds in ${}^*\mathcal{M}$. Suppose that internal $Q \subset {}^*A_1 \times {}^*A_2$. Let $q \in ({}^*P_1)(Q)$. Then $q \in {}^*A_1 \Rightarrow \exists s \in {}^*A_2$ such that $(q, s) \in Q \Rightarrow ({}^*P_1)(Q) \subset P_1(Q)$, where P_1 is a set-theoretic projection for n -ary relations $Q \in {}^*\mathcal{H}$. On the other hand, let $q \in P_1(Q)$. Then $q \in {}^*A_1$ and $\exists s \in {}^*A_2$ such that $(q, s) \in Q$. This yields, from the * -transform, that $q \in ({}^*P_1)(Q)$ and, hence, $({}^*P_1)(Q) = P_1(Q)$. The remainder of this proof follows in exactly the same manner, appropriately modified, as does the last part of the proof of Theorem 3.1.4.

Theorem 4.3.2. *Let $F(B)$ be the set of all finite subsets of B . Then nonempty internal $A \subset {}^*B$ is hyper finite if and only if $A \in ({}^*F)(B)$, where if there can be any confusion, we continue to denote the * -transform of such operators as $({}^*F)$.*

Proof. As mentioned, the notion used for finite sets is a first-order statement in our set-theory with “finitely” many $\forall$ symbols and constants. This is not the mapping notion used for the definition of hyper finite. However, it is a simple exercise to show by induction that since $\mathbb{N} \subset X_0$ that for any nonempty finite set $A \in \mathcal{H}$, there exists in $\mathcal{H}$ a bijection $f: [0, n] \rightarrow A$. Then f corresponds within our set-theory to a set that can be characterizes as having the same “number” of $\forall$ symbols and constants when the range is completely described. Consequently, whenever it is necessary, we characterize finite sets by such mappings. Next, it is necessary to establish by induction that the mapping concept used for the hyperfinite set definition is equivalent to the * -transform of the standard set-theoretic nonempty finite set mapping correspondence. We show that $\forall n \in \mathbb{N}$ if $g \in B^{[0, n]}$ and $g[[0, n]] = A$, then $\exists k_n \in \mathbb{N}$ and an injection $f \in B^{[0, k_n]}$ such that $f[[0, k_n]] = A$. For $n = 0$ then g itself is such an injection. Suppose the hypothesis holds for an arbitrary $n \in \mathbb{N}$. Let $g \in B^{[0, n+1]}$ and $g[[0, n+1]] = A'$. Our set theory states that there exists a map $g_1 = g|_{[0, n]}$ and $g_1[[0, n]] \subset A'$. By induction $\exists k_n \in \mathbb{N}$ and an injection $f_1 \in B^{[0, k_n]}$ such that $f_1[[0, k_n]] = g_1[[0, n]]$. If $g_1[[0, n]] = A'$, then f_1 suffices. If $g_1[[0, n]] \neq A'$, then $g(n+1) \notin g_1[[0, n]] = f_1[[0, k_n]]$ and $A' = g_1[[0, n]] \cup \{g(n+1)\}$. In this case, simply define $f = \{(k_n + 1, g(n+1))\} \cup f_1$. Then f is an injection defined on $[0, k_n + 1]$ onto A' . The hypothesis holds by induction. Consequently, if $\exists n \in \mathbb{N}$ such that $g \in B^{[0, n]}$, then $g[[0, n]] = A$ is finite in the standard set-theoretic mapping sense. The converse is obvious and this implies that our slightly modified mapping definition is equivalent to the * -transform of the standard mapping “definition” for nonempty finite sets.

For a given entity B there exists some X_p such that $F(B) \in X_p$. Let $\mathcal{A} = \mathcal{P}(\mathbb{N})$ and $\mathcal{B} = \{B\}$. Then as discussed just prior to example 4.3.1 there exists some $q \in \mathbb{N}$ such that for each $A \in \mathcal{A}$ and $B \in \mathcal{B}$, $B^A \in X_q$ and, hence, since the members of B^A are not individuals each $f \in B^A$ is a member of X_{q-1} . From the above proof it follows that the follow sentence

$$\forall x(x \in X_p \rightarrow (x \in F(B) \leftrightarrow \exists y \exists z(y \in \mathbb{N} \wedge z \in X_{q-1} \wedge$$

$$z \in B^{[0,y]} \wedge z[[0,y]] = x)))$$

holds in $\mathcal{M}$ if and only if it holds in $^*\mathcal{M}$. The result follows by *-transfer.

Theorem 4.3.3 *Any nonempty finite set of internal individuals or entities is internal and hyperfinite.*

Proof. This follows from Proposition 2 (iii) and Theorem 4.3.2.

Theorem 4.3.4. *Let $A \in \mathfrak{R}$ and assume that A is infinite. Then there exists a hyperfinite set F such that $F \neq A$, $F \neq ^*A$ and $A \subset F \subset ^*A$.*

Proof. Let infinite set $A \in X_p$. Consider the standard binary relation $R = \{(x, y) | x \in y \wedge y \in F(A) \wedge x \in A\}$. Observe that the domain of R is the set A . Consider $\{(x_1, y_1), \dots, (x_n, y_n)\} \subset R$. Then the set $y_1 \cup \dots \cup y_n \in F(A)$ and $\{(x_1, y), \dots, (x_n, y)\} \subset A$. Thus R is concurrent on A . Thus there exists some $F \in ^*F(A)$ such that $\forall x \in {}^\sigma A$, $x \in F$. Since $A \in \mathfrak{R}$ then ${}^\sigma A = A \subset F$ and by *-transfer $F \subset ^*A$. Since A is not internal and F is internal then $F \neq A$. Further, if $F = ^*A$, then the sentence

$$\exists x(x \in F(A) \wedge x = A)$$

holds in $\mathcal{M}$ since it holds in $^*\mathcal{M}$. This would contradict the fact that A is not finite and the proof is complete.

Theorem 4.3.5. *An infinite series, $\sum_{i=0}^\infty a_i$, converges to $r \in \mathbb{R}$ if and only if for each $\Gamma \in \mathbb{N}_\infty$ it follows that $\sum_{i=0}^\Gamma a_i \in \mu(r)$.*

Proof. This is an immediate consequence of Theorem 3.2.1.

Theorem 4.4.1. *Definition 4.4.1 for continuity and uniform continuity is equivalent to the $\delta - \epsilon$ definition.*

Proof. (i) (Continuity). Assume Definition 4.4.1. Thus for nonzero $n, m \in \mathbb{N}$, nonempty $A \subset \mathbb{R}^n$, the mapping $f: A \rightarrow \mathbb{R}^m$ has the property at $p \in A$ that $^*f[\mu(p) \cap ^*A] \subset \mu(f(p))$. Let $r \in \mathbb{R}^+$. If p is an isolated point, then $\mu(p) \cap ^*A = \{p\}$ and $^*f[\mu(p) \cap ^*A] = \{p\} \subset \mu(f(p))$. Suppose that p is not isolated. Then $\exists 0 \neq \epsilon \in \mu(0)$ and $p + \epsilon = q \in \mu(p) \cap ^*A$. Moreover, $0 < \|q - p\| < \|\epsilon\|$ and if $s \in ^*A$ and $0 < \|s - p\| < \|\epsilon\|$, then $s \in \mu(p) \Rightarrow ^*f(s) \in \mu(f(p))$. Combining these two cases it follows by *-transfer that the sentence

$$\exists x(x \in \mathbb{R}^+ \wedge \forall y(y \in A \wedge 0 \leq \|p - y\| < x \rightarrow \|f(y) - f(p)\| < r))$$

holds in $\mathcal{M}$. Hence, f is continuous at $p \in A$.

For the converse, let $r \in \mathbb{R}^+$. Then we know that there exists some $r_0 \in \mathbb{R}^+$ such that

$$\forall x(x \in A \wedge 0 \leq \|x - p\| < r_0 \rightarrow \|f(x) - f(p)\| < r)$$

holds in $\mathcal{M}$. Hence the *-transfer holds in $^*\mathcal{M}$. Now let $q \in \mu(p) \cap ^*A$. Then $\|q - p\| < r_0$ for any $r_0 \in \mathbb{R}^+$. Thus for any $r \in \mathbb{R}^+$, $\|f(q) - f(p)\| < r$. Since r is an arbitrary positive real number then it follows that for any $q \in \mu(p) \cap ^*A$, $^*f(q) - f(p) \in \mu(0)$ and $\Rightarrow ^*f[\mu(p) \cap ^*A] \subset \mu(f(p))$.

(ii) (Uniform continuity). Assume Definition 4.4.1. Thus for the map $f: A \rightarrow \mathbb{R}^m$ and if $p, q \in ^*A$, $p - q \in \mu(0)$, then $^*f(p) - ^*f(q) \in \mu(0)$. Suppose that arbitrary $r \in \mathbb{R}^+$. We know that

there exists a nonzero $\epsilon \in \mu(0)$ and for any $p, q \in {}^*A$ such that $\|p - q\| < \epsilon$, ${}^*f(p) - {}^*f(q) \in \mu(0) \Rightarrow \|{}^*f(p) - {}^*f(q)\| < r$. Hence, the sentence

$$\exists x(x \in \mathbb{R}^+ \wedge \forall y \forall z(y \in A \wedge z \in A \wedge \|y - z\| < x \rightarrow \|f(y) - f(z)\| < r))$$

holds in $\mathcal{M}$. Thus f is uniformly continuous on A .

For the converse, assume uniform continuity and let arbitrary $r \in \mathbb{R}^+$. Then we know that there exists some $r_0 \in \mathbb{R}^+$ such that for each x and each y such that $x, y \in A$ and $\|x - y\| < r_0$ then $\|f(x) - f(y)\| < r$. By $*$ -transfer we have that for any $p, q \in {}^*A$ and $p - q \in \mu(0)$ it follows that $\|{}^*f(p) - {}^*f(q)\| < r$. Since r is an arbitrary nonzero positive real number then ${}^*f(p) - {}^*f(q) \in \mu(0)$. This completes the proof.

Theorem 4.4.2. *Definition 4.4.2 for compactness is equivalent to the standard definition utilizing open covers.*

Proof. Even though there are proofs of this that, in this case, do not require the enlargement property (see Herrmann [1978]) the enlargement property leads to a much shorter demonstration.

Suppose A is compact and that there is a $q \in {}^*A$ such that $q \notin \bigcup\{\mu(r) \mid r \in A\}$. We use the characterization that $\mu(r) = \bigcap\{{}^*G \mid r \in G \in \mathcal{T}\}$, where $\mathcal{T}$ is the set of all open subsets of $\mathbb{R}^n$. Thus for each $r \in A$ there exists some $G_r \in \mathcal{T}$ such that $q \notin {}^*G_r$. Since $\{G_r \mid r \in A\}$ is an open cover for A then there exists a finite $\{G_1, \dots, G_n\} \subset \{G_r \mid r \in A\}$ such that $A \subset G_1 \cup \dots \cup G_n$. Hence, ${}^*A \subset {}^*G_1 \cup \dots \cup {}^*G_n \Rightarrow \exists i, 1 \leq i \leq n$ such that $q \in {}^*G_i \in \{G_r \mid r \in A\}$. But this contradiction implies that standard compactness yields Definition 4.4.2.

Conversely, assume that A is not compact and let $\mathcal{G}$ be an open cover of A that does not contain a finite subcover. As usual all of these objects are members of $\mathcal{H}$. Consider the binary relation

$$R = \{(x, y) \mid x \in \mathcal{G} \wedge y \in A \wedge y \notin x\}.$$

Let $\{(x_1, y_1), \dots, (x_n, y_n)\} \subset R$. Then there exists some $q \in A$ such that $q \notin x_i$ for any $i, 1 \leq i \leq n$. Hence, R is concurrent and the domain of $R = \mathcal{G}$. Thus there exists some $b \in {}^*A$ such that $(x, b) \in {}^*R$ for each $x \in \mathcal{G}$. But $x \in \mathcal{G} \leftrightarrow x = {}^*G, G \in \mathcal{G}$. Consequently, since $\mathcal{G}$ contains at least one open neighborhood for each $r \in A$ then $b \notin \bigcup\{\mu(r) \mid r \in A\}$ and this completes the proof.

NOTATION

Notice that the notation has been generalized slightly. So that there was no possibility of confusion when not carefully read, hyper-function notation such as ${}^*f$ has been presented in two forms. The $({}^*f)(A)$, to identify the function ${}^*f$ and ${}^*f(A)$, meaning ${}^*(f(A))$ when A is a standard object. The notational form $({}^*f)(A)$ is unnecessary since which meaning applies depends upon the argument. In this example, when A is standard, one needs only write ${}^*f(A)$. It is obvious that this notation has only the one possible meaning ${}^*(f(A))$ for in this form the hyper-function is also being displayed since ${}^*(f(A)) = {}^*f({}^*A)$. On the other hand, if A is simply specified as an internal object, then the notation ${}^*f(A)$ can only mean $({}^*f)(A)$. The notations are equivalent if $A = {}^*B$. In some cases under our identification process, the notations are also equivalent where we let ${}^*A = A$, where A is identified with the standard object via the constants sequences of atoms (urelements).

Appendix For Chapter 5.

In all that follows the set of natural numbers, $\mathbb{N}$, is assumed to contain zero and all functions from a domain that is a subset of $\mathbb{R}^n$ into $\mathbb{R}$ are bounded.

For an n -dimensional space $n \geq 1$, the closed set $R = [a_1, b_1] \times \cdots \times [a_n, b_n]$, $a_i < b_i$, $1 \leq i \leq n$ is called a **rectangle**. Of course, if $n = 1$, then a “rectangle” is but a closed interval. As usual, consider for each $[a_i, b_i]$ a **partition** P_i as a finite set of members of $[a_i, b_i]$ such that $a_i, b_i \in P_i$ and where P_i is considered as ordered. This is often explicitly written as $P_i = \{x_{i0}, \dots, x_{ik}\}$, $a_i = x_{i0} < x_{i1} < \cdots < x_{ik} = b_i$. This determines the closed one-dimensional subintervals $[x_{i(p-1)}, x_{ip}]$, $1 \leq p \leq k$. In brief, this process obtains a partition $P = P_1 \times \cdots \times P_n$ of R and a finite collection of closed n -dimensional subrectangles S obtained by considering $([x_{10}, x_{11}] \cup \cdots \cup [x_{1k-1}, x_{1k}]) \times \cdots \times ([x_{n0}, x_{n1}] \cup \cdots \cup [x_{nm-1}, x_{nm}])$. Each S has a measure, $v(S) \in \mathbb{R}$, assigned to it which is intuitively the product of the lengths of the sides. For simplicity of notion the definition of the measure $v(S)$ is left intuitively understood.

Probably the simplest partition to consider would be the one termed a **simple** partition. These are formed by selecting n nonzero natural numbers $m_1, \dots, m_n$ and dividing each interval $[a_i, b_i]$ into an equal length partition by adding to each successive partition point the number $(b_i - a_i)/(m_i)$. This concept is extended to the nonstandard world by selecting n infinite natural numbers $\Gamma_1, \dots, \Gamma_n$ and generating for each interval $[a_i, b_i]$ an internal hyperfinite partition, P_i , each subinterval of which has positive infinitesimal length $(b_i - a_i)/(\Gamma_i) = dx_i$. Then the partition $P = P_1 \times \cdots \times P_n$ is a **simple fine partition** of R . Such a partition yields an internal set of hyperrectangles R_q such that $*v(R_i) = dx_1 \cdots dx_n = dX \in \mu(0)^+$. You could be much more general and consider the **fine** partitions which are internal collections of hyperfinitely many members of $*[a_i, b_i]$ such that the length of any subinterval is an infinitesimal. Let P be a partition of the rectangle R and assume that P determines the set of subrectangles $\{R_q | 1 \leq q \leq m\}$. An **intermediate partition**, Q , is any finite sequence of vectors $\{\vec{v}_q\}$, where $\vec{v}_q \in R_q$ for each q such that $1 \leq q \leq m$. Let $\mathcal{P}$ be the set of all simple partitions of R and $\mathcal{P}_S$ any nonempty subset of $\mathcal{P}$. Then there exists a mapping Q on the set $\mathcal{P}_S$ such that for each $P \in \mathcal{P}_S$ the image $Q(P)$ is the set of all intermediate partitions for P . From a notational view point subrectangles are denoted by various symbol strings. In particular, S , R_q , R_S .

For any partition P of R let $\mathcal{S}(P)$ denote the set of all subrectangles generated by P . If $S \in \mathcal{S}(P)$, then let $\|S\|$ denote the length of the diagonal of the subrectangle S . As usual, define the **mesh**(P) = $\|P\| = \max\{\|S\| | S \in \mathcal{S}(P)\}$. Now let $L(f, P) = \sum_{S \in \mathcal{S}(P)} m_S v(S)$ be an **lower sum** and $U(f, P) = \sum_{S \in \mathcal{S}(P)} M_S v(S)$ be an **upper sum**, where $m_S = \inf\{f(\vec{x}) | \vec{x} \in S\}$ and $M_S = \sup\{f(\vec{x}) | \vec{x} \in S\}$. As is well-known, for any set of partitions of R the set of lower and upper sums is a bounded set. Further, let $L(f) = \sup\{L(f, P) | P \text{ is any partition of } R\}$ and $U(f) = \inf\{U(f, P) | P \text{ is any partition of } R\}$. The function f is **Darboux** integrable if $L(f) = U(f)$ and the value of this integral is $D \int_R f(\vec{x}) dX = L(f) = U(f)$. As is well-known f is Darboux integrable if and only if f is Riemann integrable in the sense of Riemann Sums and has the same value. Now the concept of the mesh extends to the $*$ -mesh and for any internal partition Q of $*R$ the $*\|Q\|$ exists by $*$ -transfer of the finitary statement dealing with maximum values that exist within a finite set of real numbers. Also, Q is a fine partition if and only if $*\|Q\| \in \mu(0)$.

Notice that if $\mathcal{P}_S$ is any nonempty subset of the set of all partitions of R , then the lower and upper sums we may consider maps $L(f, \bullet): \mathcal{P}_S \rightarrow \mathbb{R}$ and $U(f, \bullet): \mathcal{P}_S \rightarrow \mathbb{R}$. These maps extend to the maps $*L(f, \bullet): *\mathcal{P}_S \rightarrow *\mathbb{R}$ and $*U(f, \bullet): *\mathcal{P}_S \rightarrow *\mathbb{R}$.

Definition A5.1. Let bounded $f: R \rightarrow \mathbb{R}$ and $\mathcal{P}_I$ be any nonempty set of hyperfinite partitions of ${}^*R$ such that ${}^*\|P\| \in \mu(0)$ for each $P \in \mathcal{P}_I$. Then f is said to be **Z-integrable** for $\mathcal{P}_I$ if there exists some $P \in \mathcal{P}_I$ such that

$$\text{st}({}^*L(f, P)) = \text{st}({}^*U(f, P)) = Z \int_R f(\vec{x}) {}^*\|P\|.$$

Proposition 5.1. Suppose that bounded $f: R \rightarrow \mathbb{R}$. Let $Q = Q_1 \times \cdots \times Q_n$ be a hyperfinite partition of ${}^*R$, where each Q_i is a hyperfinite partition of $[a_i, b_i]$, for $1 \leq i \leq n$. If ${}^*\text{mesh}(Q_i) \in \mu(0)$, then

$$0 \leq L(f) - {}^*L(f, Q) \in \mu(0),$$

$$0 \leq {}^*U(f, Q) - U(f) \in \mu(0).$$

Proof. Consider $P_i = \{a_i = x_{i0}, \dots, x_{i(m+1)} = b_i\}$ a partition of $[a_i, b_i]$. Then since it is a finite set of real numbers it follows that ${}^*P_i = P_i$. Now let $Q = Q_1 \times \cdots \times Q_n$ be a hyperfinite partition of ${}^*R$, where each Q_i is hyperfinite and ${}^*\text{mesh}(Q_i) \in \mu(0)$. [Note that ${}^*\text{mesh}(Q_i) \in \mu(0)$ for $1 \leq i \leq n$ if and only if ${}^*\text{mesh}(Q) \in \mu(0)$.] Let $Q_i = \{a_i = y_{i0}, \dots, y_{i(\nu+1)} = b_i\}$. For each $j = 1, \dots, m$, we know there are finitely many members of Q_i such that $y_{ik(j)} \leq x_{ij} < y_{i(k(j)+1)}$. This holds for each $i = 1, \dots, n$. Thus there is a finite set $\mathcal{R}$ of infinitesimal subrectangles R_q such that $\vec{x} \in P$ if and only if there exists some $R_q \in \mathcal{R}$ such that $\vec{x} \in R_q$. Further, if $R_q \in \mathcal{R}$, then ${}^*v(R_q) \in \mu(0)$. Since $\mathcal{R}$ is also a hyperfinite set then by * -transfer we can rearrange the hyperfinite sum $\sum_{S \in \mathcal{S}(Q)} m_S v(S)$ as follows:

$${}^*L(f, Q) = \sum_{S \in \mathcal{R}} m_S {}^*v(S) + \sum_{S \in (\mathcal{S}(Q) - \mathcal{R})} m_S {}^*v(S).$$

This all implies since a finite sum of infinitesimals is infinitesimal that

$${}^*L(f, Q) \approx \sum_{S \in (\mathcal{S}(Q) - \mathcal{R})} m_S {}^*v(S).$$

Moreover, with respect to the common refinement $K = P \cup Q$ the partition points in P only finitely partition the infinitesimal subrectangles $S \in \mathcal{R}$. Hence, in a similar manner the hyperfinite sum ${}^*L(f, K)$ can be rearranged so that

$${}^*L(f, K) \approx \sum_{S \in (\mathcal{S}(Q) - \mathcal{R})} m_S {}^*v(S).$$

Thus ${}^*L(f, K) \approx {}^*L(f, Q)$.

Now K is a hyperfinite partition of R that * -refines P . By * -transfer of the elementary properties of lower sums and partitions it follows that ${}^*L(f, K) \geq L(f, P)$. Let $r \in \mathbb{R}^+$. Then there is a standard partition P_1 such that $L(f, P_1) \geq L(f) - r$. From this we have that

$$L(f) \geq {}^*L(f, Q) \approx {}^*L(f, K) \geq L(f, P_1) \geq L(f) - r.$$

Since r is arbitrary then taking the standard part operator we have that $\text{st}({}^*L(f, Q)) = L(f)$. Thus $0 \leq L(f) - {}^*L(f, Q) \in \mu(0)$. In similar manner the second conclusion follows and this completes the proof.

Proposition 5.2.

- (i) A bounded function $f: R \rightarrow \mathbb{R}$ is Darboux integrable iff it is Z-integrable for some $\mathcal{P}_I$ if and only if it is Z-integrable for all $\mathcal{P}_I$.
- (ii) If bounded $f: R \rightarrow \mathbb{R}$ is Darboux integrable or Z-integrable for $\mathcal{P}_I$, then

$$D \int_R f(\vec{x}) dX = Z \int_R f(\vec{x}) * \|P\|.$$

Proof. These results are indeed immediate from Proposition 5.1 and application of the standard part operator.

Proposition 5.3. Let bounded $f: R \rightarrow \mathbb{R}$ and $\mathcal{P}_I$ be any nonempty set of hyperfinite partitions of $*R$ such that $*\|P\| \in \mu(0)$ for each $P \in \mathcal{P}_I$. If f is Z-integrable and $P, Q \in \mathcal{P}_I$, then

$$\text{st}(*L(f, P)) = \text{st}(*L(f, Q)) = Z \int_R f(\vec{x}) * \|P\| = Z \int_R f(\vec{x}) * \|Q\|.$$

Proof. Since f is Darboux integrable then once again application of Proposition 5.1 and the standard part operator completes the proof.

Theorem 5.1.1. A bounded function $f: R \rightarrow \mathbb{R}$ is integrable if and only if it is integrable in the sense of Darboux and Riemann Sums.

Proof. Let $\mathcal{P}$ be the set of all simple partitions of R . Then there exists a mapping $\mathcal{Q}$ on the set $\mathcal{P}$ such that for each $P \in \mathcal{P}$ the image $\mathcal{Q}(P)$ is the set of all intermediate partitions for P . For each $P \in \mathcal{P}$ and each $Q \in \mathcal{Q}(P)$ let

$$S(f, P, Q) = \sum_{S \in \mathcal{S}(P), \vec{x} \in S \cap Q} f(\vec{x}) v(S).$$

First, assume that f is integrable as defined in by Definition 5.1.1. Then we know that there is some $r \in \mathbb{R}$, some $P' \in *\mathcal{P}$ such that for each $Q' \in *\mathcal{Q}(P')$, $*S(f, P', Q') \in \mu(r)$. Hence for arbitrary $A \in \mathbb{R}^+$

$$r - A < *S(f, P', Q') < r + A.$$

By *-transfer, there exists a standard simple partition $P \in \mathcal{P}$ such that for each $Q \in \mathcal{Q}(P)$ it follows that

$$r - A < S(f, P, Q) < r + A.$$

From this it follows that $r - A \leq L(f, P) \leq U(f, P) \leq r + A$. Thus $r - A \leq L(f) \leq U(f) \leq r + A$, $\Rightarrow |U(f) - r| \leq A$ and $|L(f) - r| \leq A$. Since A is arbitrary then this implies that $U(f) = L(f) = r$. Consequently, f is Darboux integrable and as is well-known f is Riemann integrable in the sense of Riemann Sums.

Conversely, assuming that f is Darboux (or Riemann) integrable then f is Z-integrable for, $\mathcal{P}_I$, the set of all fine partitions of $*R$. Thus $P' \in \mathcal{P}_I$ and $*\|P'\| = dX$, then

$$\text{st}(*L(f, P')) = \text{st}(*U(f, P')) = Z \int_R f(\vec{x}) dX = D \int_R f(\vec{x}) dX.$$

But from *-transfer, for each $Q' \in *\mathcal{Q}(P')$,

$$*L(f, P') \leq *S(f, P', Q') \leq *U(f, P').$$

The result follows by taking the standard part operator.

Theorem 5.1.2. *If bounded $f: R \rightarrow \mathbb{R}$ is integrable, then there exists a unique $r \in \mathbb{R}$ such that for every fine partition $P = \{\vec{x}_0, \dots, \vec{x}_\Omega\}$, $\Omega \in \mathbb{N}_\infty$ and every internal intermediate partition $Q = \{\vec{v}_q\}$, $1 \leq q \leq \Gamma \in \mathbb{N}_\infty$ it follows that*

$$\sum_{k=1}^{\Gamma} {}^*f(\vec{v}_q) {}^*v(R_q) \in \mu(r).$$

Proof. Since f is Darboux integrable then it is Z-integrable. Let $\mathcal{P}_I$ be the set of all fine partitions of R . Then every member of $\mathcal{P}_I$ is internal. Result follows by repeating the proof of the converse of Theorem 5.1.1 for $\mathcal{P}_I$.

Obviously, the restriction of fine partitions to the simple fine partitions is not necessary since the value of the integral defined in Chapter 5, as well as the equivalent Darboux or Riemann integral, is independent of the choice of the fine partition. Clearly, the simplicity of the simple fine partitions is often useful in applications and the customary integral notation of dX implies that each infinitesimal subrectangle has * -measure $= dX$. However, for much that follows $\mathcal{P}$ is often assumed only to be an infinite set of partitions $P = P_1 \times \dots \times P_n$ and in this case, letting $\mathcal{S}(P)$ the set of all subrectangles of $R \subset \mathbb{R}^n$ generated by $P \in \mathcal{P}$, we define the set $\mathcal{C}'_{PSR} = \{S | \exists P(P \in \mathcal{P} \wedge S \in \mathcal{S}(P))\}$. Extend the definition of simply additive to $\mathcal{P}$.

Proposition 5.4. *Let $f: R \rightarrow \mathbb{R}$. Let $\mathcal{P}$ be any infinite set of partitions (including the trivial one $\{R\}$) of R , $B: \mathcal{C}'_{PSR} \rightarrow \mathbb{R}$, and B simply additive on each $P \in \mathcal{P}$. Suppose that there exists a fine partition $Q \in {}^*\mathcal{P}$ (i.e. ${}^*\|Q\| \in \mu(0)$) such that for every $S \in {}^*\mathcal{S}(Q)$ there exists some $\vec{p} \in S$ such that*

$${}^*B(S)/({}^*v(S)) \approx {}^*f(\vec{p}),$$

then for any $r \in \mathbb{R}^+$

$$-rv(R) + {}^*L(f, Q) < B(R) < {}^*U(f, Q) + rv(R).$$

Proof. Let $r \in \mathbb{R}^+$ and assume that for the fine partition $Q \in {}^*\mathcal{P}$

$${}^*(B(R)) = {}^*B({}^*R) \geq {}^*U(f + r, Q).$$

First, we make the following standard observations. Let $P \in \mathcal{P}$ be any standard partition of R . Assume that $B(R) \geq U(f + r, P)$. The simple additivity of B yields that there must exist some $S \in \mathcal{S}(P)$ such that

$$B(S) \geq M_S v(S), \quad M_S = \sup\{f(x) + r | x \in S\}.$$

By * -transfer, there exists some $T \in {}^*\mathcal{S}(Q)$ such that

$${}^*B(T) \geq M_T {}^*v(T), \quad M_T = {}^*\sup\{{}^*f(\vec{x}) + r | x \in T\}.$$

Hence for each $\vec{p} \in T$

$${}^*B(T) \geq ({}^*f(\vec{p}) + r) {}^*v(T).$$

Therefore since ${}^*v(T) > 0$ this implies that for each $\vec{p} \in T$,

$${}^*B(T)/({}^*v(T)) - {}^*f(p) \geq r.$$

This, however, contradicts the hypotheses. Thus ${}^*(B(R)) = {}^*B({}^*R) = B(R) < {}^*U(f + r, Q) = {}^*U(f, Q) + rv(R)$.

In like manner, it follows that $-rv(R) + {}^*L(f, Q) < B(R)$.

Proposition 5.5. *Let $f: R \rightarrow \mathbb{R}$. Let $\mathcal{P}$ be any infinite set of partitions (including the trivial one $\{R\}$) of R , $B: \mathcal{C}'_{PSR} \rightarrow R$, and B simply additive on each $P \in \mathcal{P}$. Suppose that there exists a fine partition $Q \in {}^*\mathcal{P}$ such that for every $S \in {}^*\mathcal{S}(Q)$ there exists some $\vec{p} \in S$ such that*

$${}^*B(S)/({}^*v(S)) \approx {}^*f(\vec{p}),$$

then f is Z -integrable for $\{Q\}$ and

$$B(R) = Z \int_R f(\vec{x}) {}^*\|Q\|.$$

Proof. From Proposition 5.4 we have that for any $r \in \mathbb{R}^+$, $-rv(R) + {}^*L(f, Q) < B(R) < {}^*U(f, Q) + rv(R)$. Taking the standard part operator and using Proposition 5.1 it follows that

$$-rv(R) + L(f) \leq B(R) \leq U(f) + rv(R).$$

But r an arbitrary positive real number $\Rightarrow$

$$L(f) = U(f) = Z \int_R f(\vec{x}) {}^*\|Q\|.$$

Theorem 5.2.1. (An Infinite Sum Theorem.) *Let bounded $f: R \rightarrow \mathbb{R}$ and simply additive $B: \mathcal{C}_{PSR} \rightarrow \mathbb{R}$. If there exists a simple fine partition $\{R_q | 1 \leq q \leq \Gamma\}$ and for each R_q there exists some $\vec{p} \in R_q$ such that*

$${}^*B(R_q)/dX \approx {}^*f(\vec{p}), \tag{*}$$

then f is integrable and

$$B(R) = \int_R f(\vec{x}) dX.$$

Proof. Simply apply Proposition 5.5.

Definition A5.2. (Jordan-Superneareness). Let J be a nonempty Jordan-measurable subset of $R \subset \mathbb{R}^n$. A map $B: \mathcal{C}_J \rightarrow \mathbb{R}$, where $\mathcal{C}_J$ is a nonempty set of Jordan-measurable subsets of J , is **JORDAN-SUPERNEAR** to bounded $f: J \rightarrow \mathbb{R}$ if for every $K \in {}^*\mathcal{C}_J$ such that ${}^*v(K \cap R_S) \in \mu(0)^+$, where R_S is an infinitesimal subrectangle of ${}^*R$, and every $\vec{p} \in K \cap R_S$ it follows that

$${}^*B(K \cap R_S)/{}^*v(K \cap R_S) \approx {}^*f(\vec{p}), \tag{**}$$

where $v(A)$ is the Jordan content for any Jordan-measurable $A \subset J$ and ${}^*B$ is defined on $K \cap R_S$.

Proposition 5.6. *Let J be a Jordan-measurable subset of $\mathbb{R}^n$. If bounded $f: J \rightarrow \mathbb{R}$ is uniformly continuous on J and $\mathcal{C}_J$ is any nonempty set of connected Jordan-measurable subsets of J , then there exists a map $B: \mathcal{C}_J \rightarrow \mathbb{R}$ that is Jordan-superneare to f .*

Proof. Consider any rectangle R such that $J \subset R$ and as usual define for each $A \in \mathcal{C}_J$,

$$B(A) = \int_A f(\vec{x}) dX = \int_R \hat{f}(\vec{x}) dX.$$

Let $A \in \mathcal{C}_J$ and $R_0 \subset R$ be any subrectangle such that $A \cap R_0 \neq \emptyset$. Then we know that there exists some $\vec{s} \in A \cap R_0$ such that $\int_{A \cap R_0} f(\vec{x}) dX = f(\vec{s})v(A \cap R_0)$. Suppose that $Q \in {}^*\mathcal{C}_J$, ${}^*v(Q \cap R_S) \in \mu(0)^+$ and B is defined on $Q \cap R_S$, where R_S is an infinitesimal subrectangle. Then $Q \cap R_S \neq \emptyset$ and * -transfer implies that there exists some $\vec{q} \in Q \cap R_S$ such that ${}^*B(Q \cap R_S) = {}^*f(\vec{q}){}^*v(Q \cap R_S)$. Consequently,

$$({}^*B(Q \cap R_S) / {}^*v(Q \cap R_S)) \approx {}^*f(\vec{p}).$$

Since $Q \cap R_S \subset R_S$, then $\vec{x}, \vec{y} \in Q \cap R_S \Rightarrow \vec{x} \approx \vec{y}$, and uniform continuity of $f \Rightarrow {}^*f(\vec{x}) \approx {}^*f(\vec{y})$. Thus for any $\vec{p} \in Q \cap R_S$,

$${}^*B(Q \cap R_0) / {}^*v(Q \cap R_S) \approx {}^*f(\vec{q}) \approx {}^*f(\vec{p}).$$

This completes the proof.

In order for Jordan-supernearness to yield continuity on J , it appears necessary to select both of the sets J and $\mathcal{C}_J$ more carefully.

Proposition 5.7. *Let the rectangle $R \subset \mathbb{R}^n$, $f: R \rightarrow \mathbb{R}$ and $\mathcal{C}_{SR} = \{R_q | R_q \text{ is a subrectangle of } R\}$. Suppose that $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ is Jordan-supernear to f , then f is uniformly continuous on R .*

Proof. Let $\vec{p} \neq \vec{q}$, $\vec{p}, \vec{q} \in {}^*R$, $\vec{p} \approx \vec{q}$. Assuming that $\vec{p} = (x_1, \dots, x_n)$, $\vec{q} = (y_1, \dots, y_n)$ then $dx = \max\{|x_i - y_i| | 1 \leq i \leq n\} > 0$. Now letting $R_S = \{(z_1, \dots, z_n) | \forall i (1 \leq i \leq n \rightarrow (x_1 - dx \leq z_i \leq x_n + dx) \wedge (z_i \in {}^*\mathbb{R}))\}$ then for this infinitesimal subrectangle we have that $R_S \in {}^*\mathcal{C}_{SR}$, $\vec{p}, \vec{q} \in R_S$ and ${}^*v(R_S) \in \mu(0)^+$. Jordan-supernearness implies that for each $\vec{x} \in R_S$

$${}^*B(R_S) / {}^*v(R_S) \approx {}^*f(\vec{x}).$$

Since f is bounded on R and $\vec{p}, \vec{q} \in R_S$ then

$${}^*B(R_S) / {}^*v(R_S) \approx {}^*f(\vec{p}) \approx {}^*f(\vec{q}).$$

Consequently f is uniformly continuous on R .

Theorem 5.2.2. *A bounded function $f: R \rightarrow \mathbb{R}$ is continuous if and only if there exists a map $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ that is supernear to f .*

Proof. Obvious from above propositions.

Corollary 5.2.2. *Suppose that $f: R \rightarrow \mathbb{R}$ is continuous. For each $R_S \in \mathcal{C}_{SR}$ define $B(R_S) = \int_{R_S} f(\vec{x}) dX$. Then B is supernear to f .*

Proof. This is established in the proof of Proposition 5.6

Theorem 5.2.3. *Let bounded $f: R \rightarrow \mathbb{R}$. If $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ is supernear to f and simply additive on each simple partition of each R_S , then f is continuous on R and*

$$B(R_S) = \int_{R_S} f(\vec{x}) dX$$

for each $R_S \in \mathcal{C}_{SR}$.

Proof. Proposition 5.7 implies continuity of f implies that the mapping $B': \mathcal{C}_{SR} \rightarrow \mathbb{R}$ defined by $B'(R_S) = \int_{R_S} f(\vec{x}) dX$ is supernear to f . Now f is bounded on R_S . Let $\mathcal{P}_I$ be the set of all simple partitions of R_S . Then B is defined on all subrectangles generated by members of $\mathcal{P}_I$ and simply additive on $\mathcal{S}(P)$ for each $P \in \mathcal{P}_I$. Hence there exists a fine partition $Q \in {}^*\mathcal{P}_I$ and B restricted to $\mathcal{C}_{P(I)SR(S)}$ is also (Jordan) supernear to f . Of course, f is bounded on each R_S . Thus for each R_S , B satisfies the hypotheses of Proposition 5.5. Therefore,

$$B(R_S) = Z \int_{R_S} f(\vec{x}) dX = \int_{R_S} f(\vec{x}) dX = B'(R_S).$$

Corollary 5.2.3.1 *Let bounded $f: R \rightarrow \mathbb{R}$. If $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ is supernear to f and additive on $\mathcal{C}_{SR}$, then f is continuous on R and*

$$B(R_S) = \int_{R_S} f(\vec{x}) dX$$

for each $R_S \in \mathcal{C}_{SR}$.

Corollary 5.2.3.2 *Let bounded $f: R \rightarrow \mathbb{R}$. There exists one and only one map $B: \mathcal{C}_{SR} \rightarrow \mathbb{R}$ that is supernear to f and either simply additive on each simple partition of each R_S or additive on $\mathcal{C}_{SR}$.*

Appendix For Chapter 6.

Recall that a curve is a continuous map $c: [0, 1] \rightarrow \mathbb{R}^n$. This is equivalent to considering c as determined by n continuous coordinate functions $x_i = f_i(t)$, $1 \leq i \leq n$ each defined on $[0, 1] \subset \mathbb{R}$. Of course, the geometric curve C determined by these functions is usually considered as the set $\{(x_1, \dots, x_n) | t \in [0, 1]\}$. The $*$ -transform of these defining functions leads to the functions $x_i = {}^*f_i(t)$, $1 \leq i \leq n$, each defined on ${}^*[0, 1] \subset {}^*\mathbb{R}$ and they generate the “hypercurve” ${}^*C \subset {}^*\mathbb{R}^n$.

Example 4.4.1.A. Let $\Gamma \in \mathbb{N}_\infty$. Then $F = \{t_i | t_i = i/\Gamma \wedge 0 \leq i \leq \Gamma\}$ is an internal and hyperfinite subset of ${}^*[0, 1]$. By $*$ -transfer F behaves like an ordered partition of the interval $[0, 1]$ as defined in the standard sense. Such a set is termed a **fine partition** (i.e. hyperfinitely many members of $[0, 1]$ generating subintervals that are infinitesimal in length). The internal set F generates the internal set of “points” $P = \{({}^*f_1(t_i), \dots, {}^*f_n(t_i)) | t_i \in F\}$ that are members of the hypercurve ${}^*C$. Now for each $i = 0, \dots, \Gamma - 1$, and each j , $0 \leq j \leq n$ let ${}^*f_j(t_{i+1}) - {}^*f_j(t_i) = d(j, i)$. (If c is continuous, then each $d(j, i) \in \mu(0)$.) For each $i \in {}^*\mathbb{N}$ such that $0 \leq i \leq \Gamma - 1$, the internal set $\ell_i = \{(x_1, \dots, x_n) | \forall j \in {}^*\mathbb{N}, 0 \leq j \leq n, x_j = {}^*f_j(t_i) + t(d(j, i)) \wedge t \in {}^*[0, 1]\}$ is a hyperline segment connecting the two points $({}^*f_1(t_i), \dots, {}^*f_n(t_i))$, $({}^*f_1(t_{i+1}), \dots, {}^*f_n(t_{i+1}))$ on the curve ${}^*C$. From this one obtains the internal hyperpolygonal curve $\mathcal{P}_\Gamma = \bigcup \{\ell_i | 0 \leq i \leq \Gamma - 1\}$. As to the length of $\mathcal{P}_\Gamma$ simply extend the concept of length in the classical sense by defining for each $i = 0, \dots, \Gamma - 1$ the vector $\vec{v}_i = (d(1, i), \dots, d(n, i)) \in {}^*\mathbb{R}^n$. Then let the hyperfinite sum $\sum_{i=0}^{\Gamma-1} \|\vec{v}_i\| = |\mathcal{P}_\Gamma| \in {}^*\mathbb{R}$. Even though, in general, you would have a different hyperpolygon with a different hyperreal length for $\forall \Gamma \in \mathbb{N}_\infty$ we show, using the above terminology and notation that:

If c is continuously differentiable, then for all $\Gamma \in \mathbb{N}_\infty$, $|\mathcal{P}_\Gamma| \in \mu(r)$ and the real number

$$r = \int_a^b \sqrt{\sum_{j=1}^n f_j'(t)^2} dt.$$

Proof. Consider an arbitrary $\Gamma \in \mathbb{N}_\infty$ and the internal partition F of $[0, 1]$. This is a simple fine partition with $t_{i+1} - t_i = dt = 1/\Gamma$. (Actually the following proof holds for any fine partition.) By $*$ -transfer of the standard mean value theorem for the derivative, it follows that for each $i = 0, \dots, \Gamma - 1$, and each $j = 1, \dots, n$ there exists some $t'_{ji} \in (t_i, t_{i+1})$ such that

$$d(j, i) = {}^*f_j'(t'_{ji}) dt.$$

Thus $\|\vec{v}_i\| = \sqrt{\sum_{j=1}^n d(j, i)^2} = \sqrt{\sum_{j=1}^n ({}^*f_j'(t'_{ji}) dt)^2} = \sqrt{\sum_{j=1}^n ({}^*f_j'(t'_{ji}))^2} dt$. Since each f_j' is uniformly continuous then ${}^*f_j'(t_i) = {}^*f_j'(t'_{ji}) + \delta_{ji}$, $\delta_{ji} \in \mu(0)$. Thus $\|\vec{v}_i\| = \sqrt{\sum_{j=1}^n ({}^*f_j'(t_i) + \delta_{ji})^2} dt = \sqrt{\sum_{j=1}^n ({}^*f_j'(t_i))^2 + \epsilon_i} dt$, $\epsilon_i \in \mu(0)$. However, $\sum_{j=1}^n ({}^*f_j'(t_i))^2 + \epsilon_i \approx \sum_{j=1}^n ({}^*f_j'(t_i))^2 \Rightarrow \sqrt{\sum_{j=1}^n ({}^*f_j'(t_i))^2 + \epsilon_i} \approx \sqrt{\sum_{j=1}^n ({}^*f_j'(t_i))^2}$. Hence, $\|\vec{v}_i\| = \sqrt{\sum_{j=1}^n ({}^*f_j'(t_i))^2} dt + \delta_i dt$, $\delta_i \in \mu(0)$. By $*$ -transfer of the finite case, $\max\{|\delta_i| | 0 \leq i \leq \Gamma - 1\} = \delta \in \mu(0)$. Also note that the function $\sqrt{\sum_{j=1}^n f_j'(t)^2}$ is continuous on $[a, b]$. Putting the above together we have that

$$\sum_{i=0}^{\Gamma-1} \|\vec{v}_i\| = |\mathcal{P}_\Gamma| = \sum_{i=0}^{\Gamma-1} \sqrt{\sum_{j=1}^n ({}^*f_j'(t_i))^2} dt + \sum_{i=0}^{\Gamma-1} \delta_i dt.$$

But,

$$\left| \sum_{i=0}^{\Gamma-1} \delta_i dt \right| \leq \sum_{i=0}^{\Gamma-1} |\delta| dt = \delta(b-a) \in \mu(0).$$

Therefore,

$$\sum_{i=0}^{\Gamma-1} \|\vec{v}_i\| = \sum_{i=0}^{\Gamma-1} \sqrt{\sum_{j=1}^n (*f'_j(t_i))^2} dt + \lambda, \quad \lambda \in \mu(0).$$

Since $\sqrt{\sum_{j=1}^n f'_j(t)^2}$ is bounded and integrable then theorem 5.1.2 yields that

$$\text{st}\left(\sum_{i=0}^{\Gamma-1} \sqrt{\sum_{j=1}^n (*f'_j(t_i))^2} dt\right) = \int_a^b \sqrt{\sum_{j=1}^n f'_j(t)^2} dt,$$

and the proof is complete.

What the above definition shows is that in this case our definition $|\mathcal{P}_\Gamma|$ is independent of the particular $\Gamma \in \mathbb{N}_\infty$ chosen (as mentioned within the proof it is actually independent of any fine partition chosen) and also coincides with the classical one. Indeed, we have the following added proposition that shows that the classical concept of the rectifiable curve and the existence of $r \in \mathbb{R}$ such that $|\mathcal{P}_P| \in \mu(r)$ for every fine partition Q of $[0, 1]$ are equivalent concepts.

Proposition 6.1. *Consider continuous $c: [0, 1] \rightarrow \mathbb{R}^n$. Then c is rectifiable if and only if for every fine partition Q of $[0, 1]$*

$$\text{st}(*|\mathcal{P}_Q|) = L.$$

Proof. (Sketch) Let $P = \{0 = u_0 < u_1 < \dots < u_q = 1\}$ be a partition and let $\mathcal{P}_P$ denote the polygonal curve generated by the functions $f_1, \dots, f_n$ and $|\mathcal{P}_P|$ the standard length for the polygonal curve. Consider any fine partition Q of $[0, 1]$. Let $I = [t_i, t_{i+1}]$ be a subinterval generated by Q . Then for any $t \in I$, $|*f_j(t_i) - *f_j(t)|$, $|*f_j(t) - *f_j(t_{i+1})| \in \mu(0)$ for each j since each f_j is uniformly continuous on $[0, 1]$. Note that by *-transfer $*|\mathcal{P}_Q|$ is but the length of the interval hyperpolygonal curve $\mathcal{P}_Q$ as defined in Example 4.4.1.A. Now there are but finitely many infinitesimal subintervals generated by Q that contain the partition points from P . Considering the standard properties of the Euclidean norm, the internal common refinement $P \cup Q$ and the fact that the finite sum of infinitesimals is infinitesimal this leads to

$$|\mathcal{P}_P| \leq *|\mathcal{P}_{P \cup Q}| \leq *|\mathcal{P}_Q| + \epsilon, \quad \epsilon \in \mu(0).$$

Since c is rectifiable then given any $r \in \mathbb{R}^+$ there exists a partition $P(r)$ of $[0, 1]$ such that $0 \leq L - |\mathcal{P}_{P(r)}| < r$. By *-transfer we also have that $*|\mathcal{P}_Q| \leq L$. Consequently,

$$0 \leq L - *|\mathcal{P}_Q| < r + \epsilon.$$

Thus $\text{st}(*|\mathcal{P}_Q|) = L$.

Conversely, let P be any standard and Q any fine partition of $[0, 1]$ Then internal partition $P \cup Q$ is a fine partition. Thus

$$|\mathcal{P}_P| \leq *|\mathcal{P}_{P \cup Q}| \in \mu(L).$$

Hence,

$$\text{st}(|\mathcal{P}_P|) = |\mathcal{P}_P| \leq \text{st}(*|\mathcal{P}_{P \cup Q}|) = L$$

and L is an upper bound for the standard $|\mathcal{P}_P|$ and, thus, an upper bound for all internal partitions of $*[0, 1]$. Moreover, if $s \in \mathbb{R}^+$, then $0 \leq L - *|\mathcal{P}_{P \cup Q}| < s$. Thus by $*$ -transfer, for every $s \in \mathbb{R}^+$ there exists some partition K of $[0, 1]$ such that $0 \leq L - |\mathcal{P}_K| < s$. Thus c is rectifiable.

In Proposition 6.1, the number L is the length of the curve. We now proceed to establish the other Chapter 6 theorems.

Theorem 6.2.1. *Let A be a compact subset of $\mathbb{R}^n$ and the infinitesimal subrectangle $R \subset *A$. Then there exists some $\vec{p} \in A$ such that $R \subset \mu(\vec{p})$.*

Proof. Since $R \neq \emptyset$ let $\vec{x}, \vec{y} \in R$. Then $\vec{x} \approx \vec{y}$. Since A is compact then there exists some $\vec{p} \in \mathbb{R}^n$ such that $y \in \mu(\vec{p})$. Hence for each $\vec{x} \in R$, $\vec{x} \approx \vec{p} \Rightarrow R \subset \mu(\vec{p})$.

IR3

(1) We wish to measure a quantity M for a compact Jordan-measurable set $J \subset R \subset \mathbb{R}^n$, where M is defined on and, at least, additive over members of the set $\{\mathcal{C}_{SR}, R - J, J\}$. Further, if subrectangle $S \subset R - J$, $M(S) = 0$, and $M(R - J) = 0$. Let $v(J)$ denote the Jordan content.

(2) There is a generating function $f(\vec{x})$ that is related to the functional M in the following manner:

- (i) The functions f is continuous on J .
- (ii) Let P be some simple fine partition, $S \in *S(P)$ and $K = *J \cap S \neq \emptyset$. Then there exist $\vec{x}_m \in K$ and $\vec{x}_M \in K$ such that $*f_m = *f(\vec{x}_m) = *\inf\{ *f(\vec{x}) | \vec{x} \in K \} = \inf\{ *f(\vec{x}) | \vec{x} \in K \}$ and $*f_M = *f(\vec{x}_M) = *\sup\{ *f(\vec{x}) | \vec{x} \in K \} = \sup\{ *f(\vec{x}) | \vec{x} \in K \}$.
- (iii) $(*f_m) *v(S) \leq *M(S) \leq (*f_M) *v(S)$. [Note: this is the case where $*L$ is the identity map.]

Theorem 6.2.2. *If IR3 holds, then*

$$M(J) = \int_J f(\vec{x}) dX.$$

Proof. Continuity of f on J yields, since the set of discontinuities of $\hat{f}$ has Lebesgue measure zero, that $\hat{f}$ is integrable on R . Thus $\int_J f(\vec{x}) dX = \int_R \hat{f}(\vec{x}) dX$. Let P be a simple fine partition and any $S \in S(P)$ such that $K = *J \cap S \neq \emptyset$. Then we have from (iii) that

$$*f_m *v(S) = *f_m dX \leq *M(S) \leq *f_M *v(S) = *f_M dX.$$

Assume that $K \neq \emptyset$. Since $f = \hat{f}$ is uniformly continuous on J and $\vec{x}_m, \vec{x}_M \in K \Rightarrow \vec{x}_m \approx \vec{x}_M$ then $*\hat{f}(\vec{x}_m) = *f(\vec{x}_m) \approx *f(\vec{x}_M) = *\hat{f}(\vec{x}_M) \Rightarrow$

$$*M(S)/dX \approx *\hat{f}(\vec{x}_m). \quad (*)$$

For the case that ${}^*J \cap S = \emptyset$, it follows that $S \subset R - J$; which implies that ${}^*M(S) = 0$. Hence in this case ${}^*M(S) = {}^*\hat{f}(\vec{x})$, for any $\vec{x} \in S \Rightarrow$ expression (*). By application of the Infinite Sum Theorem

$$M(R) = \int_R \hat{f}(\vec{x}) dX = \int_J f(\vec{x}) dX = M(R - J) + M(J) = M(J).$$

Theorem 6.2.3. *Let compact Jordan-measurable $J \subset R \subset \mathbb{R}^n$. If continuous $f: J \rightarrow \mathbb{R}$, then for any partition P of R and any $S \in \mathcal{S}(P)$, where $K = J \cap S \neq \emptyset$ there exist $\vec{x}_m \in K$ and $\vec{x}_M \in K$ such that $f_m = f(\vec{x}_m) = \inf\{f(\vec{x}) | \vec{x} \in K\}$ and $f_M = f(\vec{x}_M) = \sup\{f(\vec{x}) | \vec{x} \in K\}$.*

Proof. This follows immediately since each such K is compact and f is continuous on it.

Theorem 6.2.4. (Self-evident Max. and Min.) *Let the rectangle $R \subset \mathbb{R}^n$ and suppose that compact Jordan-measurable $J \subset R$. Let M be defined as in (1) of IR3, continuous $f: J \rightarrow \mathbb{R}$, $\mathcal{P}$ an acceptable set of partitions and any $P \in \mathcal{P}$. If for any $S \in \mathcal{S}(P)$ such that $J \cap S \neq \emptyset$ it follows that $(f_m)v(S) \leq M(S) \leq (f_M)v(S)$, then the infinitesimalizing process IR3 holds.*

Proof. Everything stated in the hypothesis can be written in our first-order set-theoretic language. Noting such things as the *-minimum [resp. *-max.] value of ${}^*f$ on a *-compact ${}^*J \cap S$, where S is an infinitesimal subrectangle, is the same as the minimum [resp. max.] value of ${}^*f$ on ${}^*J \cap S$ by *-transfer and the fact that some simple fine partition exists the result follows from Theorem 6.2.2.

IR4–Method of Constants

(1) In what follows, let for any $A \subset \mathbb{R}^n$ “int” denote the interior of A . Let $\mathcal{A} = \{C_{SR}, \{\text{int}(J \cap S) \neq \emptyset \mid S \in \mathcal{C}_{SR}\}\}$. We wish to measure a quantity M for a Jordan-measurable set $J \subset R \subset \mathbb{R}^n$, where M is, at least, defined on and additive over the members of the set $\{A, R - J, J\}$ and for a subrectangle S , $S \subset R - \text{int}(J)$ it follows that $M(S) = 0$ and $M(R - J) = 0$. Let $v(J)$ denote the Jordan content.

(2) There is a generating function $f(\vec{x})$ that is related to the functional M in the following manner:

- (i) The function f is bounded on J .
- (ii) Let P be any arbitrary simple fine partition, $S \in {}^*\mathcal{S}(P)$ an arbitrary infinitesimal subrectangle and $\emptyset \neq K = {}^*\text{int}({}^*J \cap S)$.
- (iii) There exists some $\vec{x} \in K$ such that ${}^*M(S) = {}^*f(\vec{x}) {}^*v(S)$ or ${}^*M(S)/{}^*v(S) \approx {}^*f(\vec{x})$.

Theorem 6.4.1. *If IR4 holds, then*

$$M(J) = \int_J f(\vec{x}) dX.$$

Proof. First, f is bounded if and only if $\hat{f}$, as it is defined on R , is bounded. We confine our attention to the function $\hat{f}: R \rightarrow \mathbb{R}^n$. From (iii) of IR4, letting $\emptyset \neq K = {}^*\text{int}({}^*J \cap S)$ for $S \in {}^*\mathcal{S}(P)$ there is some $\vec{x} \in K$, hence in S , such that

$${}^*M(S) = {}^*f(\vec{x}) {}^*v(S) = {}^*\hat{f}(\vec{x}) dX \text{ or}$$

$${}^*M(S)/{}^*v(S) \approx {}^*f(\vec{x}) = {}^*\hat{f}(\vec{x}).$$

For the case that $K = \emptyset$ then $S \subset {}^*R - {}^*\text{int}({}^*J) \Rightarrow {}^*M(S) = 0 = {}^*\hat{f}(\vec{x}) dX$ for any $\vec{x} \in S$. Thus in all cases, ${}^*M(S)/dX \approx {}^*\hat{f}(\vec{x})$, $\vec{x} \in S$. Thus from the Infinite Sum Theorem

$$M(R) = \int_R \hat{f}(\vec{x}) dX = \int_J f(\vec{x}) dX = M(R - J) + M(J) = M(J)$$

and the proof is complete.

Prior to the next proof I point out one aspect of the Jordan-measurable subsets that will tacitly appear throughout many of these proofs. Let J be a Jordan-measurable. Then it is part of the definition that J is a bounded subset of $\mathbb{R}^n$. As is well-known this implies that the boundary of J , ∂J is Jordan-measurable and that $v(\partial J) = 0$. Further, $\partial(\text{int}(J)) \subset \partial J \Rightarrow$ that $\text{int}(J)$ is Jordan-measurable. Since $\text{int}(\partial J) \cap \text{int}(\text{int}(J)) = \emptyset$ and $J \subset \partial J \cup \text{int}(J) \Rightarrow v(J) \leq v(\text{int}(J)) \Rightarrow v(\text{int}(J)) = v(J)$.

Theorem 6.4.2. (Self-evident Method of Constants) *Let the rectangle $R \subset \mathbb{R}^n$ and suppose that Jordan-measurable $J \subset R$. Let M be defined as in (1) of IR4, continuous $f: R \rightarrow \mathbb{R}$, $\mathcal{P}$ an acceptable set of partitions of R and any $P \in \mathcal{P}$. If for any $S \in \mathcal{S}(P)$ such that $\emptyset \neq K = \text{int}(J \cap S)$ there exists some $\vec{x} \in K$ and some $\vec{y} \in S$ such that (i) $M(K) = f(\vec{x}) v(K)$ and (ii) $M(S) = f(\vec{y}) v(S)$, then the infinitesimalizing process IR4 holds for f restricted to J .*

Proof. Let $k = f|_J$ and $Q \in {}^*\mathcal{P}$ be a simple fine partition. Assume that $\emptyset \neq K = {}^*\text{int}({}^*J \cap S)$ for $S \in Q$. Note that k is bounded. By $*$ -transfer of the hypotheses there exists some $\vec{x} \in K$ (thus $\vec{x} \in S$ also) and some $\vec{y} \in S$ such that (i) ${}^*M(K) = {}^*k(\vec{x}) {}^*v(K) = {}^*f(\vec{x}) {}^*v(K)$ and (ii) ${}^*M(S) = {}^*f(\vec{y}) {}^*v(S)$. Thus since ${}^*v(K) \neq 0$ then

$${}^*M(K)/{}^*v(K) = {}^*f(\vec{x})$$

$${}^*M(S)/{}^*v(S) = {}^*f(\vec{y}).$$

But, both $\vec{x}, \vec{y} \in S \Rightarrow \vec{x} \approx \vec{y}$. Uniform continuity of f yields that ${}^*f(\vec{x}) \approx {}^*f(\vec{y}) \Rightarrow {}^*M(S)/{}^*v(S) \approx {}^*f(\vec{x}) = {}^*k(\vec{x}) = {}^*\hat{k}(\vec{x})$, where $\vec{x} \in S$ and thus

$$M(R) = \int_R \hat{k}(x) dX = \int_J f(\vec{x}) dX.$$

Theorem 6.4.3. (Extended Self-evident Method of Constants) *Let the rectangle $R \subset \mathbb{R}^n$ and suppose that Jordan-measurable $J \subset R$. Let M be defined as in (1) of IR4, continuous $f: R \rightarrow \mathbb{R}$, continuous $g: R \rightarrow \mathbb{R}$, $\mathcal{P}$ an acceptable set of partitions of R and any $P \in \mathcal{P}$. If for any $S \in \mathcal{S}(P)$ and $\emptyset \neq K = \text{int}(J \cap S)$ there exists some $\vec{x}_1, \vec{x}_2 \in K$ and some $\vec{y}_1, \vec{y}_2 \in S$ such that (i) $M(K) = f(\vec{x}_1) g(\vec{x}_2) v(K)$ and (ii) $M(S) = f(\vec{y}_1) g(\vec{y}_2) v(S)$, then the infinitesimalizing process IR4 holds for $f g$ restricted to J .*

Proof. Let $k = f|_J$ and $h = g|_J$. By $*$ -transfer there exists a simple fine partition $Q \in {}^*\mathcal{P}$ with all the indicated properties. Let $S \in {}^*\mathcal{S}(Q)$, $\emptyset \neq K = {}^*\text{int}({}^*J \cap S)$. Then there exist $\vec{x}_1, \vec{x}_2 \in K$, ($\vec{x}_1, \vec{x}_2 \in S$) and $\vec{y}_1, \vec{y}_2 \in S$ such that

$${}^*M(K)/{}^*v(K) = {}^*k(\vec{x}_1) {}^*h(\vec{x}_2) = {}^*f(\vec{x}_1) {}^*g(\vec{x}_2)$$

$${}^*M(S)/{}^*v(S) = {}^*f(\vec{y}_1) {}^*g(\vec{y}_2).$$

Uniform continuity of f and g imply that ${}^*f(\vec{x}_1) \approx {}^*f(\vec{y}_1)$ and ${}^*g(\vec{y}_2) \approx {}^*g(\vec{x}_1)$. Using the fact that f and g are also bounded we have that

$${}^*f(\vec{y}_1) {}^*g(\vec{y}_2) \approx {}^*f(\vec{x}_1) {}^*g(\vec{x}_1).$$

Hence,

$${}^*M(S)/{}^*v(S) \approx {}^*\hat{k}(\vec{x}_1) {}^*\hat{h}(\vec{x}_1), \quad \vec{x}_1 \in S$$

and the result follows.

[**Remark.** It is obvious that in the above two theorems (i) is not really needed if we know that the y 's are members of K . The reason that (i) is included in the hypothesis is that it is easier to argue for the location of the x 's when applications are considered.]

For a given rectangle R , recall that $F(\mathcal{C}_{SR})$ is the set of all finite sets of subrectangles of R . Extend the definition of the maps $L(f, \bullet)$, $U(f, \bullet)$ for any nonempty collection $\mathcal{F} \in F(\mathcal{C}_{SR})$, where $\emptyset \notin \mathcal{F}$. Let P be any standard partition of R . For a given $J \subset R \subset \mathbb{R}^n$, where to avoid trivialities we always assume that $\text{int}(J) \neq \emptyset$, define the following set theoretic operators each defined on the set of all partitions. The boundary subrectangles $\partial(P) = \{S \mid S \in \mathcal{S}(P) \wedge S \cap J \neq \emptyset \wedge S \cap (R - J) \neq \emptyset\}$; the exterior subrectangles $\text{ext}(P) = \{S \mid S \in \mathcal{S}(P) \wedge S \subset (R - J)\}$, and the inner subrectangles $\text{inn}(P) = \mathcal{S}(P) - (\partial(P) \cup \text{ext}(P))$. Notice that $S \in \text{inn}(P)$ if and only if $S \subset J$. Let $\mathcal{P}$ be a set of partitions of R and $I_{\mathcal{P}} = \{\cup A \mid \exists P(P \in \mathcal{P} \wedge A = \text{inn}(P))\} \cup \{S \mid \exists P(P \in \mathcal{P} \wedge S \in \text{inn}(P))\}$. When considering the nonstandard extension of these operators, we use the notation ∂ , inn and ext rather than the notation ${}^*\partial$, ${}^*\text{ext}$ and ${}^*\text{inn}$. Since any fine partition Q is hyperfinite and the sets $\partial(Q)$, $\text{inn}(Q)$ and $\text{ext}(Q)$ are internal subsets of Q then they are hyperfinite and by * -transfer of the standard finite case it follows that for bounded $f: J \rightarrow \mathbb{R}$

$${}^*L(\hat{f}, Q) = {}^*L(\hat{f}, \partial(Q)) + {}^*L(\hat{f}, \text{ext}(Q)) + {}^*L(\hat{f}, \text{inn}(Q)).$$

$${}^*U(\hat{f}, Q) = {}^*U(\hat{f}, \partial(Q)) + {}^*U(\hat{f}, \text{ext}(Q)) + {}^*U(\hat{f}, \text{inn}(Q)).$$

Applying a method similar to that used to establish Proposition 6.1, it follows from the definition of Jordan-content that J has Jordan-content (i.e. is Jordan-measurable) if and only if for every fine partition Q of ${}^*R$,

$$\sum_{S \in \partial(Q)} {}^*v(S) \approx 0.$$

Proposition 6.2. *Let Jordan-measurable $J \subset R \subset \mathbb{R}^n$, bounded $f: J \rightarrow \mathbb{R}$, and Q be any fine partition of ${}^*R$. Then*

$${}^*L(\hat{f}, \partial(Q)) \approx 0, \quad {}^*U(\hat{f}, \partial(Q)) \approx 0,$$

$${}^*L(f, \text{inn}(Q)) \approx {}^*L(\hat{f}, Q), \quad {}^*U(f, \text{inn}(Q)) \approx {}^*U(\hat{f}, Q).$$

Proof. There exist some $m, M \in \mathbb{R}$ such that for each $\vec{x} \in R$, $m \leq \hat{f}(\vec{x}) \leq M$. Thus

$$m \sum_{S \in \partial(Q)} {}^*v(S) \leq {}^*L(\hat{f}, \partial(Q)) \leq {}^*U(\hat{f}, \partial(Q)) \leq M \sum_{S \in \partial(Q)} {}^*v(S).$$

From the above observation relative to the Jordan-content, it follows that ${}^*L(\hat{f}, \partial(Q)) \approx {}^*U(\hat{f}, \partial(Q)) \approx 0$ and the second part of the conclusion follows from the fact that ${}^*\hat{f}(\vec{x}) = 0$ for each $\vec{x} \in S$ and for each $S \in \text{ext}(Q)$.

Proposition 6.3. *Suppose that Jordan-measurable $J \subset R \subset \mathbb{R}^n$ and bounded $f: J \rightarrow \mathbb{R}$. Let $Q = Q_1 \times \cdots \times Q_n$ be an internal partition of ${}^*R$, where each Q_i is an internal partition of $[a_i, b_i]$, for $1 \leq i \leq n$. If ${}^*\text{mesh}(Q_i) \in \mu(0)$ (i.e. Q is a fine partition), then*

$$L(\hat{f}) \approx {}^*L(f, \text{inn}(Q)), \quad U(\hat{f}) \approx {}^*U(f, \text{inn}(Q)).$$

Proof. From Proposition 5.1 of Appendix 5, $L(\hat{f}) \approx {}^*L(\hat{f}, Q)$ and $U(\hat{f}) \approx {}^*U(\hat{f}, Q)$ since $\hat{f}: R \rightarrow \mathbb{R}$ is bounded. The result follows by application of Proposition 6.2.

Proposition 6.4. *Suppose that $J \subset R \subset \mathbb{R}^n$ and bounded $f: J \rightarrow \mathbb{R}$. Let $\mathcal{P}$ be any infinite set of partitions, $B: I_{\mathcal{P}} \rightarrow \mathbb{R}$, and for each $P \in \mathcal{P}$, B is simply additive on each $\text{inn}(P)$. Suppose that there exists a fine partition $Q \in {}^*\mathcal{P}$ such that for every $S \in \text{inn}(Q)$ there exists some $\vec{p} \in S$ such that*

$${}^*B(S)/({}^*v(S)) \approx {}^*f(\vec{p}),$$

then for any $r \in \mathbb{R}^+$ and $A = \text{inn}(Q)$

$$-r {}^*v(\cup A) + {}^*L(f, A) < {}^*B(\cup A) < {}^*U(f, A) + r {}^*v(\cup A).$$

Proof. First, note that by * -transfer that ${}^*v$ is defined for $\cup A$, ${}^*v(\cup A) = \sum_{S \in A} {}^*v(S)$ and that ${}^*B$ is defined on $\cup A$. Now repeat the proof of Proposition 5.4 with $\text{inn}(P)$ substituted for P , $\cup \text{inn}(P)$ or $\cup \text{inn}(Q)$ substituted for R and $\cup \text{inn}(Q)$ for Q .

Proposition 6.5. (An Infinite Sum Theorem) *Suppose that Jordan-measurable $J \subset R \subset \mathbb{R}^n$ and bounded $f: J \rightarrow \mathbb{R}$. Let $\mathcal{P}$ be any infinite set of partitions, $B: I_{\mathcal{P}} \rightarrow \mathbb{R}$, and for each $P \in \mathcal{P}$, B is simply additive on $\text{inn}(P)$. Suppose that there exists a fine partition $Q \in {}^*\mathcal{P}$ such that for every $S \in \text{inn}(Q) = A$ there exists some $\vec{p} \in S$ such that*

$${}^*B(S)/({}^*v(S)) \approx {}^*f(\vec{p}),$$

then f is integrable and

$$\text{st}({}^*B(\cup A)) = \int_J f(\vec{x}) dX.$$

Proof. From Proposition 6.4, we have that for any $r \in \mathbb{R}^+$, $-r {}^*v(\cup A) + {}^*L(f, A) < {}^*B(\cup A) < {}^*U(f, A) + r {}^*v(\cup A)$. By * -transfer of the standard case, then ${}^*v(\cup A) \leq {}^*v({}^*J) = v(J) \leq {}^*v((\cup A) \cup (\partial(Q))) = {}^*v(\cup A) + {}^*v(\cup(\partial(Q)))$ implies that $\text{st}({}^*v(\cup A)) \leq v(J) \leq \text{st}({}^*v(\cup A))$ or that $\text{st}({}^*v(\cup A)) = v(J)$. Consequently,

$$\begin{aligned} -r \text{st}({}^*v(\cup A)) + \text{st}({}^*L(f, A)) &\leq \text{st}({}^*B(\cup A)) \leq \\ &\text{st}({}^*U(f, A)) + r \text{st}({}^*v(\cup A)) \end{aligned}$$

yields by Proposition 6.3 that

$$-r v(J) + L(\hat{f}) \leq \text{st}({}^*B(\cup A)) \leq U(\hat{f}) + r v(J).$$

Since r is an arbitrary member of $\mathbb{R}^+$ then

$$L(\hat{f}) = U(\hat{f}) = \int_R \hat{f}(\vec{x}) dX = \int_J f(\vec{x}) dX = \mathbf{st}(*B(\cup A)).$$

Obviously, the above Infinite Sum Theorem almost yields the result being sought. But, just as obviously the functional B would need some additional property akin to a simple property displayed by the basic Jordan-measure v . Assume that a functional such as B is defined on J and say that B is **Jordan-like** if for the fine partition Q of Proposition 6.5 $*B(\cup A) \approx B(J)$. It is not difficult to show if for each $r \in \mathbb{R}^+$ there exists a $\delta \in \mathbb{R}^+$ such that for each $P \in \mathcal{P}$ with $\|P\| < \delta$ it follows that $|B(\text{inn}(P)) - B(J)| < r$, then B is Jordan-like for each fine partition $Q \in \mathcal{P}$. (Of course, it is assumed that for each δ mentioned in this last statement that such a partition P exists such that $\|P\| < \delta$.) If B satisfies this property for a set of partitions $\mathcal{P}$ then we say that B **has an ordinary Jordan-like property**.

IR5

(1) We wish to measure a quantity M for a compact Jordan-measurable set $J \subset R \subset \mathbb{R}^n$, where for simplicity assume that M is defined on and, at least, additive over members of the set $\{I_{\mathcal{P}}, J\}$, and $\mathcal{P}$ is the set of all simple partitions of R . As usual, $v(J)$ denotes the Jordan-content.

(2) There is a generating function $f(\vec{x})$ that is related to the functional M in the following manner:

- (i) The function f is continuous on J .
- (ii) For a some simple fine partition, Q , and each $S \in \text{inn}(Q)$ there exist $\vec{x}_m \in S$ and $\vec{x}_M \in S$ such that $*f_m = *f(\vec{x}_m) = \inf\{ *f(\vec{x}) | \vec{x} \in S \} = \inf\{ *f(\vec{x}) | \vec{x} \in S \}$ and $*f_M = *f(\vec{x}_M) = \sup\{ *f(\vec{x}) | \vec{x} \in S \} = \sup\{ *f(\vec{x}) | \vec{x} \in S \}$.
- (iii) $(*f_m) *v(S) \leq *M(S) \leq (*f_M) *v(S)$.
- (iv) For Q , M is Jordan-like.

Proposition 6.6. *Suppose there exists a fine partition such that IR5 holds for Q . Then*

$$M(J) = \int_J f(\vec{x}) dX.$$

Proof. Consider the fine partition Q from (iii) of IR5. Then for each $S \in \text{inn}(Q)$

$$(*f_m) *v(S) \leq *M(S) \leq (*f_M) *v(S).$$

Now f is uniformly continuous on J and $\vec{x}_m, \vec{x}_M \in S$ imply that $\vec{x}_m \approx \vec{x}_M$ and that $*f_m \approx *f_M$. Therefore,

$$*M(S) / *v(S) \approx *f(\vec{x}_m).$$

Consequently, from (iv)

$$\int_R \hat{f}(\vec{x}) dX = \int_J f(\vec{x}) dX = \mathbf{st}(*B(\cup A)) = B(J).$$

We also have a self-evident theorem that goes with Proposition 6.6.

Proposition 6.7. (Self-evident Max. and Min.) *Let the rectangle $R \subset \mathbb{R}^n$ and suppose that compact Jordan-measurable $J \subset R$. Let M be defined as in (1) of IR5, continuous $f: J \rightarrow \mathbb{R}$, $\mathcal{P}$ is the set of all simple partition on R and any $P \in \mathcal{P}$. If for any $S \in \text{inn}(P)$ it follows that $(f_m)v(S) \leq M(S) \leq (f_M)v(S)$, and M has the ordinary Jordan-like property, then the infinitesimalizing process IR3 holds.*

Proof. For each fine partition $Q \in \mathcal{P}$ the functional M is Jordan-like. Since there exists a fine partition $Q \in {}^*\mathcal{P}$ then the *-transfer of Theorem 6.2.3 and the hypotheses yields that (ii), (iii), and (iv) hold for Q . The conclusion follows from Proposition 6.6.

The Method of Constants can be greatly simplified for functionals M that are Jordan-like.

IR6–Method of Constants

(1) We wish to measure a quantity M for a Jordan-measurable set $J \subset R \subset \mathbb{R}^n$, where for simplicity assume that M is defined on and, at least, additive over members of the set $\{I_P, J\}$, and $\mathcal{P}$ is the set of all simple partitions of R . As usual, $v(J)$ denotes the Jordan-content.

(2) There is a generating function $f(\vec{x})$ that is related to the functional M in the following manner:

- (i) The function f is bounded on J .
- (ii) For some simple fine partition, Q , and each $S \in \text{inn}(Q)$ there exists some $\vec{x} \in S$ such that ${}^*M(S) = {}^*f(\vec{x}) {}^*v(S)$ or ${}^*M(S)/{}^*v(S) \approx {}^*f(\vec{x})$.
- (iii) The functional M is Jordan-like for Q .

Proposition 6.8. *Suppose there exists a fine partition such that IR6 holds for Q . Then*

$$M(J) = \int_J f(\vec{x}) dX.$$

Proof. This is immediate from Proposition 6.5 and that M is Jordan-like for Q .

Proposition 6.9. (Self-evident Method of Constants) *Let the rectangle $R \subset \mathbb{R}^n$ and suppose that Jordan-measurable $J \subset R$. Let M be defined as in (1) of IR6, bounded $f: J \rightarrow \mathbb{R}$, where $\mathcal{P}$ is the set of all simple partitions on R and any $P \in \mathcal{P}$. If for any $S \in \text{inn}(P)$ there exists some $\vec{x} \in S$ such that $M(S) = f(\vec{x})v(S)$, and M has the ordinary Jordan-like property, then the infinitesimalizing process IR3 holds.*

Proof. Again this is immediate from *-transfer and Proposition 6.5.

The advantages of having a Jordan-like functional are obvious when one compares our last proposition in the section with Theorem 6.4.3.

Proposition 6.10. (Extended Self-evident Method of Constants). *Let the rectangle $R \subset \mathbb{R}^n$ and suppose that Jordan-measurable $J \subset R$. Let M be defined as in (1) of IR6, continuous $f, g: J \rightarrow \mathbb{R}$, where $\mathcal{P}$ is the set of all simple partitions on R and any $P \in \mathcal{P}$. If for any $S \in \text{inn}(P)$ there exists some $\vec{x}_1 \in S$ and some $\vec{x}_2 \in S$ such that $M(S) = f(\vec{x}_1)g(\vec{x}_2)v(S)$, and M has the ordinary Jordan-like property, then the infinitesimalizing process IR6 holds.*

Proof. The proof follows from Proposition 6.5 and the observation that for a fine partition Q and $S \in \text{inn}(Q)$ if $\vec{x}_1, \vec{x}_2 \in S$, then $\vec{x}_1 \approx \vec{x}_2$ and ${}^*f(\vec{x}_1) {}^*g(\vec{x}_2) \approx {}^*f(\vec{x}_1) {}^*g(\vec{x}_1)$.

Appendix For Chapter 7.

Theorem 7.2.2. *Let Jordan-measurable $J \subset R \subset \mathbb{R}^n$, bounded $f: J \rightarrow \mathbb{R}$, and Q be any fine partition of ${}^*R$. Let hyperfinite ${}^*\mathcal{S}(Q) = \{S_i \mid 0 \leq i \leq \Gamma - 1\}$. Assume that there exists a hyperfinite sequence U_i , where $U_i \in S_i$ for each $S_i \in \mathcal{S}(Q)$. Then*

$$\text{st}\left(\sum_{S(i) \in \text{inn}(Q)} {}^*f(U_i) {}^*v(S_i)\right) = \text{st}\left(\sum_{S(i) \in \mathcal{S}(Q)} {}^*\hat{f}(U_i) {}^*v(S_i)\right).$$

Proof. First, recall the following definitions. The boundary subrectangles $\partial(Q) = \{S \mid S \in {}^*\mathcal{S}(Q) \wedge S \cap {}^*J \neq \emptyset \wedge S \cap ({}^*\mathbb{R}^n - {}^*J) \neq \emptyset\}$; the exterior subrectangles $\text{ext}(Q) = \{S \mid S \in {}^*\mathcal{S}(Q) \wedge S \subset ({}^*\mathbb{R}^n - {}^*J)\}$, and the inner subrectangles $\text{inn}(Q) = \mathcal{S}(Q) - (\partial(Q) \cup \text{ext}(Q))$. Notice that we have used the notation ∂ , inn and ext rather than the notation ${}^*\partial$, ${}^*\text{ext}$ and ${}^*\text{inn}$ even though these sets are generated by obvious extended standard set-theoretic operators. It follows directly from the definition of Jordan-content that J has Jordan-content if and only if for every fine partition Q

$$\sum_{S(i) \in \partial(Q)} {}^*v(S_i) \approx 0.$$

Since there exists $m, M \in \mathbb{R}$ such that for all $\vec{x} \in R$, $m \leq \hat{f}(\vec{x}) \leq M$ then noting that $\partial(Q)$ is internal and, hence, a hyperfinite set we have from * -transfer of the finite case that

$$m \sum_{S(i) \in \partial(Q)} {}^*v(S_i) \leq \sum_{S(i) \in \partial(Q)} {}^*\hat{f}(U_i) {}^*v(S_i) \leq M \sum_{S(i) \in \partial(Q)} {}^*v(S_i).$$

Thus $\sum_{S(i) \in \partial(Q)} {}^*\hat{f}(U_i) {}^*v(S_i) \approx 0$.

The sets $\text{inn}(Q)$ and $\text{ext}(Q)$ are also hyperfinite sets and from the properties of finite addition and the fact that ${}^*\hat{f}(U_i) = 0$ for each $U_i \in S_i \in \text{ext}(Q)$ it follows that

$$\begin{aligned} \sum_{S(i) \in \mathcal{S}(Q)} {}^*\hat{f}(U_i) {}^*v(S_i) &= \sum_{S(i) \in \text{inn}(Q)} {}^*\hat{f}(U_i) {}^*v(S_i) + \sum_{S(i) \in \partial(Q)} {}^*\hat{f}(U_i) {}^*v(S_i) + \\ &\quad \sum_{S(i) \in \text{ext}(Q)} {}^*\hat{f}(U_i) {}^*v(S_i) \approx \sum_{S(i) \in \text{inn}(Q)} {}^*\hat{f}(U_i) {}^*v(S_i) + \sum_{S(i) \in \text{inn}(Q)} {}^*f(U_i) {}^*v(S_i). \end{aligned}$$

Since

$$m \sum_{S(i) \in \mathcal{S}(Q)} {}^*v(S_i) \leq \sum_{S(i) \in \mathcal{S}(Q)} {}^*\hat{f}(U_i) {}^*v(S_i) \leq M \sum_{S(i) \in \mathcal{S}(Q)} {}^*v(S_i)$$

and

$$\sum_{S(i) \in \mathcal{S}(Q)} {}^*v(S_i) = v(R)$$

then

$$\sum_{S(i) \in \mathcal{S}(Q)} {}^*\hat{f}(U_i) {}^*v(S_i)$$

is a limited number and the result follows.

Appendix For Chapter 8.

Theorem 8.3.1. *For each $\epsilon \in \mu(0)$ the set $o(\epsilon) = \{\epsilon h \mid h \in \mu(0)\}$ is an ideal in $\mu(0)$.*

Proof. First, we show that $o(\epsilon)$ is a subring of the ring $\mu(0)$. Since $o(\epsilon) \subset \mu(0)$, then consider arbitrary $\epsilon h_1 \in o(\epsilon)$ and $\epsilon h_2 \in o(\epsilon)$. Observe that $(\epsilon h_1) - (\epsilon h_2) = \epsilon(h_1 - h_2) = \epsilon h_3$, where $h_3 \in \mu(0)$, and $(\epsilon h_1)(\epsilon h_2) = \epsilon(\epsilon h_1 h_2)$, where $(\epsilon h_1 h_2) \in \mu(0)$. Thus $o(\epsilon)$ is a subring of $\mu(0)$. Let $\delta \in \mu(0)$. Then $\delta(\epsilon h_1) = \epsilon(\delta h_1)$ implies that $o(\epsilon)$ is an ideal in $\mu(0)$.

Theorem 8.3.2. *Let $\epsilon \in \mu(0)^+$. Suppose that $w \in {}^*\mathbb{R}$ and $0 \leq w \leq \epsilon h \in o(\epsilon)$. Then $w \in o(\epsilon)$.*

Proof. Obviously, if $w = 0$ or $w = \epsilon h$, then result holds. Hence, suppose that $0 < w < \epsilon h$. Now $0 < w/\epsilon < h$ implies that $w/\epsilon \in \mu(0)$ implies that $\epsilon(w/\epsilon) = w \in o(\epsilon)$.

Theorem 8.3.3. *Let $\epsilon, \delta \in \mu(0)^+$. If $0 \leq \delta \leq \epsilon$, then $o(\delta) \subset o(\epsilon)$ and $o(\delta)$ is a ideal in $o(\epsilon)$.*

Proof. We know that $o(\delta)$ is an ideal in $\mu(0)$. Thus let $\delta h \in o(\delta)$ and $\epsilon h_1 \in o(\epsilon)$. Then $\delta(\epsilon h h_1) \in o(\delta)$ implies that $o(\delta)$ is an ideal in $o(\epsilon)$. To show that $o(\delta) \subset o(\epsilon)$ we need only consider the nonnegative elements. Let $0 \leq \delta h \in o(\delta)$. Then $0 \leq \delta h < \epsilon h$ implies by Theorem 8.3.2 that $\delta h \in o(\epsilon)$.

Are there order ideals such that $o(\delta) \subset o(\epsilon)$ and $o(\delta) \neq o(\epsilon)$?

(1) Let $0 < \delta \leq \epsilon$, $\delta, \epsilon \in \mu(0)^+$. Then $o(\delta\epsilon) \subset o(\epsilon)$ and $o(\delta\epsilon) \neq o(\epsilon)$.

Proof. $\delta\epsilon \in o(\epsilon)$, but $\delta\epsilon \notin o(\delta\epsilon)$ for $\delta\epsilon h = \delta\epsilon \Rightarrow h = 1$. Indeed, if $r \in \mathcal{O} - \mu(0)$, then $\delta\epsilon r \in o(\epsilon)$, but $\delta\epsilon r \notin o(\delta\epsilon)$.

(2) Let $n \in \mathbb{N}^+ = \mathbb{N} - \{0\}$, $\epsilon \in \mu(0)^+$. Then $o(\epsilon^n) \subset o(\epsilon^{n-1}) \subset \dots \subset o(\epsilon)$ and $o(\epsilon^i) \neq o(\epsilon^j)$; $1 \leq i, j \leq n$; $i \neq j$.

Proof. From (1).

Let $\{\epsilon_1, \dots, \epsilon_k\} \subset \mu(0)$. Define $o(\epsilon_1, \dots, \epsilon_k) = \{\epsilon_1 h_1 + \dots + \epsilon_k h_k \mid h_i \in \mu(0) \wedge 1 \leq i \leq k\}$.

Theorem 8.3.4. *Let $\epsilon = \max\{|\epsilon_1|, \dots, |\epsilon_k|\}$. Then $o(\epsilon_1, \dots, \epsilon_k) = o(\epsilon)$.*

Proof. Assume that $x = \sum_{i=1}^n \epsilon_i h_i \in o(\epsilon_1, \dots, \epsilon_n)$. Let $\delta \in \mu(0)$ and $\delta \geq \max\{|\epsilon_1|, \dots, |\epsilon_n|\}$. Then

$$0 \leq \left| \sum_{i=1}^n \epsilon_i h_i \right| \leq \sum_{i=1}^n |\epsilon_i| |h_i| \leq n\epsilon\delta = \epsilon(n\delta) \in o(\epsilon).$$

Thus $o(\epsilon_1, \dots, \epsilon_n) \subset o(\epsilon)$.

On the other hand, let $\epsilon h \in o(\epsilon)$. Since $\epsilon = \epsilon_j$ for some j such $1 \leq j \leq n$ then $\epsilon h \in o(\epsilon_1, \dots, \epsilon_n) \Rightarrow o(\epsilon) \subset o(\epsilon_1, \dots, \epsilon_n)$.

Theorem 8.3.5. *Let $\epsilon = \max\{|\epsilon_1|, \dots, |\epsilon_k|\}$. Then*

$$o(\epsilon_1, \dots, \epsilon_k) = o(\sqrt{\epsilon_1^2 + \dots + \epsilon_k^2}) = o(\epsilon).$$

Proof. By *-transfer $\sqrt{\sum_{i=1}^n \epsilon_i^2} \leq \sum_{i=1}^n |\epsilon_i|$. Now for $h \in \mu(0)$ it follows that $|h|\sqrt{\sum_{i=1}^n \epsilon_i^2} \leq n|h|\epsilon$; in which case by Theorem 8.3.3 $|h|\sqrt{\sum_{i=1}^n \epsilon_i^2} \in o(\epsilon) \Rightarrow h\sqrt{\sum_{i=1}^n \epsilon_i^2} \in o(\epsilon)$. Hence, $o(\sqrt{\sum_{i=1}^n \epsilon_i^2}) \subset o(\epsilon)$.

Conversely, let $\epsilon h \in o(\epsilon)$. Then $\epsilon|h| \in o(\epsilon)$. But $|\epsilon h| = \epsilon|h| = |\epsilon_j||h| \leq |h|\sqrt{\sum_{i=1}^n \epsilon_i^2}$. Once again Theorem 8.3.3 implies that $\epsilon h \in o(\sqrt{\sum_{i=1}^n \epsilon_i^2})$.

Theorem 8.4.1. *Let $1 \leq n \in \mathbb{N}$. Suppose that $f^{(n-1)}: [a, nb] \rightarrow \mathbb{R}$ and that $f^{(n)}: (a, nb) \rightarrow \mathbb{R}$, where $f^{(k)}$ denotes the k th derivative of f . Then there exists some $t \in (a, nb)$ such that $\Delta^n f(a, b) = f^{(n)}(t) b^n$.*

Proof. By induction. We know the result holds for $n = 1$, therefore, assume that result holds for $n-1$ and let $g(x) = f(x+b) - f(x)$, $x \in [a, a+(n-1)b]$. Now $g^{(n-1)}(x) = f^{(n-1)}(x+b) - f^{(n-1)}(x) \in \mathbb{R}$ for each $x \in [a, a+(n-1)b]$. Thus, by induction, there exists some $t_0 \in (a, a+(n-1)b)$ such that $\delta^{(n-1)} g(a, b) = g^{(n-1)}(t_0) b^{n-1} = (f^{(n-1)}(t_0+b) - f^{(n-1)}(t_0)) b^{n-1}$. Applying the mean value theorem yields that there exists some $t \in (t_0, t_0+b)$ such that $g^{(n-1)}(t_0) = f^{(n-1)}(t_0+b) - f^{(n-1)}(t_0) = f^n(t) b$. Thus, $\Delta^n f(a, b) = \Delta^{(n-1)} g(a, b) = f^n(t) b^n$, where $t \in (a, a+nb)$.

Corollary 8.4.1.1 *Let $1 \leq n \in \mathbb{N}$. Suppose that $f^{(n-1)}: [a, b] \rightarrow \mathbb{R}$ and that $f^{(n)}: (a, b) \rightarrow \mathbb{R}$, then for each $dx \in \mu(0)^+$ and $c \in {}^*[a, b]$, there exists some $t \in (c, c+ndx)$ such that $\Delta^n {}^*f(c, c+dx) = {}^*f^{(n)}(t) (dx)^n$.*

Proof. This follows from *-transfer and the fact that $[c, c+ndx] \subset {}^*[a, b]$.

Theorem 8.4.2. *Let $1 \leq n \in \mathbb{N}$. Suppose that $f^{(n-1)}: [a, b] \rightarrow \mathbb{R}$ and that $f^{(n)}: (a, b) \rightarrow \mathbb{R}$. If $c \in (a, b)$, then for each $dx \in \mu(0)$, $dx \geq 0$ [resp. $dx < 0$]*

$$f^n(c) (dx)^n \approx \Delta^n {}^*f(c, c+dx), [\text{resp. } f(c+dx, c)] \pmod{o((dx)^n)}.$$

Proof. Let $c \in (a, b)$ and consider any $r \in \mathbb{R}^+$ such that $[c, c+nr] \subset [a, b]$. Define $g: [c, c+(n-1)r] \rightarrow \mathbb{R}$ by $g(y) = f(y+r) - f(y)$. Theorem 8.4.1 implies that there exists some $t_1 \in (c, c+(n-1)r)$ such that $\Delta^{(n-1)} g(c, r) = g^{(n-1)}(t_1) r^{n-1}$. By *-transfer it follows that for $dx \in \mu(0)^+$ there exists some $t \in (c, c+(n-1)dx)$ such that $\Delta^{(n-1)} {}^*g(c, dx) = {}^*g^{(n-1)}(t) dx^{n-1}$. Consequently, by *-transfer, one obtains

$$\Delta^n {}^*f(c, dx) = \frac{{}^*f^{(n-1)}(t+dx) - {}^*f^{(n-1)}(t)}{dx} dx^n.$$

Consequently,

$$\frac{\Delta^n {}^*f(c, dx)}{dx^n} = \frac{{}^*f^{(n-1)}(t+dx) - {}^*f^{(n-1)}(t)}{dx} =$$

$$\begin{aligned}
 & \left(\frac{{}^*f^{(n-1)}(t+dx) - f^{(n-1)}(c)}{(t+dx) - c} \right) \left(\frac{(t+dx) - c}{dx} \right) + \\
 & \quad \left(\frac{f^{(n-1)}(c) - {}^*f^{(n-1)}(t)}{c - t} \right) \left(\frac{c - t}{dx} \right) = \\
 & \left(\frac{{}^*f^{(n-1)}(c+dx_1) - f^{(n-1)}(c)}{dx_1} \right) \left(\frac{dx_1}{dx} \right) + \\
 & \quad \left(\frac{{}^*f^{(n-1)}(c+dx_2) - f^{(n-1)}(c)}{dx_2} \right) \left(\frac{-dx_2}{dx} \right),
 \end{aligned}$$

where $dx_1 = t + dx - c$, $dx_2 = t - c \in \mu(0)$. We observe that $1 < dx_1/dx < n$ and that $0 < dx_2/dx < n - 1$. Recall that

$$\begin{aligned}
 \text{st} \left(\left(\frac{{}^*f^{(n-1)}(c+dx_1) - f^{(n-1)}(c)}{dx_1} \right) \left(\frac{dx_1}{dx} \right) \right) &= f^n(c) \text{st} \left(\frac{dx_1}{dx} \right) \\
 \text{st} \left(\left(\frac{{}^*f^{(n-1)}(c+dx_2) - f^{(n-1)}(c)}{dx_2} \right) \left(\frac{-dx_2}{dx} \right) \right) &= f^n(c) \text{st} \left(\frac{-dx_2}{dx} \right)
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 & \left(\frac{{}^*f^{(n-1)}(c+dx_1) - f^{(n-1)}(c)}{dx_1} \right) \left(\frac{dx_1}{dx} \right) + \\
 & \quad \left(\frac{{}^*f^{(n-1)}(c+dx_2) - f^{(n-1)}(c)}{dx_2} \right) \left(\frac{-dx_2}{dx} \right) \approx \\
 & f^n(c) \left(\text{st} \left(\frac{dx_1}{dx} \right) + \text{st} \left(\frac{-dx_2}{dx} \right) \right) = \\
 & f^n(c) \left(\text{st} \left(\frac{dx_1}{dx} - \frac{dx_2}{dx} \right) \right) = f^n(c).
 \end{aligned}$$

From page 71. A simple proof shows that if $r, r_0 \in {}^*[-1, 1]$, then the best we can say is that

$$\| {}^*c(t + r \epsilon) - \vec{g}(t + r_0 \epsilon) \| \in o(\epsilon). \quad (6)$$

Proof. $\| {}^*c(t + r \epsilon) - \vec{g}(t + r_0 \epsilon) \| = \| (h_1, \dots, h_p) r \epsilon \| = \lambda \epsilon, \quad \lambda \in \mu(0)$.

From page 72. Considering any $(s, t) (s_0, t_0) \in {}^*[-1, 1] \times {}^*[-1, 1]$ a simple proof yields that

$$\| {}^*\vec{r}(u_0 + s \delta, v_0 + t \epsilon) - {}^*\vec{k}(u_0 + s_0 \delta, v_0 + t_0 \epsilon) \| \in o(\max\{\delta, \epsilon\}) = o(\lambda). \quad (4)$$

Proof. $\| {}^*\vec{r}(u_0 + s \delta, v_0 + t \epsilon) - {}^*\vec{k}(u_0 + s_0 \delta, v_0 + t_0 \epsilon) \| = \sqrt{s^2 \delta^2 h_s^2 + t^2 \epsilon^2 h_t^2}$. Hence

$$|s \delta h_s| \leq \sqrt{s^2 \delta^2 h_s^2 + t^2 \epsilon^2 h_t^2} \leq |s \delta h_s| + |t \epsilon h_t|,$$

$$|t \epsilon h_t| \leq \sqrt{s^2 \delta^2 h_s^2 + t^2 \epsilon^2 h_t^2} \leq |s \delta h_s| + |t \epsilon h_t|,$$

implies that

$$\left| \frac{s \delta h_s}{2} \right| + \left| \frac{t \epsilon h_t}{2} \right| \leq \sqrt{s^2 \delta^2 h_s^2 + t^2 \epsilon^2 h_t^2} \leq |s \delta h_s| + |t \epsilon h_t|.$$

Consequently,

$$\lambda_s \delta + \lambda_t \epsilon \leq \sqrt{s^2 \delta^2 h_s^2 + t^2 \epsilon^2 h_t^2} \leq \gamma_s \delta + \gamma_t \epsilon,$$

where $\lambda_s, \lambda_t, \gamma_s, \gamma_t \in \mu(0)$. The result follows from Theorem 8.3.4.

From page 75. Prior to discussing the McShane integral, a general theorem that establishes that λ -fine partitions exist for each L -gauge is a useful addition to this appendix.

Proposition 8.1. *Let X be a nonempty connected topological space. Let $\mathcal{C}$ be a collection of subsets of X with the following two properties.*

- (i) *If $x \in X$, then there exists some $C \in \mathcal{C}$ such that $x \in \text{int}(C)$.*
- (ii) *If $C_1, C_2 \in \mathcal{C}$ and $C_1 \cap C_2 \neq \emptyset$, then $C_1 \cup C_2 \in \mathcal{C}$.*

The following two statements are equivalent.

- (A) *The space X is compact.*
- (B) *The set X is an element of every collection of subsets of X that satisfy (i) and (ii).*

Proof. Assume that X is compact and the set $\mathcal{C}$ satisfies (i) and (ii) above. Then there exist a finite subset $\{C_1, \dots, C_n\}$ of $\mathcal{C}$ such that $X = \bigcup \{\text{int}(C_i) \mid 1 \leq i \leq n\} = \bigcup \{C_i \mid 1 \leq i \leq n\}$ from (i) and compactness. Consider C_1 and assume that $n > 1$. Then there exists some C_k such that $C_1 \cap C_k \neq \emptyset$ from the connectedness of X . From (ii) $A_1 = C_1 \cup C_k \in \mathcal{C}$. If $n > 2$, then there exists some C_{k_2} , where $C_{k_2} \notin \{C_1, C_k\}$, such that $A_1 \cap C_{k_2} \neq \emptyset$. Again from (ii) $A_2 = A_1 \cup C_{k_2} \in \mathcal{C}$. Continuing this finite process leads to the conclusion that $X = \bigcup \{C_i \mid 1 \leq i \leq n\} \in \mathcal{C}$.

Conversely, assume that (B) holds and let $\mathcal{G}$ be an open cover of X . Recall that $F(\mathcal{G})$ is the set of all finite subsets of $\mathcal{G}$. Let $\mathcal{C} = \{A \mid A \subset X \wedge \exists Y(Y \in F(\mathcal{G}) \wedge (A \subset \bigcup Y))\}$. Obviously, since $\mathcal{G} \subset \mathcal{C}$, (i) holds. Moreover, if $C_1, C_2 \in \mathcal{C}$ and $C_1 \subset \bigcup Y_1$ and $C_2 \subset \bigcup Y_2$, then $Y_1 \cup Y_2 \in F(\mathcal{G})$ and $C_1 \cup C_2 \subset \bigcup \{Y_1 \cup Y_2\}$ imply that $C_1 \cup C_2 \in \mathcal{C}$. Thus (ii) holds. Hence, $X \in \mathcal{C}$ implies that X is compact. This completes the proof.

I point out that the partitions that are used in the gauge type integrals and other generalizations of the Riemann integral often are not considered to generate closed subrectangles; but, rather are considered to generate “left-closed” or “right-closed” subrectangles. This is what is done by McShane in his definition of the McShane integral. Thus referring to Chapter 5, Section 5.1, replace the set of closed subrectangles obtained by considering the expansion of $([x_{10}, x_{11}] \cup \dots \cup [x_{1k-1}, x_{1k}]) \times \dots \times ([x_{n0}, x_{n1}] \cup \dots \cup [x_{nm-1}, x_{nm}])$ by the collection $\mathcal{A} = \{A_i \mid 1 \leq i \leq m\}$ of n -dimensional right-closed subrectangles obtained by considering the expansion of $((x_{10}, x_{11}] \cup \dots \cup (x_{1k-1}, x_{1k}]) \times \dots \times ((x_{n0}, x_{n1}] \cup \dots \cup (x_{nm-1}, x_{nm}])$. This gives a collection of right-closed rectangles that subdivides $(x_{10}, x_{1k}] \times \dots \times (x_{n0}, x_{nm}] = R_{rc}$ into nonoverlapping right-closed subrectangles. However, when considering an intermediate partition Q , then it is often (but not always) required that each $\vec{x}_i \in Q$ also has the property that $\vec{x}_i \in \overline{A}_i$. This is the case with the δ -fine partitions discussed in Mawhin [1985], and Swartz and Thomson [1988]. This is not the case with the λ -fine partition as defined by McShane.

Let the closed rectangle $R = \overline{R}_{rc} \subset \mathbb{R}^n$. Recall that $B \subset \mathbb{R}^n$ is a nonempty **open n-cell** if $B = (a_1, b_1) \times \dots \times (a_n, b_n)$, $a_i < b_i$, $1 \leq i \leq n$. Let $\mathcal{B}$ be the set of all open n -cells for $\mathbb{R}^n$. In order

to establish results with the least amount of notation, call a map $\lambda: \mathbb{R}^n \rightarrow \mathcal{B}$ a **L -gauge** if for each $\vec{x} \in \mathbb{R}^n$, $\vec{x} \in \lambda(\vec{x})$. (In all cases and without further mention, when one concentrates upon a domain such as R then the gauge is considered to be the above map restricted to R .) Intuitively, think of a L -gauge as simply carving out specific open n -cell neighborhoods for each member of $\mathbb{R}^n$. Now there is associated with gauge integration the pairs (Q, P) which are termed a **P-partition of R** if

(i) $P = \{A_1, \dots, A_k\}$ is a set of right-closed subrectangles that subdivides R_{rc} into nonoverlapping right-closed subrectangles in the usual manner and

(ii) the finite sequence $Q = \{\vec{x}_1, \dots, \vec{x}_k\}$ has the properties that each $\vec{x}_i \in R$. More significantly, however, are the **λ -fine partitions** that are P-partitions with the additional property that

(iii) each $A_i \subset \lambda(\vec{x}_i)$, for $1 \leq i \leq k$, where λ is a L -gauge.

What needs to be determined, however, is whether or not there exists at least one λ -fine partition for a given L -gauge.

Proposition 8.2. *Let $\lambda: \mathbb{R}^n \rightarrow \mathcal{B}$ be an L -gauge and $R \subset \mathbb{R}^n$. Then there exists at least one λ -fine partition of R .*

Proof. Let $\mathcal{C}$ be the set of all (closed) subrectangles of compact R for which there exists an λ -fine partition. First, we show that $\mathcal{C}$ satisfies (i) of Proposition 8.1. Let $\vec{x} \in R$. Then $\vec{x} \in \lambda(\vec{x})$ and $\lambda(\vec{x})$ is an open neighborhood of $\vec{x}$. Hence, $R \cap \lambda(\vec{x})$ is a nonempty R -open neighborhood of $\vec{x}$. Moreover, since there exists a closed subrectangle $I \subset \mathbb{R}^n$ such that $\vec{x} \in \text{int}_{\mathbb{R}^n}(I) \subset I \subset \lambda(\vec{x})$, it follows that $\vec{x} \in R \cap \text{int}_{\mathbb{R}^n}(I) \subset \text{int}_R(R \cap I) \subset I \subset \lambda(\vec{x})$. Now simply consider the λ -fine partition $\{(\vec{x}, R_{rc} \cap I_{rc})\}$ of $(R \cap I)_{rc}$. Since $R \cap I = \overline{(R \cap I)_{rc}}$ is a (closed) subrectangle of R then $R \cap I \in \mathcal{C}$ and $\mathcal{C}$ satisfies (i) of Proposition 8.1.

Let $C_1, C_2 \in \mathcal{C}$ and $C_1 \cap C_2 \neq \emptyset$. Let $P_\lambda = \{A_1, \dots, A_k\}$ and $P'_\lambda = \{A'_1, \dots, A'_p\}$ be the two subdividing collections of right-closed subrectangles and Q_λ, Q'_λ the corresponding finite sequences for C_1 and C_2 , respectively. These exist from the definition of $\mathcal{C}$. Now $P_\lambda = \{A_1, \dots, A_k\}$ and $P'_\lambda = \{A'_1, \dots, A'_p\}$ are generated by partition P and P' , respectively. Considering the common refinement P^* then it follows, since any new right-closed subrectangle generated by this common refinement is a subset of some member of $P_\lambda \cup P'_\lambda$, that a finite sequence of members of $Q_\lambda \cup Q'_\lambda$ may be chosen in such a manner that selection yields a possible new pair $(P_\lambda^*, Q_\lambda^*)$ which forms a λ -fine partition for the closed subrectangle $C_1 \cup C_2$. Thus $\mathcal{C}$ satisfies part (ii) of Proposition 8.1. Since R is compact then $R \in \mathcal{C}$ and the proof is complete.

In order to show the existence of a L -micropartition, a special type of internal function, $\hat{\lambda}$, termed a **L-microgauge** is needed.

Proposition 8.3. *Let $L(\mathbb{R}^n, \mathcal{B})$ be the set of all L -gauges. Then there exists an internal map $\hat{\lambda}: {}^*\mathbb{R}^n \rightarrow {}^*\mathcal{B}$ such that for each $\vec{x} \in {}^*\mathbb{R}^n$ and every $\lambda \in L(\mathbb{R}^n, \mathcal{B})$,*

$$\hat{\lambda}(\vec{x}) \subset {}^*\lambda(\vec{x}).$$

Proof. Consider the internal binary relation

$$B = \{(x, y) \mid x, y \in {}^*(L(\mathbb{R}^n, \mathcal{B})) \wedge \forall z(z \in {}^*\mathbb{R}^n \rightarrow y(z) \subset x(z))\}.$$

We show that B is concurrent, at least, on the set ${}^\sigma(L(\mathbb{R}^n, \mathcal{B}))$. Let $\{(x_1, y_1), \dots, (x_n, y_n)\} \subset B$ and $\{x_1, \dots, x_n\} \subset {}^\sigma(L(\mathbb{R}^n, \mathcal{B})) = \{{}^*\lambda \mid \lambda \in L(\mathbb{R}^n, \mathcal{B})\}$. By $*$ -transfer of the result that for any

$\vec{x} \in \mathbb{R}^n$ the intersection of any nonempty finite subset of $\mathcal{B}$ each member of which contains $\vec{x}$ is itself a member of $\mathcal{B}$, we can define the internal map $f: {}^*\mathbb{R}^n \rightarrow {}^*\mathcal{B}$ as follows: For each $\vec{x} \in {}^*\mathbb{R}^n$ let $f(\vec{x}) = x_1(\vec{x}) \cap \dots \cap x_n(\vec{x})$. This yields $f \in {}^*(L(\mathbb{R}^n, \mathcal{B}))$ and $\{(x_1, f), \dots, (x_n, f)\} \subset B$. Consequently, B is concurrent, at least, on ${}^\sigma(L(\mathbb{R}^n, \mathcal{B}))$. But, we are working in a polysaturated enlargement. This implies that there exists some internal $\hat{\lambda} \in {}^*(L(\mathbb{R}^n, \mathcal{B}))$ such that for each $\vec{x} \in {}^*\mathbb{R}^n$ and each $\lambda \in L(\mathbb{R}^n, \mathcal{B})$ it follows that $\hat{\lambda}(\vec{x}) \subset {}^*\lambda(\vec{x})$ and completes the proof.

The object $\hat{\lambda}$ that exists by Proposition 8.3 is called a **L-microgauge** and has many interesting properties. For example, if $\vec{p} \in R$, then $\hat{\lambda}(\vec{p}) \subset \mu(\vec{p})$, but $\hat{\lambda}(\vec{p}) \neq \mu(\vec{p})$ since $\hat{\lambda}(\vec{p})$ is an internal $*$ -open member of ${}^*\mathcal{B}$; while $\mu(\vec{p})$ is an external set which is the union of $*$ -open members of ${}^*\mathcal{B}$. [Herrmann [1978]] However, our major use of a L-microgauge $\hat{\lambda}$, in this appendix, is to obtain a **L-micropartition** $(Q_{\hat{\lambda}}, P_{\hat{\lambda}})$ which now exists by $*$ -transfer of the results in Proposition 8.2. Observe that $(Q_{\hat{\lambda}}, P_{\hat{\lambda}})$ is also a $*$ λ -fine partition for each standard L -gauge λ since each internal $A_i \subset \hat{\lambda}(\vec{x}_i) \subset {}^*\lambda(\vec{x}_i)$.

McShane defined his integral in the following manner. Let λ be a L -gauge. Then there exists a map $\mathcal{L}$ such that $\mathcal{L}(\lambda)$ is the nonempty set of all λ -fine partitions of R . Let $f: R \rightarrow \mathbb{R}$. Then f is **McShane integrable over R** if there exists a real number J such that: for each positive real number r there corresponds a L -gauge λ such that for each $(Q_\lambda, P_\lambda) \in \mathcal{L}(\lambda)$,

$$\left| \sum_{i=1}^k f(\vec{x}_i) v(A_i) - J \right| < r,$$

where $v(A_i) = v(\overline{A_i})$. Since McShane's integral is equivalent to the Lebesgue integral, our final results in this particular appendix are stated in terms of the Lebesgue integral.

Proposition 8.4. *If $f: R \rightarrow \mathbb{R}$ is Lebesgue integrable with $\int_R f(\vec{x}) dX = J$, then for each L -microgauge $\hat{\lambda}$ and for each $(Q_{\hat{\lambda}}, P_{\hat{\lambda}}) \in {}^*\mathcal{L}(\hat{\lambda})$,*

$$\sum_{i=1}^\Gamma {}^*f(\vec{x}_i) {}^*v(A_i) \in \mu(J).$$

Proof. Suppose that $f: R \rightarrow \mathbb{R}$ is Lebesgue integrable with $\int_R f(\vec{x}) dX = J$. Consider arbitrary positive real r . The following sentence

$$\exists x (\forall y (x \in L(\mathbb{R}^n, \mathcal{B}) \wedge y \in \mathcal{L}(x) \rightarrow \left| \sum_{i=1}^k f(\vec{x}_i) v(A_i) - J \right| < r))$$

holds in ${}^*\mathcal{M}$ by $*$ -transfer. Now let $\hat{\lambda}$ be a L-microgauge and $(Q_{\hat{\lambda}}, P_{\hat{\lambda}}) \in {}^*\mathcal{L}(\hat{\lambda})$. From the above observation $(Q_{\hat{\lambda}}, P_{\hat{\lambda}}) \in {}^*\mathcal{L}({}^*\lambda)$ for any standard L -gauge λ . Thus, independent of any particular L -gauge and hence any particular positive real r

$$\left| \sum_{i=1}^\Gamma {}^*f(\vec{x}_i) {}^*v(A_i) - J \right| < r,$$

for each $(Q_{\hat{\lambda}}, P_{\hat{\lambda}}) \in {}^*\mathcal{L}(\hat{\lambda})$. Consequently, $\sum_{i=1}^\Gamma {}^*f(\vec{x}_i) {}^*v(A_i) \in \mu(J)$ for each $(Q_{\hat{\lambda}}, P_{\hat{\lambda}}) \in {}^*\mathcal{L}(\hat{\lambda})$ and the proof is complete.

Proposition 8.5. *Suppose that $f: R \rightarrow \mathbb{R}$ and there exists a real number J , and a L -microgauge $\hat{\lambda}$ such that for each $(Q_{\hat{\lambda}}, P_{\hat{\lambda}}) \in {}^*\mathcal{L}(\hat{\lambda})$,*

$$\sum_{i=1}^{\Gamma} {}^*f(\vec{x}_i) {}^*v(A_i) \in \mu(J).$$

Then f is Lebesgue integrable on R and $\int_R f(\vec{x}) dX = J$.

Proof. Let r be any positive real number. Then $\left| \sum_{i=1}^{\Gamma} {}^*f(\vec{x}_i) {}^*v(A_i) - J \right| < r$. Thus the sentence

$$\exists x (\forall y (x \in {}^*(L(\mathbb{R}^n, \mathcal{B})) \wedge y \in {}^*\mathcal{L}(x) \rightarrow \left| \sum_{i=1}^k {}^*f(\vec{x}_i) {}^*v(A_i) - J \right| < r))$$

holds in ${}^*\mathcal{M}$, hence, in $\mathcal{M}$ and the result follows.

Appendix For Chapter 10.

Proposition 10.1 *Let standard $A \in \mathcal{H}$. Then ${}^\sigma A$ is internal if and only if A is finite.*

Proof. If A is empty, then ${}^*A = {}^\sigma A = \emptyset$ and thus ${}^\sigma A$ is internal. If $A = \{a_1, \dots, a_k\}$, a nonempty finite set, then ${}^*A = \{{}^*a_1, \dots, {}^*a_k\} = {}^\sigma A$ is internal. Conversely, assume that A is infinite. Then there some $B \subset A$ and a bijection $f: B \rightarrow \mathbb{N}$. Then for each $b \in B$ and there exists $n \in \mathbb{N}$ such that $({}^*b, {}^*n) \in {}^*f$. Further, for each $n \in \mathbb{N}$ there exists some $b \in B$ such that $({}^*b, {}^*n) \in {}^*f$ from Theorem 3.1.3 part (v). Hence, ${}^*f[{}^\sigma B] = {}^\sigma N$. If ${}^\sigma A$ is internal, then ${}^\sigma A \cap {}^*B = {}^\sigma B$ is internal. But, the image of an internal set under an internal map is internal. This contradicts the result that ${}^\sigma N$ is external.

The following is the very important **Cauchy Principle** for normed linear spaces.

Theorem 10.1.1. (Cauchy Principle) *Let $P(x)$ be a bounded formal first-order expression in one free variable and employing internal constants (i.e. a bounded formal internal property as discussed in Appendix 4, within the proof of Theorem 4.2.2.) If $P(\epsilon)$ holds for each $\epsilon \in \mu^n(\vec{0})$, then there exists an open ball B about $\vec{0}$ such that $P(\vec{x})$ holds for each $\vec{x} \in {}^*B$.*

Proof. (In the following proof, as an illustration, we retain the σ notation rather than assume the identification. This will show how the embedding technically allows one to go back-and-forth between the standard model and the σ model, in many cases. Some modern texts in this subject do not utilize our identification process for objects in $\mathfrak{R}$.) Let $\Phi(z) = z \in ({}^*\mathbb{N} - \{0\}) \wedge \forall y (y \in {}^*\mathbb{R}^n \wedge \|y\| < 1/z \rightarrow P(y))$. Then $\Phi(z)$ is a bounded internal first-order expression and from the hypothesis $\Phi(z)$ holds for each $n \in \mathbb{N}_\infty$. Let $A = \{z \mid z \in {}^*\mathbb{N} \wedge \neg \Phi(z)\}$. From the internal definition theorem, A is an internal subset of ${}^*\mathbb{N}$ and $A \subset {}^\sigma \mathbb{N}$. Thus $A = {}^\sigma C$, where $C \subset \mathbb{N}$. From proposition 10.1, C is finite. Thus C has a greatest member, say k . Hence $\Phi(z)$ holds for each $n \in {}^*\mathbb{N}$ such that $n \geq {}^*k + {}^*1$. Now simply consider the standard ball B about $\vec{0}$ with radius $1/(k+1)$. Then ${}^*B$ is but the ball in ${}^*\mathbb{R}^n$ about ${}^*\vec{0}$ with radius $1/(k+1) = 1/({}^*k + {}^*1)$.

Theorem 10.1.2. *Two internal maps f and g are equivalent if and only if there is a local map α such that for each nonzero $\vec{x} \in \mu^n(\vec{0})$*

$$f(\vec{x}) = g(\vec{x}) + (\alpha(\vec{x}))\|\vec{x}\|.$$

Proof. Define

$$\alpha(\vec{x}) = \frac{f(\vec{x}) - g(\vec{x})}{\|\vec{x}\|},$$

for nonzero $\vec{x} \in \mu^n(\vec{0})$, and $\alpha(\vec{0}) = \vec{0}$. Let $f \sim g$. Then for each nonzero $\vec{x} \in \mu^n(\vec{0})$ we have that $(f(\vec{x}) - g(\vec{x}))/\|\vec{x}\| \in \mu^m(\vec{0}) \Rightarrow \alpha$ is a local map.

Conversely, assume that for nonzero $\vec{x} \in \mu^n(\vec{0})$ that $f(\vec{x}) = g(\vec{x}) + (\alpha(\vec{x}))\|\vec{x}\|$, where α is a local map. The result follows from the definition of a local map.

If the maps f and g of Theorem 10.1.2 are locally linear maps, then the equation in the conclusion of Theorem 10.1.2 holds for $\vec{x} = \vec{0}$ as well.

Theorem 10.1.3. *Suppose that f is any locally linear map. Then there exists a unique internal linear map $T_f: {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}^m$ such that $\|T\| \in \mathcal{O}$ and there exists an open $E \subset \mathbb{R}^n$, where $\vec{0} \in E$, such that $f(\vec{x}) = T_f(\vec{x})$ for each $\vec{x} \in {}^*E$.*

Proof. First, we define an internal function by means of the Internal Definition Theorem. Let fixed $\Lambda \in \mathbb{R}_\infty$ and $\epsilon = 1/\Lambda$. Notice that for nonzero $\vec{x} \in {}^*\mathbb{R}^n$, $\epsilon(\vec{x}/\|\vec{x}\|) \in \mu^n(\vec{0})$. Next, consider

$$T_f = \{(x, y) \mid x \in {}^*\mathbb{R}^n \wedge x \neq \vec{0} \wedge y \in {}^*\mathbb{R}^m \wedge \\ y = \Lambda\|x\|f\left(\epsilon\left(\frac{x}{\|x\|}\right)\right)\} \cup (\vec{0}, \vec{0}).$$

We know that for any $\vec{x} \in {}^*\mathbb{R}^n$ there is a nonzero $\alpha \in \mu(0)$ such that $\alpha\vec{x} \in \mu^n(\vec{0})$. Consider any $\vec{x} \in {}^*\mathbb{R}^n$ and an $\alpha \in \mu(0)$ such that $\alpha\vec{x} \in \mu^n(\vec{0})$. Then

$$T_f(\vec{x}) = \alpha\|\vec{x}\| \left(\frac{f\left(\epsilon\frac{\vec{x}}{\|\vec{x}\|}\right)}{\alpha\epsilon} \right) = \\ \frac{f\left(\frac{\alpha\epsilon\|\vec{x}\|\vec{x}}{\|\vec{x}\|}\right)}{\alpha\epsilon} = \frac{f(\alpha\epsilon\vec{x})}{\alpha\epsilon} = \frac{f(\epsilon\alpha\vec{x})}{\alpha\epsilon} = \frac{\epsilon f(\alpha\vec{x})}{\epsilon\alpha} = \frac{f(\alpha\vec{x})}{\alpha}. \quad (*)$$

Obviously, if $\vec{x} \in \mu^n(\vec{0})$, then taking any nonzero $\alpha \in \mu(0)$ we have that

$$T_f(\vec{x}) = \frac{f(\alpha\vec{x})}{\alpha} = \frac{\alpha f(\vec{x})}{\alpha} = f(\vec{x}).$$

Consider the bounded internal first-order expression $P(x) = x \in {}^*\mathbb{R}^n \wedge T_f(x) = f(x)$. Since $P(x)$ holds for all $\vec{x} \in \mu^n(\vec{0})$ then the Cauchy Principle implies that there exists an open ball E about $\vec{0}$ such that for each $\vec{x} \in {}^*E$, $T_f(\vec{x}) = f(\vec{x})$.

To establish that T_f is a linear transformation, let $\vec{x}, \vec{y} \in {}^*\mathbb{R}^n$ and $\alpha \in \mu(0)$ be such that $\alpha\vec{x}, \alpha\vec{y} \in \mu^n(\vec{0})$. Then using (*) and the locally linear property it follows that

$$T_f(\vec{x} + \vec{y}) = \frac{1}{\alpha}(f(\alpha\vec{x} + \alpha\vec{y})) = \frac{1}{\alpha}(f(\alpha\vec{x}) + f(\alpha\vec{y})) = T_f(\vec{x}) + T_f(\vec{y}).$$

Now let $r \in {}^*\mathbb{R}$. Then there is a nonzero $\alpha \in \mu(0)$ such that $\alpha\vec{x}, r\alpha\vec{x} \in \mu^n(\vec{0})$. Consequently,

$$T_f(r\vec{x}) = \frac{1}{\alpha}f(r\alpha\vec{x}) = \frac{r}{\alpha}f(\alpha\vec{x}) = rT_f(\vec{x}).$$

For uniqueness, let $f(\vec{x}) = T(\vec{x}) = G(\vec{x})$ for each $\vec{x} \in \mu^n(\vec{0})$, where T, G are internal linear transformations from ${}^*\mathbb{R}^n$ into ${}^*\mathbb{R}^m$. [Note: We, of course, mean that they are linear transformations over the vector space ${}^*\mathbb{R}^n$ into ${}^*\mathbb{R}^m$ where the field in question is ${}^*\mathbb{R}$.] Then by linearity for any $\vec{x} \in {}^*\mathbb{R}^n$

$$T(\vec{x}) = \Lambda\|\vec{x}\|T\left(\epsilon\left(\frac{\vec{x}}{\|\vec{x}\|}\right)\right) = f(\vec{x}) = \Lambda\|\vec{x}\|G\left(\epsilon\left(\frac{\vec{x}}{\|\vec{x}\|}\right)\right) = G(\vec{x}).$$

Lastly, we show that $\|T_f\|$ is limited. Recall that if T is a standard linear transformation, then $\|T\| = \sup\{T(\vec{x}) \mid \vec{x} \in \mathbb{R}^n \wedge \|\vec{x}\| = 1\}$. The ${}^*\sup$ operator extends, in general, to internal subsets of ${}^*\mathbb{R}$ and, indeed, is the same operator as a sup defined on some subsets of the ordered field ${}^*\mathbb{R}$ even though it may not exist for certain bounded sets such as $\mu(0)$. Assume that $\|T_f\|$ is not limited. Then by * -transfer of the standard case, there is some $\vec{x} \in {}^*\mathbb{R}^n$, $\|\vec{x}\| = 1$, such that $\|T_f(\vec{x})\|$ is infinite since the set $\{T_f(\vec{x}) \mid \vec{x} \in \mathbb{R}^n \wedge \|\vec{x}\| = 1\}$ is internal and not * -bounded. Consequently, $\vec{x}/\|T_f(\vec{x})\| \in \mu^n(\vec{0})$. From (*) one obtains

$$T_f(\vec{x}) = \|T_f(\vec{x})\|f\left(\frac{\vec{x}}{\|T_f(\vec{x})\|}\right),$$

which implies that

$$\|T_f(\vec{x})\| = \|T_f(\vec{x})\| \left\| f\left(\frac{\vec{x}}{\|T_f(\vec{x})\|}\right) \right\|.$$

Hence, $\|f(\vec{x}/\|T_f(\vec{x})\|)\| = 1 \notin \mu(0)$ implies the contradiction that f is not a local map and this completes the proof.

Theorem 10.1.4. *Let f, g be locally linear maps. Then $f \sim g$ if and only if $T_f(\vec{x}) \approx T_g(\vec{x})$ for each $\vec{x} \in {}^*\mathbb{R}^n$ such that $\|\vec{x}\| = 1$.*

Proof. First, assume that $T_f(\vec{x}) \approx T_g(\vec{x})$ for each $\vec{x} \in {}^*\mathbb{R}^n$ such that $\|\vec{x}\| = 1$. Consider any $\vec{y} \in \mu^n(\vec{0})$. Then from Theorem 10.1.3, it follows that

$$\frac{f(\vec{y}) - g(\vec{y})}{\|\vec{y}\|} = \frac{T_f(\vec{y}) - T_g(\vec{y})}{\|\vec{y}\|} = T_f\left(\frac{\vec{y}}{\|\vec{y}\|}\right) - T_g\left(\frac{\vec{y}}{\|\vec{y}\|}\right) \in \mu^m(\vec{0}).$$

Thus $f \sim g$.

Conversely, assume that $f \sim g$, $\|\vec{x}\| = 1$ and that positive $\alpha \in \mu(0)$. Then from theorem 10.1.3,

$$T_f(\vec{x}) - T_g(\vec{x}) = \frac{1}{\alpha}(T_f(\alpha\vec{x}) - T_g(\alpha\vec{x})) = \frac{f(\alpha\vec{x}) - g(\alpha\vec{x})}{\|\alpha\vec{x}\|} \in \mu^m(\vec{0}).$$

Proposition 10.2. *Let $F, G: \mathcal{V} \rightarrow \mathcal{W}$ be two linear transformations and $\mathcal{V}, \mathcal{W}$ be (nontrivial) linear spaces over the real or complex fields, where $\mathcal{V}$ is normed. If there exists an open ball $B \subset \mathcal{V}$, about $\vec{0}$, such that $F(\vec{x}) = G(\vec{x})$ for each $\vec{x} \in B$, then $F = G$.*

Proof. Let nonzero $\vec{x} \in \mathcal{V}$, and positive $r \in \mathbb{R}$ be the radius of the ball B . Then $(r\vec{x})/2\|\vec{x}\| \in B$ and $T((r\vec{x})/(2\|\vec{x}\|)) = (r/(2\|\vec{x}\|))T(\vec{x}) = G((r\vec{x})/(2\|\vec{x}\|)) = (r/(2\|\vec{x}\|))G(\vec{x}) \Rightarrow T(\vec{x}) = G(\vec{x})$. Obviously, $\vec{0} = T(\vec{0}) = G(\vec{0})$ and the proof is complete.

Theorem 10.1.5. *If f is a differential, then there exists a unique bounded linear transformation $F: \mathbb{R}^n \rightarrow \mathbb{R}^m$ and an open set $E \subset \mathbb{R}^n$ such that $\vec{0} \in E$ and ${}^*F(\vec{x}) = f(\vec{x})$ for all $\vec{x} \in {}^*E$ and, in particular, $F(\vec{x}) = f(\vec{x})$ for each $\vec{x} \in E$.*

Proof. Since f is a differential then there exists a bounded linear transformation $F: \mathbb{R}^n \rightarrow \mathbb{R}^m$ and a unique internal linear transformation $T_f: {}^*\mathbb{R}^n \rightarrow {}^*\mathbb{R}^m$ such that ${}^*F(\vec{x}) = f(\vec{x}) = T_f(\vec{x})$ for each $\vec{x} \in \mu^n(\vec{0})$. Consider the bounded internal first-order expression $P(x) = x \in {}^*\mathbb{R}^n \wedge {}^*F(x) = f(x) = T_f(x)$. Then since $P(x)$ holds for all $\vec{x} \in \mu^n(\vec{0})$ then the Cauchy Principle implies that there exists some open ball E about $\vec{0}$ such that for each $\vec{x} \in {}^*E$, ${}^*F(\vec{x}) = f(\vec{x}) = T_f(\vec{x})$. Hence, in particular, ${}^*F(\vec{x}) = F(\vec{x}) = f(\vec{x})$ for each $\vec{x} \in E$. The fact that F is unique comes from the * -transfer of Proposition 10.2.

Theorem 10.1.6. *If f and g are differentials and $f \sim g$, then there exists some open $E \in \mathbb{R}^n$ such that $\vec{0} \in E$ and $f(\vec{x}) = g(\vec{x})$ for each $\vec{x} \in E$.*

Proof. There are two bounded standard transformations $T, G: \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that $f(\vec{x}) = {}^*T(\vec{x})$ and $g(\vec{x}) = {}^*G(\vec{x})$ for each $\vec{x} \in \mu^n(\vec{0})$. By the uniqueness property of Theorem 10.1.3 and Theorem 10.1.4 ${}^*T(\vec{y}) \approx {}^*G(\vec{y})$ for each $\vec{y} \in {}^*\mathbb{R}^n$ such that $\|\vec{y}\| = 1$. Considering any $\vec{z} \in \mathbb{R}^n$ such that $\|\vec{z}\| = 1$ and taking the standard part operator this yields $T(\vec{z}) = G(\vec{z})$. Taking any nonzero $\vec{x} \in \mathbb{R}^n$, we have that

$$T(\vec{x}) = \|\vec{x}\|T\left(\frac{\vec{x}}{\|\vec{x}\|}\right) = \|\vec{x}\|G\left(\frac{\vec{x}}{\|\vec{x}\|}\right) = G(\vec{x}).$$

Thus $f(\vec{x}) = g(\vec{x})$ for each $\vec{x} \in \mu^n(\vec{0})$. Once again application of the Cauchy Principle yields the result.

Theorem 10.1.7. *Let nonempty open $G \subset \mathbb{R}^n$, and $\vec{c} \in G$. A function $f: G \rightarrow \mathbb{R}^m$ is continuous at $c \in G$ if and only if the local increment map $\Delta^* f_{\vec{c}}$ is a local map.*

Proof. This follows immediately from Definition 4.4.1 and the fact that if G is open and $\vec{c} \in G$, then $\mu^n(\vec{c}) \subset {}^*G$.

Theorem 10.1.8. *Let noninfinitesimal $\vec{a} \in \mathcal{O}^n$ and suppose that $\vec{a} - \vec{b} \in \mu^n(\vec{0})$. Then $\vec{a}$ is almost parallel to $\vec{b}$.*

Proof. Since $\vec{a} \in \mathcal{O}^n$ and $\vec{0} \neq \vec{a}$ then there exists some nonzero $\vec{c} \in \mathbb{R}^n$ such that $\vec{a} \in \mu^n(\vec{c})$. Consequently, $\vec{b} \in \mu^n(\vec{c})$. Further, $\text{st}(\|\vec{a}\|), \text{st}(\|\vec{b}\|) \neq 0$ and $|\|\vec{a}\| - \|\vec{b}\|| \leq \|\vec{c} - \vec{b}\| \Rightarrow \text{st}(\|\vec{a}\|) = \text{st}(\|\vec{b}\|)$. Hence,

$$\frac{\text{st}(\vec{a})}{\text{st}(\|\vec{a}\|)} = \text{st}\left(\frac{\vec{a}}{\|\vec{a}\|}\right) = \frac{\text{st}(\vec{b})}{\text{st}(\|\vec{b}\|)} = \text{st}\left(\frac{\vec{b}}{\|\vec{b}\|}\right)$$

and the proof is complete.

Theorem 10.4.1. *Let $f: G \rightarrow \mathbb{R}$, where nonempty open $G \subset \mathbb{R}^n$ and standard $\vec{v} \in G$. Suppose that $y = x_i$, $1 \leq i \leq n$, f_y is defined on G and continuous at $\vec{v}$. Let $C_y \subset {}^*G$, where internal C_y is $*$ -convex in the direction y . If for $h \in \mu(0)$, such that $\vec{p} = (p_1, \dots, p_i, \dots, p_n), \vec{q} = (p_1, \dots, p_i + h, \dots, p_n) \in C_y$ and $\vec{v} \approx \vec{p}$, then there exists $\epsilon \in \mu(0)$ such that*

$${}^*f(\vec{q}) - {}^*f(\vec{p}) = f_y(\vec{v})h + h\epsilon.$$

Proof. Recall that for the hypothesized behavior of f and f_y the mean value theorem for the partial derivative f_y states that for any set $D_y \subset G$ convex in the direction y if $\vec{u} = (u_1, \dots, u_i, \dots, u_n), \vec{w} = (u_1, \dots, u_i + r, \dots, u_n) \in D_y, r \in \mathbb{R}$ then there exists $\vec{s}$ in the line segment with end points $\vec{u}, \vec{w}$ such that $f(\vec{w}) - f(\vec{u}) = r f_y(\vec{s})$. Thus $\vec{s} \in D_y$ and $\|\vec{u} - \vec{s}\| \leq \|\vec{u} - \vec{w}\| = |r|$. By $*$ -transfer it follows that there exists some $\vec{t} \in C_y$ such that $\vec{t} \in C_y$ and $\|\vec{p} - \vec{t}\| \leq |h|$ and ${}^*f(\vec{q}) - {}^*f(\vec{p}) = h {}^*f_y(\vec{t})$. But, $\vec{t} \approx \vec{p} \approx \vec{v}$ and the continuity of f_y at $\vec{v}$ imply that ${}^*f_y(\vec{t}) \approx f_y(\vec{v})$. Thus there exists some $\epsilon \in \mu(0)$ such that ${}^*f_y(\vec{t}) = f_y(\vec{v}) + \epsilon$. The result follows by substitution.

NOTES

[1] **Theorem N.1.** *Two unit vectors, $\vec{v}, \vec{u} \in {}^*\mathbb{R}^n$ have the property that $\vec{v} \approx \pm \vec{u}$ if and only if $\vec{v} \bullet \vec{u} \approx \pm 1$.*

Proof. First note that $\vec{v}, \vec{u} \in \mathcal{O}^n$. For the necessity, assume that $\vec{v} \approx \pm \vec{u}$. Then $\vec{v} \pm \vec{u} \in \mu^n(\vec{0}) \Rightarrow \vec{v} \bullet \vec{v} \pm \vec{u} \bullet \vec{v} = 1 \pm \vec{u} \bullet \vec{v} \in \mu(0) \Rightarrow \pm \vec{u} \bullet \vec{v} \approx 1 \Rightarrow \vec{u} \bullet \vec{v} \approx \pm 1$.

For the converse, assume that $\vec{v} \bullet \pm \vec{u} \approx 1$ but that $\vec{v} \not\approx \pm \vec{u}$. Thus there exist $\vec{0} \neq \vec{r}_{\pm} \in \mathbb{R}^n$ and $\epsilon_{\pm} \in \mu^n(\vec{0})$ such that $\vec{v} = \pm \vec{u} + \vec{r}_{\pm} + \epsilon_{\pm}$. Hence, $\vec{v} \bullet \pm \vec{u} = \pm \vec{u} \bullet \pm \vec{u} + \vec{r}_{\pm} \bullet \pm \vec{u} + \epsilon_{\pm} \bullet \pm \vec{u} = 1 + \vec{r}_{\pm} \bullet \pm \vec{u} + \epsilon_{\pm} \bullet \pm \vec{u} \approx 1 + \vec{r}_{\pm} \bullet \pm \vec{u}$.

We now show that $\vec{r}_{\pm} \bullet \pm \vec{u} \not\approx 0$. We know that $\vec{r}_{\pm} \bullet \pm \vec{u} = (1/4)(\|\pm \vec{u} + \vec{r}_{\pm}\|^2 + \|\pm \vec{u} - \vec{r}_{\pm}\|^2)$. Assume that $\vec{r}_{\pm} \bullet \pm \vec{u} \approx 0 \Rightarrow \|\pm \vec{u} + \vec{r}_{\pm}\|^2 \approx \|\pm \vec{u} - \vec{r}_{\pm}\|^2$. Now $\vec{v} = \pm \vec{u} + \vec{r}_{\pm} + \epsilon_{\pm} \Rightarrow \|\vec{v}\|^2 = 1 \approx$

$\|\pm \vec{u} + \vec{r}_\pm\|^2 \Rightarrow 2 \approx \|\pm \vec{u} + \vec{r}_\pm\|^2 + \|\pm \vec{u} - \vec{r}_\pm\|^2 = 2\|\pm \vec{u}\|^2 + 2\|\vec{r}_\pm\|^2 = 2 + 2\|\vec{r}_\pm\|^2 \Rightarrow 0 \approx \|\vec{r}_\pm\|$; a contradiction. Thus $\vec{r}_\pm \bullet \pm \vec{u} \not\approx 0 \Rightarrow \vec{v} \bullet \pm \vec{u} \not\approx 1$. The result follows from this contradiction.

[2] Let $c: [a, b] \rightarrow \mathbb{R}^n$ be a continuous differentiable curve with graph C and $c'(t) \neq \vec{0}$ for each $t \in [a, b]$. Assume that uniformly continuous $F: E \rightarrow \mathbb{R}^n$, open $E \supset C$. Then $*F(\ell_j(t')) \bullet *v_j = *F(*c(t')) \bullet *v_j + \epsilon_j \|v_j\|$, $\epsilon_j \in \mu(0)$, and $\sum_{j=1}^\Gamma *F(\ell_j(t')) \bullet *v_j \approx \sum_{j=1}^\Gamma *F(*c(t')) \bullet *v_j$, where the symbols have the same meaning as in the derivation for Application 8.2.1.

Proof. For an interior point of $t \in (a, b)$ we use the concept of uniform differentiability and the result [Stroyan and Luxemburg [1976:94-97]] that for any t_{j-1} , $t_j \approx t$ that there exists some $\vec{e}_j \in \mu^n(\vec{0})$ such that $\vec{v}_j = *c(t_j) - *c(t_{j-1}) = c'(t)(t_j - t_{j-1}) + |t_j - t_{j-1}|\vec{e}_j$. Assuming that $*c(t_j) - *c(t_{j-1}) = \vec{0} \Rightarrow c'(t)(t_j - t_{j-1}) + |t_j - t_{j-1}|\vec{e}_j = \vec{0} \Rightarrow c'(t)((t_j - t_{j-1})/|t_j - t_{j-1}|) + \vec{e}_j = \vec{0} \Rightarrow \|c'(t)((t_j - t_{j-1})/|t_j - t_{j-1}|)\| = \|\vec{e}_j\| \Rightarrow \text{st}(\|c'(t)\|) = \|c'(t)\| = \text{st}(\|\vec{e}_j\|) = 0$; a contradiction. Thus $\vec{v}_j = *c(t_j) - *c(t_{j-1}) \neq \vec{0}$. Now the same proof that appears in Stroyan and Luxemburg [1976:94-97] shows that this result also holds if $t = a$ or $t = b$. Next observe that $\ell_j(t') \approx \ell_j(t_j) = *c(t_j) \approx *c(t') \approx c(t)$, where $t \in [a, b]$. Thus $\ell_j(t')$, $*c(t') \in \mu^n(c(t)) \subset *E$ and the uniform continuity of $*F \Rightarrow *F(\ell_j(t')) = *F(*c(t')) + \vec{\lambda}_j$. Hence, $*F(\ell_j(t')) \bullet (\vec{v}_j/\|\vec{v}_j\|) = *F(*c(t')) \bullet (\vec{v}_j/\|\vec{v}_j\|) + \vec{\lambda}_j \bullet (\vec{v}_j/\|\vec{v}_j\|) = *F(*c(t')) \bullet (\vec{v}_j/\|\vec{v}_j\|) + \epsilon_j$ and the first result follows.

For the second result follow the usual method and notice that there exists infinitesimal $\delta = \max\{\epsilon_j\}$ and that $|\sum_{j=1}^\Gamma \epsilon_j \|\vec{v}_j\|| \leq |\delta| \sum_{j=1}^\Gamma \|\vec{v}_j\| \Rightarrow \sum_{j=1}^\Gamma \epsilon_j \|\vec{v}_j\| \in \mu(0)$ since $\text{st}(\sum_{j=1}^\Gamma \|\vec{v}_j\|) = \text{length of the curve } c$.

REFERENCES

- Aerts, D.** [1984], The missing elements of reality in the description of quantum mechanics of the E.P.R. paradox situation, *Helvetica Physica*, **57**: 421-428.
- Apostol, Tom M.** [1957], *Mathematical Analysis*, Addison-Wesely, Reading, MA.
- Barwise, Jon (ed.)** [1977], *Handbook of Mathematical Logic*, North-Holland, Amsterdam.
- Berkeley** [1734], *The Analyst*, London.
- Cesari** [1956], *Surface Area*, Princeton University Press, Princeton, NJ.
- Cutland N. J.** [1986], Private communication.
- De Lillo, Nicholas J.** [1982], *Advanced Calculus with Applications*, Macmillan, New York, NY.
- Gauss, Karl F.** [1827], *General Investigation of Curved Surfaces*, Raven Press, Hewlett, NY.
- Henstock, R.** [1961], Definitions of the Riemann type of variational integral, *Proc. London Math. Soc.*, (3) **11**: 402-418.
- Herrmann, Robert A.** [1976], The Q-topology, Whyburn type filters and the cluster set map, *Proc. Amer. Math. Soc.*, **59**: 161-166.
- Herrmann, Robert A.** [1978], (2003) *Nonstandard Analysis - A Simplified Approach*, <http://www.arxiv.org/abs/math.GM/0310351>
- Herrmann, Robert A.** [1980], A nonstandard approach to pseudotopological compactifications, *Z. Math. Logik Grundlagen Math.*, **26**: 361-384.
- Herrmann, Robert A.** [1985], Supernear functions, *Math. Japanica*, **30**: 169-185.
- Herrmann, Robert A.** [1986], [1987], (1993) *The Theory of Ultralogics* <http://www.arxiv.org/abs/math.GM/9903081>
<http://www.arxiv.org/abs/math.GM/9903082>
- Herrmann, Robert A.** [1989], Fractals and ultrasmooth microeffects, *J. Math. Phys.*, **30**(4), April 1989: 805-808.
- Herrmann, Robert A.** [1995], *Nonstandard Analysis Applied to Special and General Relativity - The Theory of Infinitesimal Light-Clocks* <http://arxiv.org/abs/math/0312189>
- Hurd, A.E. and P.A. Loeb** [1985], *An Introduction to Nonstandard Real Analysis*, Academic Press, Orlando, FL.
- Keisler, H. Jerome** [1986], *Elementary Calculus - An Infinitesimal Approach*, (Second edition), Prindle, Weber & Schmidt, Boston.
- Jarnik, J., J. Kurzweil and S. Schwabik** [1983], On Mawhin's approach to multiple nonabsolutely convergent integrals, *Časopis Pěst. Mat.*, **108**: 157-167.
- Jeck, Thomas J.** [1971], *Lectures Notes in St Theory, Lecture Notes in Mathematics #217*, Macmillan, Springer-Verlag, New York, NY.
- Leibniz G. W.** [1701], *Mémoire de M.G.G Leibniz touchant son sentiment sur le calcul différentiel*, *Journal de Trévoux*, Mathematische Schriften, ed. C.I. Gerhardt, Vol. 5 (1858).
- Lorentz, H. A.** [1915], *The Theory of Electrons*, (Dover, New York, NY, 1952.)
- Luxemburg, W.A.J.** [1962], *Non-Standard Analysis - Lectures on A. Robinson's Theory of Infinitesimals and Infinitely Large Numbers*, Math. Dept., California Institute of Technology, Pasadena, CA.
- Luxemburg, W.A.J.** [1973] What is nonstandard analysis? in *Papers in the Foundations of Mathematics*, No. 13 *Slaught Memorial Papers*, Amer. Math. Monthly **80**: 38-67.
- Machove, M. and J. Hirschfeld** [1969], *Lectures on Non-Standard Analysis*, Lecture Notes in Mathematics V.94, Springer-Verlag, New York, NY.
- Mawhin, J.** [1986], Nonstandard analysis and generalized Riemann integrals, *Časopis Pěst. Mat.*,

111:34-47.

Maxwell James Clark [1890] *The Scientific Papers of James Clark Maxwell*, Cambridge University Press, Cambridge (Dover, New York, NY, 1965.)

McShane, E. J. [1973], A unified theory of integration, *Amer. Math. Monthly*, **80**: 349-359.

Morley, Arthur [1942], *Mechanics for Engineers*, Longmans, Green and Co. London.

Newton, Isaac [Summer 1665], The calculus becomes an algorithm, in *The Mathematical Papers of Isaac Newton*, (ed. D.T Whiteside, V. I, Cambridge University Press, New York, NY, 1967): 298-368.

Newton, Isaac [Oct. 1665 - May 1666], The general problem of tangents, curvature and limit-motion analysed by the method of fluxions, in *The Mathematical Papers of Isaac Newton*, (ed. D.T Whiteside, V. I, Cambridge University Press, New York, NY, 1967): 369-399.

Newton, Isaac [Oct. 1666], The October 1666 tract on fluxions, in *The Mathematical Papers of Isaac Newton*, (ed. D.T Whiteside, V. I, Cambridge University Press, New York, NY, 1967): 400-448.

Newton, Isaac [Winter 1670 - 1671], The tract 'De methodis serierum et fluxionum', in *The Mathematical Papers of Isaac Newton*, (ed. D.T Whiteside, V. III, Cambridge University Press, New York, NY, 1969): 32-353.

Newton, Isaac [1686], *Mathematical Principles of Natural Philosophy*, (Revised translation by Florian Cajori, University of California Press, Berkeley, CA, 1934).

Robinson, Abraham [1961], Non-standard analysis, *Nederl. Akad. Wetensch. Proc. Ser. A 64, and Indag. Math.* **23**: 432-440.

Robinson, Abraham [1966], *Non-standard Analysis*, North-Holland, Amsterdam.

Robinson, A. and E. Zakon [1969], A set-theoretic characterization of enlargements, in *Applications of Model Theory to Algebra, Analysis, and Probability* (ed. W.A.J. Luxemburg, Holt, Rinehart and Winston, New York, NY): 109-122.

Sears, F. W. and M. W. Zemansky [1952], *College Physics*, Addison-Wesley, Cambridge, MA.

Simhony, M. [1987], *The Electron-Positron Lattice Space*, Physics Section 5, The Hebrew University, Jerusalem.

Spivak, Michael [1965], *Calculus on Manifolds*, W. A. Benjamin, New York, NY.

Stroyan, K.D. and W.A.J. Luxemburg [1976], *Introduction to the Theory of Infinitesimals*, Academic Press, New York, NY.

Struik, D. J. [1961], *Differential Geometry*, Addison-Wesley, Reading, MA.

Suppes, Patrick [1960], *Axiomatic Set Theory*, D. Von Nostrand, New York, NY.

Swartz, Charles and Brian S. Thomson [1988], More on the Fundamental Theorem of Calculus, *Amer. Math. Monthly*, **95**: 644-648.

Synge, J. L and B. A. Griffith [1959], *Principles of Mechanics*, McGraw-Hill, NY.

Special Symbols

(Alphabetically listed by first symbol letter.)

Symbol.....	Name, if any	Page no.
$\mathcal{A}$		50
BHF	Basic Hyperfinite	
	Subsets of ${}^*\mathbb{N}$	30
$\mathcal{C}({}^*\mathcal{H})$	Internal Constants.....	21
$\mathcal{C}(\mathcal{H})$	Standard Constants	21
$\mathcal{C}_{PSR}$		37
$\mathcal{C}_{SR}$		38
$\mathcal{C}'_{PSR}$		123
$\mathcal{C}_J$	A Set Of Jordan	
	Measurable subsets	125
$D \int_R f(\vec{x}) dX$	Darboux Integral	125
$\text{ext}(Q)$	Exterior Subrectangles.....	63, 132
$F(B)$	Set Of All Finite	
	Subsets of B	31
$\mathcal{H}$	The Superstructure.....	17
${}^*\mathcal{H}$	Hyperstructure	21
$\text{int}(A)$	Interior Points in A	50
$\text{inn}(Q)$	Inner Subrectangles.....	63, 132
J	Jordan-measurable set	42
$L[S]$	m-dimensional	
	Element	47
$L(f)$		120
$L(f, P)$	Lower Sum	120
$L(\mathbb{R}^n, \mathcal{B})$	Set Of L -gauges	141
$\mathcal{L}(\lambda)$	δ -fine Partitions	142
M_1	Infinitesimals	11
$\mu'(\vec{p})$	Deleted Monad	77
$\mu(0)$	Infinitesimals	11
$\mu(r)$	Monad about r	12
$\mu(\vec{v}) = \mu^n(\vec{v})$	Monad about $\vec{v}$	13, 88
$\mu(0)^+$	Nonnegative	
	Infinitesimals	69
$(\text{mod } o(\cdot))$		69
$m(x_1\delta_1, \dots, x_p\delta_p)$	Magnification Operator.....	72
$\hat{\delta}$	Microguage.....	75
$\mathbb{N}$	Natural Numbers	10
$\mathbb{N}_\infty$	Infinite Natural No.s.....	19
o	Infinitesimals	11
$\mathcal{O}$	Limited numbers	11
$\mathcal{O}^n$	Limited Vectors	13
$o(\epsilon)$	Order Ideal.....	68

$o^n(\epsilon_1, \dots, \epsilon_n)$	71
$\mathcal{P}_\Gamma$	Internal Hyper- polygonal Curve 33
$\mathcal{P}(W)$	Power Set 17
$P_i(R)$	i'th Projection 18
P_i	Hyperfinite Partition of $[a_i, b_i]$ 35
$P = P_1 \times \dots \times P_n$	Simple Fine Partition 36
P_δ	Gauge Partition 75
(Q_δ, P_δ)	δ -fine partition 75
$(Q_{\hat{\delta}}, P_{\hat{\delta}})$	Micropartition 75
Q	Intermediate Partition 120
$\mathcal{Q}$	Intermediate Partition Map 122
$^*\mathbb{R}$	Extended reals 10
$^*\mathbb{R}$	Hyperreals 10
$^*\mathbb{R}$	Star-reals 10
$\mathbb{R}^+$	Positive Reals 10
$\mathbb{R}$	Real Numbers 10
$^*\mathbb{R} - \mathcal{O}$	Infinite Hyperreals 11
$^*\mathbb{R}_\infty$	Infinite Hyperreals 11
$\mathbb{R}^n$	Euclidean n-space 12
$\mathbb{R}$	19
$\mathbb{R}_\infty^+$	Positive Infinite Hyperreals 26
R	Rectangle 35
R_q	Subrectangle 35
R_S	Subrectangle 37
$\text{st}()$	Standard Part 14
$\mathcal{S}(P)$	37
S	Subrectangle 48
$S(f, P, Q)$	Riemann Sum 122
$\mathcal{U}$	Basic Universe 17
$U(f)$	120
$U(f, P)$	Upper Sum 120
$v(R_q)$	Volume 85
X_{n+1}	17
$Z \int_R f(\vec{x}) * \ P\ $	Z-integral 121

(Non-alphabetical listing.)

$\approx$	Infinitely Close 11
$\ \cdot\ $	Euclidean Norm 11
∂A	Boundard Points of A 55

$\partial(Q)$	Boundary Subrectangles	
		62
$\ P\ $		120
$\ S\ $		120

**Some Applications of Nonstandard Analysis
to Advanced Undergraduate Mathematics**

◇ Very Elementary Physics◇

Robert A. Herrmann

A 1991 Instructional Development Project from the
Mathematics Department
United States Naval Academy
572C Holloway Road
Annapolis, Maryland 21402-5002

IMPORTANT NOTICE

Since the writing of this book was financed entirely by a designated grant from the Federal Government that was specifically obtained for this sole purpose then a copyright for this specific book cannot be obtained by its author. Any portion of its contents can be copied and used without seeking permissions from the author. However, when such copying or use is made of this material, it is necessary that the author and the U. S. Naval Academy be indicated as the source of the material being used. Further note that certain new results that appear in this book will be published under the author's name in scholarly journals.

CONTENTS

Chapter 1

Introduction

1.1	Brief Comments	165
1.2	Manual Structure	165

Chapter 2

Mechanics

2.1	Instantaneous Velocity	167
2.2	Acceleration	171
2.3	Forces and Newton's Law	172
2.4	Vectors	174
2.5	Energy and Force Fields	175
2.6	General Impulse	176

Chapter 3

Slightly Less Basic Mechanics

3.1	Mass	181
3.2	Moments and Center of Mass	183
3.3	Point Masses	185
3.4	Standard Rules	187
	References	189
	Additional Special symbols	190

To be continued by members of the physics community.

INTRODUCTION

1.1 Brief Comments

Since the time of Archimedes the major applications of infinitesimal reasoning have been in the general discipline of geometry and what we now categorize as the subject matter of Physics. All of the applications that appear in the first volume in this series – Infinitesimal Modeling – are taken from these two disciplines. The methods employed within this manual are based exclusively upon those that appear in the our major reference the *Some Applications of Nonstandard Analysis to Undergraduate Mathematics – Infinitesimal Modeling*. From time-to-time, a portion of certain applications, discussions and conclusions are directly retrieved from the Infinitesimal Modeling manual so that this Elementary Physics manual will present, from the view point of applications, a continuous and cohesive structure that parallels the standard first undergraduate physics course that requires the Calculus as a prerequisite. **One important feature of this physics manual is that many of the basic rigorous derivations are followed immediately by additional derivations that have been translated into the classical language used in most undergraduate calculus courses. This will enable most undergraduate students to more easily comprehend a derivation's logical sequence.**

1.2 Manual Structure

We will not replicate an actual physics course in this manual but rather present representative derivations using the rules established within the Infinitesimal Modeling manual for some of the more significant integral and differential equation models for the behavior of will-known natural systems.

This presentation will only be for mechanics. Obviously, we can only make a minute sampling from these very broad categories. However, it is hoped that, if care is exercised, the examples chosen will lead the instructor to seek more rigorous derivations for the more complex and refined aspects of system behavior. The individual physics instructor is certainly more intuitively and academically prepared than you author for a penetrating and rigorous investigation of the more subtle aspects of this subject.

It is the belief of your author that the most expedient approach is to train the scientific community in the rudiments of rigorous infinitesimal analysis by such devices as the Infinitesimal Modeling manual, the new infinitesimal calculus courses that have been introduced throughout the world, and manuals similar to this Physical Manual. Once individual scientists achieve a working knowledge of the basic principles then those who specialize in a given subject area are the appropriate ones to continue a more in depth exploration. What is discovered by an in depth rigorous infinitesimal approach is that simple fundamental observations lead to simple standard expressions. These expressions, after being transferred to the nonstandard model, yield a simple view of a new world called the nonstandard physical world (NSP-world). These transferred simple processes also lead to NSP-world processes that when applied within the NSP-world lead back again to standard integral or differential equation models that mirror natural system behavior. Usually, one acquires knowledge about the appropriate NSP-world processes through observation of simple or idealized natural system behavior and then accepts those NSP-world views that lead to verified predictions. It is by means of this back-and-forth approach that we gain useful knowledge about the NSP-world.

The following notation indicates the beginning and ending of each derivation. For the rigorous analytical derivation using the language of The Basic Manual, the beginning and ending are marked by a $\diamond$ symbol. A second or third derivation is also be denoted by $\diamond$; but, each is included within a subsection marked by $\{Second\ Derivation.....\}$ or $\{Third\ Derivation.....\}$

Chapter 2.

MECHANICS

2.1. Instantaneous Velocity.

In 1686, Newton [Newton [1934]] gives what he claims is the easily comprehended notion of the “ultimate velocity,” or what we now term the **instantaneous velocity**, for an actual real material object. *But by the same argument it may be alleged that a body arriving at a certain place, and there stopping, has no ultimate velocity; because the velocity, before the body comes to the place, is not its ultimate velocity; when it has arrived, there is none. But the answer is easy; for by the ultimate velocity is meant that with which the body is moved, neither before it arrives at its last place and the motion ceases, nor after, but at the very instant it arrives; that is, the velocity with which the body arrives at its last place, and with which the motion ceases.* [Scholium to Lemma XI in Book 1] For the case of nonzero instantaneous velocity, one might gather from this the power of Newton’s mental vision and his intuitive comprehension of future behavior. Since even though the object may not appear to move at the “instant” (i.e. an instant of time) one observes the hands of clock point at a numerical representation for the time, the object did arrive at a space location and has the capacity to change its position. It is claimed, incorrectly, that this type of change in position is noted when a second observation is made and the hands of the same clock are assumed to point at a different numerical representation for the time.

Newton’s modeling of this idea is firmly rooted in his concept of the relation between geometry (the basic mathematical structure of the 1600’s) and its relation to mechanics. *Geometry does not teach us to draw lines, but requires them to be drawn, for it requires that the learner should first be taught to describe these accurately before he enters geometry, then it shows how by these operations problems may be solved. To describe right lines and circles are problems, but not geometrical problems. The solution of these problems is required from mechanics,...therefore geometry is founded in mechanical practice, and is nothing but that part of universal mechanics which accurately proposes and demonstrates the art of measure.* [Newton [1934:xvii]] Newton’s claim is that our observations and intuitive comprehension of mechanics comes first in our education. These concepts are then abstracted to include the vague notion that objects have certain “capacities or potentials to do things”- the capacity or potential idea. We are told that it is after experimentation, observation and reflection that the mathematical structure is evoked and these “easy” capacity concepts are modeled.

The abstract notion of instantaneous velocity may have been “easy” for Newton to grasp, but it was incomprehensible to Berkeley and many others who believed that such abstractions could not be applied to actual real material objects. The paramount philosophy of science for Berkeley was a science of the material and directly observed universe. Any arguments that relied upon such abstractions would need to be rejected.

How can we communicate such an abstract idea to students who do not necessarily possess Newton’s obvious mental ability? Tipler [1982:26] writes: *At first glance, it might seem impossible to define the velocity of a particle at a single instant, i.e., at a specific time. At a time t_1 , the particle is at a single point x_1 . If it is at a single point, how can it be moving? On the other hand, if it is not moving, shouldn’t it stay at the same point? This is an age-old paradox, which can be resolved when we realize that to observe motion and thus define it, we must look at the position of the object at more than one time. It is then possible to define the velocity at an instant by a limiting process.* Unfortunately, Tipler has reversed Newton’s original notion, that something exists prior to the

modeling of motion and this something is the capacity to move. This capacity then leads to the need to seek various observations from which a numerical value can be “defined.” Is it now possible to derive the well-known derivative expression for instantaneous velocity and, at least partially, retain Newton’s capacity concept by infinitesimally modeling what is indeed easily observed behavior? An affirmative answer to this question depends upon your acceptance fundamental properties the Galilean theory of uniform velocity, infinitesimal analysis and its associated interpretations.

◇ Assuming that we are working in the laboratory setting with a fixed standard for the measure of (linear) distance and time, let increasing $d: I \rightarrow \mathbb{R}$, where $I = [a, b]$, $a \neq b$, $a, b \in \mathbb{R}$, represent the distance an object travels over the time interval $[a, b]$. Of course, a lot has been assumed, even that it makes sense to consider time as representable by a continuum such as $[a, b]$. [Remark. Recent work [Herrmann [1989]] has shown that if time is not a continuum then there are internal functions that relate time to a continuum and these functions are infinitely close to any discrete (discontinuous) time concept.] Suppose that we extend our “observations” of finitely many cases and accept for a very simple motion that $d(t_2) - d(t_1)/(t_2 - t_1) = v = \|\vec{v}\|$ a constant (the scalar velocity), for any $t_1, t_2 \in [a, b]$, $t_1 < t_2$. If such a motion persisted, then, of course, v can be used to calculate a change in the distance over a change in time. Let $t_1 \in (a, b)$ and Δt any positive real number such that $t_1 + \Delta t \in (a, b]$. Then, assuming a constant scalar velocity, one has that $d(t_1 + \Delta t) = d(t_1) + v \Delta t$ or that

$$d(t_1 + \Delta t) - d(t_1) = v \Delta t. \quad (1)$$

What if the distance expression was more complex than the linear type expressed by (1)? We are seeking an appropriate definition for v that extends this case. First, it follows immediately that v would be a function in $t \in [a, b]$. Since a constant function is the simplest in the collection of continuous functions, then, at this stage of our analysis, we simply require v to be continuous on $[a, b]$. However, we know from the Extreme Value Theorem that for any $[t_1, t_2] \subset [a, b]$, $t_2 = t_1 + \Delta t$, $\Delta t > 0$, there exists $t_m, t_M \in [t_1, t_2]$ such that for each $t \in [t_1, t_2]$, $v(t_m) \leq v(t) \leq v(t_M)$. Hence,

$$(t_2 - t_1)v(t_m) \leq (t_2 - t_1)v(t) \leq (t_2 - t_1)v(t_M), \quad (2)$$

for each $t \in [t_1, t_2]$.

Now consider the physical processes involved and correspond equation (2) to these processes. For the time span $t_2 - t_1$, it appears reasonable to state, using the case where t_m and t_M are constant and our intuitive notion of distance traveled, that the actual distance moved $d(t_2) - d(t_1)$ has the property that

$$(t_2 - t_1)v(t_m) \leq d(t_2) - d(t_1) \leq (t_2 - t_1)v(t_M), \quad (3)$$

If you accept the model for distance expressed by (3), then from the Intermediate Value Theorem there would necessarily exist some $t' \in [t_1, t_2]$ such that $d(t_2) - d(t_1) = v(t')(t_2 - t_1)$. What this means is that the distance can be calculated, knowing $v(t')$, as if it were created by a constant scalar velocity. Transfer the above results by intuitive *-transfer to the infinitesimal NSP-world (i.e. they “hold” true for the infinitesimals). Thus, for any positive $\epsilon \in \mu(0)$ and any $t_1 \in (a, b)$, it follows that there exists some $t' \in [t_1, t_1 + \epsilon]$ such that

$$(*d(t_1 + \epsilon) - d(t_1))/\epsilon = {}^*v(t'). \quad (4)$$

The same argument shows that if negative $\epsilon \in \mu(0)$, then there exists some $t' \in [t_1 + \epsilon, t_1]$ such that (4) as well.

Unfortunately, we do not know the value of t' in (4). But, once again, continuity of v at t_1 does allow us to write that $*v(t') \approx v(t_1)$. Hence,

$$(*d(t_1 + \epsilon) - d(t_1))/\epsilon \approx v(t_1). \quad (5)$$

Obviously, since ϵ is an arbitrary nonzero infinitesimal then if there exists a distance function, d , that satisfies (3) for all such time intervals, then d must be differentiable at t_1 and application of the standard part operator implies that $d'(t_1) = v(t_1)$. $\diamond$

{*Second derivation.* In what follows, the above derivation for the instantaneous velocity function is reworded into a quasi-classical description using slightly modified calculus terminology. The ground rules for this second derivation are:

(i) As is done in Internal Set Theory, the “*” notation is removed from the functions since whether they are nonstandard extensions of standard functions is clear from the function’s argument (i.e. preimage).

(ii) The symbols “ $\approx$ ” is translated by the term “infinitely close.” This relation can be physically characterized by stating that no standard machine can measure any difference between the quantity on the left and the quantity on the right no matter how small the machine error.

(iii) Except for $\mathbb{N}_\infty$, hyperreal numbers are usually limited. Hence, simply call such a hyperreal number by the single word term “number.”

(iv) We use the fact that functions defined and continuous on $[a, b]$ preserve the infinitely close concept for these numbers. That is if $t, t_1 \in [a, b]$ and $t \approx t_1$, then $f(t) \approx f(t_1)$. Infinitesimals may be called the “infinitely or very small.” These numbers can be physically characterized as measures that are smaller than any standard machine error — measures that appear to a machine to be zero.

(v) Rather than use the standard part operator, where applicable, use the simple term “limit” in its place, since it has the same operative properties. Also use the fact that limits of two infinitely close numbers are equal.

The modified classical derivation is exactly the same until after equation (3). Then it continues as follows:

$\diamond$ If you accept the model for distance expressed by (3), then from the Intermediate Value Theorem there would necessarily exist some $t' \in [t_1, t_2]$ such that $d(t_2) - d(t_1) = v(t')(t_2 - t_1)$. What this means is that the distance can be calculated, knowing $v(t')$, as if it was created by a constant scalar velocity. Now (3) and these facts hold for the infinitely small. Thus, for any positive infinitely small Δt and any $t_1 \in (a, b)$, it follows that there exists some t' such that $t_1 \leq t' \leq t_1 + \Delta t$ (i.e., $t' \in [t_1, t_1 + \Delta t]$ and

$$(d(t_1 + \Delta t) - d(t_1))/\Delta t = v(t'). \quad (4)$$

The same argument shows that if Δt is a negative infinitely small number, then there is some number t' such that $t_1 + \Delta t \leq t' \leq t_1$ and once again (4) holds.

Unfortunately, we do not know the value of t' in (4). But, since $t' \approx t_1$ then continuity of v at t_1 allows us to write that $v(t') \approx v(t_1)$. Hence,

$$(d(t_1 + \Delta t) - d(t_1))/\Delta t \approx v(t_1). \quad (5)$$

Obviously, since nonzero Δt is an arbitrary and infinitely small, then if there exists a distance function, d , that satisfies (3) for all such time intervals, then d must be differentiable at t_1 and the limit of the left hand side of (5), [as Δt varies could be added, but is not necessary] must equal the limit of the right hand side which is the constant $v(t_1)$. This implies that $d'(t_1) = v(t_1)$. $\diamond$

{*Third derivation — entirely classical.* As discussed in the Infinitesimal Modeling manual, many derivations for differential equation models require an infinitesimalizing process for behavior that is only approximated within the standard world for "small" quantities. One of the simplest illustrations of this is the differential equation model for Newton's Law of Cooling. The necessity for this special process comes from the experiential evidence that the observed behavior holds only for small measures of the independent variables and as the measures are reduced such behavior is more closely approximated by a standard functional expression. This is in direct contrast to an expression such as (3) and the discussion that follows where the results appear to hold for all intervals $[t_1, t_2]$. In the case of such concepts as the instantaneous velocity, it is possible to present an entirely classical derivation. This classical derivation begins with equation (3) and continues as follows:

◇ If you accept the model for distance expressed by (3), then from the Intermediate Value Theorem there would necessarily exist some $t' \in [t_1, t_2]$ such that $d(t_2) - d(t_1) = v(t')(t_2 - t_1)$. What this means is that the distance can be calculated, knowing $v(t')$, as if it were created by a constant scalar velocity. Thus for $t_1 \in (a, b)$ and for every positive Δt such $t_2 = t_1 + \Delta t \in (a, b)$ there exists some t' such that

$$(d(t_1 + \Delta t) - d(t_1))/\Delta t = v(t'), \quad (4)$$

and $t_1 \leq t' \leq t_1 + \Delta t$.

Using the Axiom of Choice, we can consider a function f defined on the respective Δt such that $f(\Delta t) = t'$. Repeating the process for the negative Δt such that $[t_1 + \Delta t, t_1] \subset (a, b)$ and extending the function f to include these negative Δt leads to the conclusion that $\lim_{\Delta t \rightarrow 0} f(\Delta t) = t_1$. From the assumed continuity of v it follows that

$$\lim_{\Delta t \rightarrow 0} \left(\frac{d(t_1 + \Delta t) - d(t_1)}{\Delta t} \right) = \lim_{\Delta t \rightarrow 0} v(f(\Delta t)) = v(t_1). \quad (5)$$

This all implies that under the conditions stated $d'(t_1) = v(t_1)$. ◇}

How the above derivations improve our comprehension of the concept of instantaneous velocity is discussed at the conclusion of this section. Returning to the concept of the capacity to move, Theorem 9.1.2 of the Infinitesimal Modeling manual indicates that for each $\epsilon \in \mu(0)$

$$\Delta^* d_{t_1}(\epsilon) = {}^*d(t_1 + \epsilon) - d(t_1) = v(t_1)\epsilon + \epsilon\lambda(\epsilon) = {}^*T(\epsilon) + \epsilon\lambda(\epsilon), \quad (6)$$

where λ is a local function defined by (6) and linear $T: \mathbb{R} \rightarrow \mathbb{R}$. Thus for each $\epsilon \in \mu(0)$, $\Delta^* d_{t_1}(\epsilon)$ and $v(t_1)\epsilon$ are not just infinitely close, but they are infinitely close of the first order. (See the Infinitesimal Modeling manual Section 8.3.)

Definition 2.1. (Infinitely Close of Order One). Two hyperreal valued functions defined on $\mu(0)$ are said to be **Infinitely Close of Order One or of the First Order** if for each $\epsilon \in \mu(0)$ there exists some $t \in \mu(0)$ such that

$$f(\epsilon) = g(\epsilon) + \epsilon t.$$

In which case, this is denoted by $f \sim_1 g$. Further, if the two functions are considered to be measuring some physical properties, then we often say that the natural world effects of these properties are **indistinguishable (at level one or on the first level)**.

It is a simple matter to show that $\sim_1$ is an equivalence relation on the set of all hyperreal valued functions defined on $\mu(0)$. The capacity to move concept is now represented in the monadic

environment by noting that

$$\Delta *d_{t_1} \sim_1 *T = d'(t_1)(\cdot). \quad (7)$$

Or in words, *within the monadic world the distance represented by $\Delta *d_{t_1}$ is indistinguishable from (has the same effect as) that produced by a scalar velocity $d'(t_1)$* . Of course, the function d' is termed the **instantaneous velocity**. As Newton claimed, if mechanics leads to geometry, then this is what motivated the geometric concepts of the rectifiable curve, tangents, curvature and the like that appear in the Infinitesimal Modeling manual Chapter 7, section 7.2, Chapter 8, sections 8.5 – 8.6 and Chapter 9, sections 9.5 – 9.7.

Within the foundations of any discipline it is often difficult to refine even slightly what may have been assumed previously to be an elementary and not dissectible assertion. Thus, until now, this has been with the idea of instantaneous velocity. However,

- (i) let the distance function, $d(t)$, and an unknown continuous scalar velocity function, $v(t) = \|\vec{v}(t)\|$, be related by expression (3).
- (ii) Let (3) hold for every time subinterval $[c, d] = [t_1 + (\Delta t)_1, t_1] \subset (a, b)$ and $[c, d] = [t_1, t_1 + (\Delta t)_2]$, where $(\Delta t)_1 < 0$, $(\Delta t)_2 > 0$.
- (iii) There exists some $t' \in [c, d]$ such that the actual distance traveled $d(t_2) - d(t_1) = v(t')(t_2 - t_1)$.

Then d must be differentiable at t_1 and the only standard scalar velocity function that satisfies (i), (ii) and (iii) is the function $v(t_1) = d'(t_1)$. The instantaneous velocity is not obtained by simply postulating a definition but is derived from more elementary observations.

2.2. Acceleration

For twenty years, Galileo struggled with the problem of representing the velocity of a falling body in terms of distance [Gillispie [1960:42]]. After failing in every attempt, a new idea began to ferment - an idea that today seems so common place since, as illustrated in the previous section, we are taught to think of elementary velocity as expressed in a time coordinate. But, it was Galileo's revolutionary concept of time as an independent abstract entity that led to the theory of motion that Newton applied in his dynamical geometry and Mathematical Principles of Natural Philosophy [Newton [1686]]. The remarkable insight exemplified by Galileo should not be underestimated. His struggle once again indicates the difficulty that scientists often face when, through reflection, they alter some well entrenched but erroneous elementary precept.

◇ As seen in the previous section, the only elementary standard function the preserves our intuitive understanding about the elementary measure of distances traveled is the instantaneous velocity. Further, in the monadic environment, the Galilean properties for constant or uniform scalar velocities and how they are compounded to yield distances traveled are indistinguishable from the actual quantities. But, now that we have accepted the instantaneous velocity, $v = \|\vec{v}\|$, as the appropriate elementary nonconstant velocity concept, we can certainly apply a section 1 type analysis to v . This requires the strict application of the Galilean theory of motion for a constant acceleration on one hand, and Newton's concept of the nonconstant acceleration (produced by a force) that leads to nonuniform velocity on the other hand.

Thus for time interval $[a, b]$ substitute in expression (3) of section 1, v for d and a representation for a continuous scalar acceleration $a = \|\vec{a}\|$ for v and obtain

$$(t_2 - t_1)a(t_m) \leq v(t_2) - v(t_1) \leq (t_2 - t_1)a(t_M). \quad (1)$$

The expression on the left of (1) applies the concept of a constant acceleration and measures the minimum possible [linear] uniform change in the velocity and that on the right the maximum possible uniform change. This implies, as in the case of instantaneous velocity and under parallel hypotheses as stated in section 1 (i), (ii), that there exists a unique scalar acceleration function a such that $a(t_1) = v'(t_1) = d^{(2)}(t_1)$ - the **instantaneous acceleration**. $\diamond$ As with the case of the instantaneous velocity we also have

$$\Delta^* v_{t_1} \sim_1 {}^*T_a = v'(t_1)(\cdot), \quad (2)$$

which indicates that within the monadic environment that the change in velocity in the N-world is indistinguishable on the first level from that produced by a constant acceleration. However, application of Theorem 8.4.2 yields that an associated change in distance satisfies the two, difficult to visualize, statements

$$\begin{aligned} \Delta^2 {}^*d_{t_1} \sim_1 {}^*T_a = v'(t_1)(\cdot) = d^{(2)}(\cdot), \text{ and} \\ \Delta^2 {}^*d_{t_1} \sim_2 {}^*T_a = {}^*d^{(2)}(\cdot), \end{aligned} \quad (3)$$

where $\sim_2$ is defined in the obvious manner by replacing ϵ in the last term in the equation in definition 2.1 by ϵ^2 and the second equation in (3) is the best that we can state, in generally.

2.3. Forces and Newton's "Law"

The story is told that a student asked Max Planck to explain how he perceived nuclear forces? Planck is said to have replied that he would perceive them as someone pulling on his coat sleeve. Whether or not this story is factual, experience indicates that a change in velocity is better comprehended by considering such a change as the effect of a more easily sensed "force" that may be the cause of the change. This brings use the Newton's so-called Second Law of Motion.

It is not true that Newton formulated his Second Law as it is taught in our elementary physics courses where we are told that the scalar force is proportional to the instantaneous acceleration. It is also not true that he formulated his Second Law in terms of something equivalent to the derivative of the momentum. First, Newton defined the "quantity of motion" as follows: *The quantity of motion is the measure of the same, arising from the [scalar] velocity and the quantity of matter conjointly* [Newton [1934:1]] Thus the quantity of motion is the momentum. As indicated by the explanation that follows his statement, his Second Law was the observation that *The [uniform] change of [the quantity of] motion is proportional to the [constant] motive force impressed; and is made in the direction of the right [i.e., straight] line in which the force is impressed.* [[Newton [1934: 13]] He proceeds in the Scholium to that section to apply this Second Law and his idea that the total effect of finitely many constant forces is additive over time to establish Galileo's discovery that the *...descent of bodies varies as the square of the time.* [Newton [1934:21]] In terms of a constant impulse notion, Newton's argument does not include the mass, but rather leaves the mass as the constant of proportionality. In summation notation, the argument has the following form. The time $[a, b]$ is subdivided into n equal time intervals Δt . The scalar force during these time intervals is a constant F . Then the force times the length of time impressed (i.e. the **impulse**) is proportional to the uniform change in scalar velocity, Δv , and all such uniform changes in the scalar velocity are equal in value. Hence, the composition of such forces would yield a total effect

$$\sum F(\Delta t) = F \sum \Delta t = F(b - a) \propto \sum \Delta v = v(b) - v(a) = v, \quad (1)$$

where F now becomes the *whole force*, the constant of proportionality is the same for each summand and from the additivity of velocities in a straight line motion v is the *whole velocity*. Now translating

into our symbols, Newton writes $F(b - a) \propto v$ as $(b - a) \propto v$. He then states that the *spaces* [i.e., distance traveled] in *proportional times are as the product of the velocities and times*;.... [Newton [1934:21]] Thus such a distance $d \propto v(b - a) \propto (b - a)^2$.

In his Principles, Newton does not utilize his method of fluxions, even though in other communications he does, to develop his theory of motions of material bodies. He first presents arguments delineating the “ultimate ratios” between geometric measures - arguments that employ those intuitive concepts acceptable and apparently comprehensible by geometers, expressions such as *magnitude diminished in infinitum*. He then correlates time to the measure of one leg of a right triangle and velocity to the other leg. Considering his geometric notion of the ultimate ratio of the areas of these triangles as the length of the sides *diminish in infinitum*, which he previously established was as to the squares of the *homologous sides*, he draws the conclusion that *the spaces which a body by an finite force urging it, whether that force is determined and immutable, or is continually augmented or continually diminished* [with respect to time], *are in the very beginning of the motion to each other as the squares of the times*. [Newton [1934:34]] In corollary *iv* he writes: *And therefore the forces are directly as the spaces described in the very beginning of the motion, and inversely as the squares of the times*. [Newton [1934:35]] The expression *very beginning* is used to describe the ultimate ratio concept and that this is a point force associated with an instant of time. Further notice that he has replaced the idea of constant forces over a time subinterval with forces that are being altered *continually*.

The logical vagueness of Newton’s arguments can be eliminated by modern infinitesimal analysis. $\diamond$ Consider the time interval $[a, b]$ and any $[t_1, t_2] \subset [a, b]$. Assume that the scalar force F is a continuous real valued function defined on $[a, b]$. We look at Newton’s observations relative to constant scalar forces and there relation to uniform changes in the scalar velocity. First, from continuity, there exists $t_m, t_M \in [t_1, t_2]$ such that

$$F(t_m)(t_2 - t_1) \leq F(t)(t_2 - t_1) \leq F(t_M)(t_2 - t_1) \quad (2)$$

for each $t \in [t_1, t_2]$. Let C be a constant of proportionality which, for this particular simplistic problem, is considered absolutely constant in character. Let q denote a real valued function that measures, with respect to time, the notion of the quantity of motion. Then from the actual stated Second Law it follows that the minimal possible change in momentum over the $[t_1, t_2]$ is $q(t_m) = F(t_m)(t_2 - t_1)$ and the maximal change is $q(t_M) = F(t_M)(t_2 - t_1)$. Now the actual velocity function in terms of time is, at present, unknown. But, whatever it may be, we consider the actual quantity of motion q to be an extension of the constant case and, hence, it is proportional, for a given object, to the actual change in velocity or to $C(v(t_2) - v(t_1))$. Assuming that such a change lies somewhere between the minimal and maximal changes then applying the continuity of the force function one obtains that there exists some $t' \in [t_1, t_2]$ such that

$$F(t')(t_2 - t_1) = C(v(t_2) - v(t_1)). \quad (3)$$

Letting $t_1 \in (a, b)$ and noticing that the above is assumed to hold for *any* subinterval $[t_1, t_1 + \Delta t]$, $\Delta t > 0$ of (a, b) or *any* subinterval $[t_1 + \Delta t, t_1]$, $\Delta t < 0$ of $[a, b]$ then *-transfer yields that for nonzero $\epsilon \in \mu(0)$ there exists some $t' \approx t_1$ such that

$$*F(t') = \frac{C(*v(t_1 + \epsilon) - v(t_1))}{\epsilon} \approx F(t_1). \quad (4)$$

Taking the standard part operator we arrive at a derivation of our modern Newton's Second Law of Motion. That for all of this to occur there must exist a velocity function that is differentiable at t_1 and the only relation between such a point force function and such a velocity function is

$$F(t_1) = Cv'(t_1) = Ca(t_1), \quad (5)$$

where a is the **instantaneous acceleration**. $\diamond$ As previously, this can be further related to the change in momentum for an infinitesimal time by the expression

$$C(\Delta * v) \sim_1 F(t_1)(\cdot). \quad (6)$$

Or, as stated in words, that the change in momentum over an infinitesimal time is indistinguishable on the first level from that produced by a constant force applied to infinitesimal time periods. Moreover, the easily grasped concept of the impulse was the starting point in Newton's original arguments. Unfortunately, some modern textbooks do not introduce the impulse relative to constant forces as a first principle.

{*Second derivation.* Start the modification of the above after equation (3).

$\diamond$ Letting $t_1 \in (a, b)$ and noticing that the above is assumed to hold for *any* subinterval $[t_1, t_2]$ of $[a, b]$ then all of the above holds for infinitely small subintervals. Thus for nonzero infinitely small Δt there exists some $t' \approx t_1$ such that

$$F(t') = \frac{C(v(t_1 + \Delta t) - v(t_1))}{\Delta t} \approx F(t_1). \quad (4)$$

Taking the limit of the left hand side [as Δt varies] we arrive at a derivation of our modern Newton's Second Law of Motion.

$$C \lim \frac{(v(t_1 + \Delta t) - v(t_1))}{\Delta t} = F(t_1). \quad (5)$$

Thus for all of this to occur there must exist a velocity function that is differentiable at t_1 and the only relation between such a point force function and such a velocity function is

$$F(t_1) = Cv'(t_1) = Ca(t_1), \quad (6)$$

where a is the **instantaneous acceleration**. $\diamond$

The reader may replicate the third derivation along with the discussion of properties (i), (ii) and (iii) as they appear in Section 2.1 for the instantaneous acceleration and Newton's Second Law.

Notice that one derivation method – the maximum and minimum method – yields the instantaneous velocity, instantaneous acceleration and the modern Second Law of Motion from what may be more fundamental observations.

2.4. Vectors

For constant forces, Newton's Corollary I to his three laws establishes for constant forces the idea that bodies move in $\mathbb{R}^2$ by the vector composition of two scalar forces acting simultaneously upon a particle or fixed point in a material body. [Newton [1934:14]] Newtonian mechanics may appear to begin with the idea that the position of a particle in an n -dimensional coordinate system

is dependent upon the composition of n forces (the cause); but, in actually, it is the position (i.e. the effect) that is the fundamental concept. The position of a particle is defined in terms of n coordinate functions, each expressed in the independent parameter — time. This leads to the **position vector (i.e. radius vector)** $\vec{r} = (x_1(t), \dots, x_n(t))$, $t \in [a, b]$. Applying the previous section to these coordinate functions independently, leads to the vector form for the instantaneous velocity, instantaneous acceleration, and force.

In the Infinitesimal Modeling manual, the geometric concept of the length of a continuous curve is fully discussed and, where possible, this length is correlated to the integral by means of our integral rules. Following Newton's notion of a dynamic geometry, the same conclusions evolve from the instantaneous velocity. Suppose that a point force $\vec{F}(t)$, $t \in [a, b]$, is the cause that induces an instantaneous acceleration $\vec{a}(t)$ upon a particle. Since $\vec{F}$ was assumed continuous on some time interval $[a, b]$ then $\vec{v}'$ is continuous on $[a, b]$. Consequently, $\vec{v}$ and $\|\vec{v}\|$ are continuous on $[a, b]$. Assuming we are in $\mathbb{R}^n$ then, noting that for $t \in [a, b]$, $\|v(t)\| = \sqrt{(x'_1(t))^2 + \dots + (x'_n(t))^2} = d'(t)/dt$, we obtain

$$d(b) - d(a) = \int_a^b \sqrt{(x'_1(t))^2 + \dots + (x'_n(t))^2} dt, \quad (1)$$

which is the same result obtained in the Infinitesimal Modeling manual by considering hyperpolygonal representations for the length of a continuously differentiable geometric curve.

Let the path of motion be represented by continuous $c: [a, b] \rightarrow \mathbb{R}^n$, where $c(t) = (x_1(t), \dots, x_n(t))$. By Theorems 9.5.1 and 9.5.2 of Chapter 9 and Theorem N.1 at the end of Chapter 10 in the Infinitesimal Modeling manual it is shown that if $\vec{v}(t) = \vec{c}'(t) \neq \vec{0}$, $t \in (a, b)$, then the unit tangent $\vec{T} = \pm \vec{c}'(t)/\|\vec{c}'(t)\|$ and that $\vec{T}$ is almost parallel to every hyperpolygonal directed line segment $(^*c(t+dx) - c(t))/\|{}^*c(t+dx) - c(t)\|$ for each nonzero $dx \in \mu(0)$. This implies that the unit instantaneous velocity vector $\vec{v}(t)/\|\vec{v}(t)\|$ is not only one of the two possible unit tangent vectors but also has the property of being almost parallel to each $(^*c(t+dx) - c(t))/\|{}^*c(t+dx) - c(t)\|$. Since the effect of vectors as a model for natural world (i.e. N-world) behavior is often related to the physical concept of “direction” then as far as the N-world effects are concerned the direction of the standard velocity vector attached to the point $c(t)$ is indistinguishable from the direction of $\pm(^*c(t+dx) - c(t))$, $0 \neq dx \in \mu(0)$. The indistinguishableness of such effects is beneficial when energy is to be considered.

2.5. Energy and Force Fields

In the Infinitesimal Modeling manual, the line integral is employed to measure the *energy expended within a forced field while moving along a curve* (Application 8.2.1) and the customary method of line integral evaluation obtained. We examine here the first portion of this derivation.

◇ In experimental physics, the concept of “work” (energy expended) is introduced. **All one needs to do is to establish its properties for a polygonal curve.** Suppose we have continuous force field $F: E \rightarrow \mathbb{R}^n$. Let $\mathcal{P}_k \subset \mathbb{R}^n$ be any finite polygonal curve, ℓ_j one of the line segment portions of $\mathcal{P}_k$ with $\vec{v}_j$ denoting this line segment considered as a directed line segment in the direction of motion through the field F . If F is constant on ℓ_j , then the work done moving along ℓ_j is defined as $W(\ell_j) = F \bullet (\vec{v}_j/\|\vec{v}_j\|)\|\vec{v}_j\|$, where length of $\ell_j = \|\vec{v}_j\|$. What if the force field is not constant? Consider $\mathcal{P}_k$ as represented by a continuous $\ell: [a, b] \rightarrow \mathbb{R}^n$ and assume that F is defined on ℓ . Then for a given $\ell_j = \{(x_1(t), \dots, x_n(t)) \mid t \in [t_{j-1}, t_j]\}$ there exists some t_m, t_M such that $W_m(\ell_j) = F(\ell(t_m)) \bullet \vec{v}_j \leq W(\ell_j) = F(\ell(t)) \bullet \vec{v}_j = W_M(\ell_j) = F(\ell(t_M)) \bullet \vec{v}_j$ for each $t \in [t_{j-1}, t_j]$. Let's make the one assumption that the actual amount of work expended moving along the line

segment ℓ_j is $W(\ell_j)$ and that $W_m(\ell_j) \leq W(\ell_j) \leq W_M(\ell_j)$. Then from continuity there exists some $h'_j \in [t_{j-1}, t_j]$ such that $W(\ell_j) = F(\ell_j(h'_j)) \bullet \vec{v}_j$. The idea of the nonconstant force field over a line segment is embedded into the the NSP-world by *-transfer assuming that what has been established above holds for all such polygonal curves. Hence, let $\mathcal{P}_\Omega$ be a hyperpolygonal representation for the curve generated by a fine partition, ℓ_j an hyperline segment in $\mathcal{P}_\Omega$. Since F is continuous on E then $*F$ is defined on $\mathcal{P}_\Omega$. It follows that $*W(\ell_j) = *F(\ell_j(h'_j)) \bullet \vec{v}_j$. For polygonal curves, in general, the work done is an additive function. Thus for the hyperpolygonal curve $\mathcal{P}_\Omega$

$$*W(\mathcal{P}_\Omega) = \sum_{j=1}^{\Omega} *F(\ell_j(h'_j)) \bullet \vec{v}_j. \quad (1)$$

The following is established within this derivation and in the Infinitesimal Modeling manual. Let $c: [a, b] \rightarrow \mathbb{R}^n$ be a continuous differentiable curve with graph C and assume that $c'(t) \neq 0$ for each $t \in [a, b]$. Assume that uniformly continuous $F: E \rightarrow \mathbb{R}^n$, open $E \supset C$. The work done in moving through the force field on the path C is

$$W(C) = \text{st} \left(\sum_{j=1}^{\Gamma} *F(\ell_j(h'_j)) \bullet \vec{v}_j \right) = \int_C F \bullet d\vec{R}, \quad (2)$$

where the integral is the line integral over c and $\mathcal{P}_\Gamma$ is any hyperpolygonal representation for the curve C . Thus, under the hypotheses given, the work done in the NSP-world moving along these hyperpolygonal curves is indistinguishable, in general, from what is accepted as the N-world work in traveling along the curve. $\diamond$

{*Second derivation.* $\diamond$ See the second derivation for the general impulse in section 2.6 and modify the above accordingly. $\diamond$ }

It is significant to realize that even though the line integral will exist under less constraints and one could extend the concept of energy, to say curves that are rectifiable but not smooth, this need not imply that there is a meaningful correlation between this extended concept and its simplistic restriction to polygonal curves. On the other hand, the above derivation once again utilizes a maximum and minimum approach relative to a basic geometric element.

2.6 General Impulse

The energy or work done is considered to be the standard part of any one of a collection of hyperfinite sums each term of which is modeled by the amount of energy expended moving along a hyperline segment through a constant force field. The hyperfinite sums may be manipulated internally as if they were finite sums and preserve the intuitive concept of finite additivity. Unfortunately, when an object is represented by a simple differential equation obtained by such methods as illustrated in section 2.1, 2.2, and 2.3 then many authors solve such expressions by elementary differential equations methods without given full infinitesimal meaning to the process involved. The Fundamental Theorem of Integral Calculus is relevant to the existence of such solutions; but, it seems, that in order to grasp the NSP-world significance of the concept being measured that due consideration should be given to the integral solution rather than simply expressing the result by means of a rote procedure. As an illustration of this consider the extension of the concept of the impulse.

Previously the impulse was a natural notion restricted to constant force fields. A scalar force F , when applied for a period of time Δt produces an altered momentum Δp . This leads to the

expression $F\Delta t = \Delta p$. Then $F\Delta t = I$ is defined as the impulse. The extension to vector notation is obvious $\vec{I} = \vec{F}\Delta t = \Delta\vec{p}$, with a meaningful measure being the Euclidean norm. Applying the infinitesimalizing process (2), (3), (4) and (5) of section 2.3, where $C(v(t_2) - v(t_1))$ is replaced by $\vec{p}(t_2) - \vec{p}(t_1)$ leads to the customary elementary derivative expression

$$\vec{F} = \frac{d\vec{p}}{dt}. \quad (1)$$

Of course, the vector $\vec{F} = (p'_1, \dots, p'_n)$ where the p_i are the components for the momentum vector $\vec{p}$.

As far as a generalization for the impulse is consider there are various approaches. The Self-evident Max. and Min. Theorem 6.2.4 in the Infinitesimal Modeling manual applied to the components of $\vec{p}$ leads to the conclusion, if one has an intuitive comprehension of the basic additive properties for the impulse concept, that the proper expression for the general impulse for a continuous time dependent force field $\vec{F}$ over the time interval $[a, b]$ should be $\vec{I} = \int_a^b \vec{F} dt$. However, it may be more motivational and instructive to consider, once again, hyperpolygonal paths of motion since momentum is modeled after the motion concept and the concept of hyperfinite summation only. The summation of the measures of elementary physical processes still remains a primary modeling procedure that dictates the overall physical effect.

◇ Suppose, as in section 2.5, that uniformly continuous force field $F: E \rightarrow \mathbb{R}^n$, open $E \subset \mathbb{R}^n$, $F = (F_1, \dots, F_n)$. For a line segment $\ell' \subset E$, determine by a linear system of equations over the time interval $[t_1, t_2]$ and for a component F_i equation (2) section 2.3 can be re-expressed in terms of an impulse vector $\vec{I} = (I_1, \dots, I_n)$ as $F_i(t_m)(t_2 - t_1) = I_{im} \leq F_i(t)(t_2 - t_1) \leq F_i(t_M)(t_2 - t_1) = I_{iM}$. Thus there is some $t'_2 \in [t_1, t_2]$ such that the actual i 'th component of the impulse over that line segment is $I_i(t'_2) = F_i(t'_2)(t_2 - t_1)$. By *-transfer this holds for any hyperline segment. **Over a finite polygonal line ℓ it is assumed that the total i 'th component of the impulse is the simple sum of the i 'th component over the individual line segments.** Considering the impulse to be a function defined over the set of all finite polygonal curves in E , then for any hyperpolygonal curve $\mathcal{P}_\Gamma$

$$^*I_i(\mathcal{P}_\Gamma) = \sum_{j=1}^{\Gamma} ^*F_i(t'_j)(t_j - t_{j-1}), \quad (2)$$

where $\{t_0, \dots, t_\Gamma\}$ is the fine partition of $^*[a, b]$ that generates $\mathcal{P}_\Gamma$.

Now from uniform continuity, $^*F_i(t'_j) = ^*F_i(t_j) + \epsilon_j$, $\epsilon_j \in \mu(0)$. Since any finite set of real numbers contains a maximum, then any hyperfinite set of hyperreal numbers contains a maximum. Thus there exists some $\epsilon = \max\{\epsilon_1, \dots, \epsilon_\Gamma\} \in \mu(0)$ and following the elemental derivation process from the Infinitesimal Modeling manual, section 8, we have

$$\left| \sum_{j=1}^{\Gamma} \epsilon_j(t_j - t_{j-1}) \right| \leq \sum_{j=1}^{\Gamma} |\epsilon_j| (t_j - t_{j-1}) \leq |\epsilon| \sum_{j=1}^{\Gamma} (t_j - t_{j-1}) = |\epsilon| (b - a) \Rightarrow$$

$$\lambda_i = \sum_{j=1}^{\Gamma} \epsilon_j(t_j - t_{j-1}) \in \mu(0). \quad (3)$$

Substitution into equation (2) yields

$$^*I_i(\mathcal{P}_\Gamma) = \left(\sum_{j=1}^{\Gamma} ^*F_i(t_j)(t_j - t_{j-1}) \right) + \lambda_i, \quad (4)$$

Now F_i is continuous. Thus $\text{st}(*I_i) = \text{st}(\sum_{j=1}^{\Gamma} *F_i(t_j)(t_j - t_{j-1}))$ exists, by Theorem 5.1.2 of the Infinitesimal Modeling manual, since $\{t_1, \dots, t_{\Gamma}\}$ is an internal intermediate partition and $\text{st}(*I_i) = \int_a^b F_i(t) dt$. Notice that the hyperpolygonal curve was only an auxiliary notion since this last result is determined by a fine partition of $*[a, b]$ and is, indeed, the same for all such fine partitions. Extending the impulse to the vector $\vec{I}$, then $\vec{I} = \text{st}(*\vec{I}) = \int_a^b \vec{F}(t) dt$. $\diamond$

{*Second derivation.* First, it should be obvious that there is no complete classical counterpart to the nonstandard language used in an elemental derivation. Recall, however, how it is possible to use a quasi-classical language in the second derivation given in section 2.1. (i) As is done in Internal Set Theory, the “*” notation is removed from the functions since whether they are nonstandard extensions of standard functions is clear from the function’s argument (i.e. preimage).

(ii) The symbols “ $\approx$ ” is translated by the term “infinitely close.” This relation can be physically characterized by stating that no standard machine can measure any difference between the quantity on the left and the quantity on the right no matter how small the machine error.

(iii) Except for $\mathbb{N}_{\infty}$, hyperreal numbers are usually limited. Hence, simply call such a hyperreal number by the single word term “number.”

(iv) We use the fact that functions defined and continuous on $[a, b]$ preserve the infinitely close concept for these numbers. That is if $t, t_1 \in [a, b]$ and $t \approx t_1$, then $f(t) \approx f(t_1)$. Infinitesimals may be called the “infinitely or very small.” These numbers can be physically characterized as measures that are smaller than any standard machine error — measures that appear to a machine to be zero.

(v) Rather than use the standard part operator, where applicable, use the simple term “limit” in its place, since it has the same operative properties. Also use the fact that limits of two infinitely close numbers are equal.

For the elemental derivation method some additions to this quasi-classical language are necessary. These additions will necessarily be somewhat less precise for the concepts of the first-order property and the internal object will not be mentioned.

(vi) Let the region J be any of those studied in elementary calculus.

(vii) Hyperline segments are termed “infinitely small line segments,” which means line segments of infinitesimal length.

(viii) Hyperpolygonal curves may be called “infinitesimal polygonal curves” and defined as “polygonal curves with each line segment being an infinitely small segment.”

(ix) Call a fine partition of an interval an “infinitesimal partition”. This means that it has a $\Gamma + 1$ number of elements that determine Γ subintervals each of which is infinitely small in length.

(x) Hyperfinite sums are termed as “special finite sums” and behave as if they are finite sums.

(xi) Hyperfinite sets are termed as “special finite sets” and they also behave as if they are finite.

(xii) For the special finite sets, or special finite sums, Γ denotes the “number” of elements or terms, respectively.

(xiii) The product of an infinitesimal and a number [assuming limited] is an infinitesimal.

(xiv) Any nonnegative number less than or equal to an infinitesimal is an infinitesimal.

$\diamond$ Suppose, as in section 2.5, that uniformly continuous force field $F: E \rightarrow \mathbb{R}^n$, open $E \subset \mathbb{R}^n$, $F = (F_1, \dots, F_n)$. For a line segment $\ell' \subset E$, determine by a linear system of equations over the time interval $[t_1, t_2]$ and for a component F_i equation (2) section 2.3 can be re-expressed in terms of an impulse vector $\vec{I} = (I_1, \dots, I_n)$ as $F_i(t_m)(t_2 - t_1) = I_{im} \leq F_i(t)(t_2 - t_1) \leq F_i(t_M)(t_2 - t_1) = I_{iM}$. Thus there is some $t'_2 \in [t_1, t_2]$ such that the actual i ’th component of the impulse over that line segment is $I_i(t'_2) = F_i(t'_2)(t_2 - t_1)$. This result holds for any infinitesimal line segment. **Over a finite polygonal line ℓ it is assumed that the total i ’th component of the impulse is the simple**

sum of the i 'th component over the individual line segments. Considering the impulse to be a function defined over the set of all finite polygonal curves in E , then this i 'th component of the impulse is defined on an infinitesimal polygonal curve $\mathcal{P}_\Gamma$ with Γ many sides and its value is the special finite sum

$$I_i(\mathcal{P}_\Gamma) = \sum_{j=1}^{\Gamma} F_i(t'_j)(t_j - t_{j-1}), \quad (2)$$

where $\{t_0, \dots, t_\Gamma\}$ is an infinitesimal partition of $[a, b]$ with infinitely small subintervals that generates $\mathcal{P}_\Gamma$.

Now since the above results hold for these subintervals and $t' \in [t_{j-1}, t_j]$ for $1 \leq j \leq \Gamma$ then $t' \approx t_j$. Hence, $F_i(t'_j) \approx F_i(t_j)$ and from the definition of $\approx$ for each j there exists some infinitely small ϵ_j such that $F_i(t'_j) = F_i(t_j) + \epsilon_j$. Since any finite set of real numbers contains a maximum, then any special finite set of numbers contains a maximum. But, $\{\epsilon_1, \dots, \epsilon_\Gamma\}$ is a special finite set. Thus there exists some infinitely small $\epsilon = \max\{\epsilon_1, \dots, \epsilon_\Gamma\}$. Recall that what holds true for finite sums holds true for special finite sums. Thus the triangle inequality holds. Consequently

$$\left| \sum_{j=1}^{\Gamma} \epsilon_j(t_j - t_{j-1}) \right| \leq \sum_{j=1}^{\Gamma} |\epsilon_j| (t_j - t_{j-1}) \leq |\epsilon| \sum_{j=1}^{\Gamma} (t_j - t_{j-1}) = |\epsilon| (b - a)$$

But, since $|\epsilon|(b - a)$ is infinitely small then

$$\lambda_i = \sum_{j=1}^{\Gamma} \epsilon_j(t_j - t_{j-1}) \quad (3)$$

is infinitely small. Substitution into equation (2) and again using the fact that a special finite sum behaves like a finite sum yields

$$\begin{aligned} I_i(\mathcal{P}_\Gamma) &= \left(\sum_{j=1}^{\Gamma} (F_i(t_j) + \epsilon_j)(t_j - t_{j-1}) \right) = \\ &= \left(\sum_{j=1}^{\Gamma} F_i(t_j)(t_j - t_{j-1}) \right) + \lambda_i, \end{aligned} \quad (4)$$

Hence,

$$I_i(\mathcal{P}_\Gamma) \approx \left(\sum_{j=1}^{\Gamma} F_i(t_j)(t_j - t_{j-1}) \right). \quad (5)$$

Now F_i being continuous implies from the definition of the integral that $\lim \left(\sum_{j=1}^{\Gamma} F_i(t_j)(t_j - t_{j-1}) \right) = \int_a^b F_i(t) dt$. Taking the limit of expression (5) yields that $I_i = \int_a^b F_i(t) dt$. Extending the impulse to the vector $\vec{I}$, then $\vec{I} = \lim \vec{I} = \int_a^b \vec{F}(t) dt$. $\diamond$

There are, at least, two ways used to arrive at the relation between the change of momentum $\vec{p}(b) - \vec{p}(a)$ and the impulse $\vec{I}$. The first is the usual one of simply stating that $\vec{I} = \int_a^b \vec{F}(t) dt = \int_a^b (d\vec{p}/dt) dt = \vec{p}(b) - \vec{p}(a)$ from the Fundamental Theorem of Integral Calculus. There is a second method utilizing hyperfinite summation that incorporates the fact that F_i may be considered

as uniformly continuous on $[a, b]$. Since this notion is significant for the proper intuitive understanding of the underlying physical processes, we give an illustration using the momentum vector $\vec{p} = (p_1, \dots, p_n)$.

◇ Start with equation (2) and substitute ${}^*p'_i(t'_j)$ for $F_i(t'_j)$. This obtains

$${}^*I_i(\mathcal{P}_\Gamma) = \sum_{j=1}^{\Gamma} {}^*p'_i(t'_j)(t_j - t_{j-1}), \quad (5)$$

Then by the nonstandard mean value theorem there exists some $h'_j \in (t_{j-1}, t_j)$ such that

$${}^*p_i(t_j) - {}^*p_i(t_{j-1}) = {}^*p'_i(h'_j)(t_j - t_{j-1}). \quad (6)$$

But uniform continuity of p'_i on $[a, b]$ implies that there exists some $\delta_j \in \mu(0)$ such that ${}^*p'_i(h'_j) = {}^*p'_i(t'_j) + \delta'_j$. The elemental derivation process then yields

$$\delta_i + \sum_{j=1}^{\Gamma} {}^*p'_i(h'_j)(t_j - t_{j-1}) = \sum_{j=1}^{\Gamma} {}^*p'_i(t'_j)(t_j - t_{j-1}). \quad (7)$$

Substituting (6) into (7) and again using the elemental derivation process yields

$$\delta_i + p_i(b) - p_i(a) = \sum_{j=1}^{\Gamma} {}^*p'_i(t'_j)(t_j - t_{j-1}) = {}^*I_i(\mathcal{P}_\Gamma), \quad \delta_i, \lambda_i \in \mu(0). \quad (8)$$

Consequently, $p_i(b) - p_i(a) = \mathbf{st}({}^*I_i(\mathcal{P}_\Gamma))$ implies the result sought that $\vec{p}(b) - \vec{p}(a) = \mathbf{st}({}^*\vec{I}(\mathcal{P}_\Gamma)) = \vec{I}$. ◇

{*Second derivation.* ◇ Modify the second derivation for the general impulse integral.} ◇

Once again, the above illustrates that the total impulse is not dependent upon the hyperpolygonal path along which the object moves, but is simply the change in momentum. However, by physical intuition, an object has changed its momentum by traversing a physical path C and we are using a hyperpolygonal representation for such a path of motion. For any two such hyperpolygonal curves $\mathcal{P}_\Gamma, \mathcal{P}_\Lambda$, it follows from the hypotheses that ${}^*\vec{I}(\mathcal{P}_\Gamma) \approx {}^*\vec{I}(\mathcal{P}_\Lambda)$. Now the basic geometric measure for C , the length, is from the derivation in the Infinitesimal Modeling manual completely determined by the * -length of each of these hyperpolygonal representations. Therefore, it seems appropriate to consider the standard impulse over the curve C to be the unique effect $\mathbf{st}({}^*\vec{I}(\mathcal{P}_\Gamma))$, where $\mathcal{P}_\Gamma$ is any hyperpolygonal representation for C .

Why does the elemental derivation process work and what is it indicating about integral styled quantities? All of the basic quantities in any expression prior to considering their hyperfinite sum must not only be infinitely close but must be, in the above case, infinitely close of order one. The hyperfinite summation of infinitesimals need not be infinitesimal or even limited; but, depending on the type of limited objects this type of special infinite closeness utilizes, then the elemental derivation process does imply that this particular hyperfinite sum of infinitesimals is infinitesimal. In order to guarantee that this is the case, strong hypotheses were required for the functions involved. Except for the possibility of restricting the hyperfinite summation to special sets of partitions, such as in the case of the gauge type integral discussed in section 8.9 of the Infinitesimal Modeling manual, it, at present, has not been possible to obtain rigorous derivations for integral expressions without these strong hypotheses.

SLIGHTLY LESS BASIC MECHANICS

3.1 Mass

Since the Infinite Sum Theorems, the rules IR1–IR6 and the Self-Evident Theorems that appear in the Infinitesimal Modeling manual, have not been known to the physicist previously then almost all of the elementary definitions or derivations that involve infinitesimal quantities and the integral have relied upon elemental methods. These methods refer directly to the “particle”, point charges, point masses and other such notions. This process obviously forces the measure to have the same properties as the integral and the converse of starting with a functional, considering fundamental properties and showing that such a functional must be a measure as defined by some integral need not be considered. The mathematician might find this converse approach as appealing. But, it is more of a global view for a particular scenario and elementary instruction in the physical sciences tends to force upon a student an atomistic view in the sense that complex observed behavior is conceived of as the effects produce by minuscule objects. This is the prevalent textbook approach.

The Self-Evident Theorems that appear in the Infinitesimal Modeling manual require, at least, three global assumptions as well as a strong additivity property for the measure under consideration. The simplifications for regions that have Jordan-content, Propositions 6.7 and 6.10 as they appear in appendix 6, are very easy to apply in the laboratory setting since one need only investigate these properties on a rectangular interior to the region. However, this does require the assumption that the measure being considered is Jordan-like. Theorem 7.2.2 in the Infinitesimal Modeling manual implies that the integral over a Jordan-measurable region is an ordinary Jordan-like measure. Further, all of the customary geometric regions used within elementary physics are all compact Jordan-measurable sets and these simplified self-evident theorems may be applied. But the technical difficulties of establishing these simplified self-evident theorems should not be underestimated as amply illustrated by the formal proofs that appear in appendix 6 of the Infinitesimal Modeling manual. These proof methods are well beyond almost all undergraduates who require even advance physics courses as part of their curriculum.

In the Infinitesimal Modeling manual the Infinite Sum Theorem is the only approach used to obtain the mass of an object as determined by a continuous density function and this approach does not appear in any present day physics textbook. Consequently, it is obvious that, for the present, the elemental method with the notion of hyperfinite summation is more appropriate. As discussed towards the end of section 2.6, unless we restrict integral modeling to rectangular regions then, even within the elemental derivation, it is necessary to assume that the quantity being measured is Jordan-like. A brief review of this concept is useful.

Let R be an n -dimensional rectangle and $R \subset \mathbb{R}^n$. Suppose that Jordan-measurable $J \subset R \subset \mathbb{R}^n$, where to avoid trivialities we always assume that the interior of J is nonempty. Let $\mathcal{P}$ be a set of partitions of R , where, as usual, if $P \in \mathcal{P}$, then each $S \in P$ is but an n -dimensional subrectangle in R . Define on $\mathcal{P}$ the following map “inn.” For each $P \in \mathcal{P}$, let $\text{inn}(P) = \{S \mid S \in P \text{ and } S \subset J\}$. Of course, the set $\text{inn}(P)$ is called a set of **inner subrectangles**. Let $I_{\mathcal{P}} = \{\cup A \mid \exists P(P \in \mathcal{P} \wedge A = \text{inn}(P))\} \cup \{S \mid \exists P(P \in \mathcal{P} \wedge S \in \text{inn}(P))\}$. When considering the nonstandard extension of inn to a fine partition Q we usually do not write this as ${}^*\text{inn}(Q)$ but rather retain the original notation $\text{inn}(Q)$. Since any fine partition Q is hyperfinite and the set $\text{inn}(Q)$ is an internal subset of Q then it is also hyperfinite. For simplicity, as far as the functional (i.e. measure) $B: \{I_{\mathcal{P}}, J\} \rightarrow \mathbb{R}$ is concerned,

when the elemental derivation method is used it is assumed that B is, at least, simply additive on each $\text{inn}(P)$, when $P \in \mathcal{P}$. This means that if $P \in \mathcal{P}$ and $A = \text{inn}(P)$, then $B(\cup A) = \sum_{S \in A} B(S)$.

In general, if a generating function for the above functional B is constant over the entire Jordan-measurable J , then it is necessary that B display certain Jordan-measure properties. In particular, if v denotes the Jordan-measure, then for any fine partition Q of ${}^*R$, it follows from Theorem 7.2.2 that $v(J) = \text{st}(\sum_{S \in A} {}^*v(S)) = \text{st}({}^*v(\cup S))$. Thus, $v(J) \approx {}^*v(\cup A)$. Of course, ${}^*v(S)$ is but the product of the infinitesimal lengths of the sides of the n -dimensional subrectangle S . Bearing in mind the basic properties of Jordan-measure, we assume that for a fine partition Q that the functional B has the property that $B(J) \approx {}^*B(\cup A)$, where $A = \text{inn}(Q)$. This property is termed the **Jordan-like** property and a standard criterion for the functional B to be Jordan-like for all fine partitions is given in the Infinitesimal Modeling manual. Of course, these technical matters need made be discussed in most elementary courses. *Indeed, in the second derivations one only needs to state that the region J is one of those studied in the elementary calculus.*

◇ Let compact and Jordan-measurable $J \subset R \subset \mathbb{R}^3$ and S be an inner rectangle as defined above. Let $\rho: J \rightarrow \mathbb{R}^3$ be a continuous density function. If $\rho(x, y, z)$ is constant over a subrectangle S of J , then from the definition of ρ the mass $m(S) = \rho v(S)$, where $v(S)$ is the volume of S . Suppose that ρ is not constant. Then there exists $(x_m, y_m, z_m) \in S$ and $(x_M, y_M, z_M) \in S$ such that $\rho_m = \rho(x_m, y_m, z_m) \leq \rho(x, y, z) \leq \rho(x_M, y_M, z_M) = \rho_M$ for each $(x, y, z) \in S$. Hence, if the mass of $S = m(S)$, then observation and the above property for constant density indicates that, at least, from the macroscopic viewpoint

$$\rho_m v(S) \leq m(S) \leq \rho_M v(S). \quad (1)$$

The Intermediate Value Theorem for connected subsets of R yields that there is some $(x_1, y_1, z_1) \in S$ such that $m(S) = \rho(x_1, y_1, z_1) v(S)$. If we have more than one such inner rectangle, say $\{S_1, \dots, S_n\}$, than observation indicates that the total mass, $m(A)$, of the configuration $A = S_1 \cup \dots \cup S_n$ is the finite sum

$$m(A) = \sum_{j=1}^n m(S_j) = \sum_{j=1}^n \rho(x_j, y_j, z_j) v(S_j). \quad (2)$$

By * -transfer, these results hold for infinitesimal subrectangles of J and hyperfinite summation. Now let Q be any fine partition of ${}^*R$. Then $\text{inn}(Q)$ is an internal hyperfinite set of infinitesimal subrectangles. Hence,

$${}^*m(\cup \{S_j | S_j \in \text{inn}(Q)\}) = \sum_{j=1}^{\Gamma} {}^*\rho(x_j, y_j, z_j) {}^*v(S_j). \quad (3)$$

Even though each point $\vec{v}_j = (x_j, y_j, z_j) \in S_j$, this does not imply that the intermediate partition $\{\vec{v}_j \mid 1 \leq j \leq \Gamma\}$ is internal. However, letting $\vec{c}_j$ denote the corner of S_j nearest or equal to $(0, 0, 0)$ then $\{\vec{c}_j \mid 1 \leq j \leq \Gamma\}$ is an internal intermediate partition. Uniform continuity of ρ on J yields that for each j there exists some $\epsilon_j \in \mu(0)$ such that ${}^*\rho(x_j, y_j, z_j) = {}^*\rho(\vec{c}_j) + \epsilon_j$. Hence,

$${}^*m(\cup \{S_j | S_j \in \text{inn}(Q)\}) = \sum_{j=1}^{\Gamma} ({}^*\rho(\vec{c}_j) + \epsilon_j) {}^*v(S_j). \quad (4)$$

The elemental derivation process yields

$${}^*m(\cup\{S_j | S_j \in \text{inn}(Q)\}) \approx \sum_{j=1}^{\Gamma} ({}^*\rho(\vec{c}_j) {}^*v(S_j)). \quad (5)$$

Since ρ is integrable over J then taking the standard part and applying Theorem 7.2.2 of the Infinitesimal Modeling manual one obtains

$$\text{st}({}^*m(\cup\{S_j | S_j \in \text{inn}(Q)\})) = \int_J \rho(x, y, z) dX. \quad (6)$$

As mentioned above, one final step is required. It must be assumed that the concept of the mass of an object has the same Jordan-like quality as does the concept of the volume. Making this last assumption implies that $m(J) = \int_J \rho(x, y, z) dX$. $\diamond$

A physical interpretation of the Jordan-like quality of the mass is also possible. For every fine partition Q , the NSP-world effects of the mass of a simple internal rectangular configuration $\cup\{S_j | S_j \in \text{inn}(Q)\}$ is indistinguishable from the effects of the mass of J at the first (order) level, if no comparison is to be made with other such measures. When such a comparison is made, as pointed out in the Infinitesimal Modeling manual, Section 8.3, the effects of ${}^*m(\cup\{S | S \in \text{inn}(Q)\})$ and $m(J)$ would be infinitely close of order 3 and indistinguishable at that level. Of course, all that is being derived for the case $\mathbb{R}^3$ holds for ${}^*m(\cup\{S | S \in \text{inn}(Q)\})$ and any $\mathbb{R}^n$, $n \geq 1$.

In all of the previous derivations, equations of form (1) are the essential modeling requirements. The effect of the elemental derivation process is to eliminate the additional additivity property for such functionals m . However, if such additivity is assumed, then the integral expression follows immediately from the self-evident theorems that appear in the Infinitesimal Modeling manual.

{*Second derivation.* Now that we have the essential procedures needed to modify a rigorous infinitesimally styled derivation and obtained a quasi-classical one, it is not necessary to present an entire second derivation for equation (6). It is useful to conjoin the following to our list of alterations. The set J can simply be described as one of the regions studied in the elementary calculus and for which an integral expression for volume is obtained. For the necessity of expressing $\rho(x_j, y_j, z_j)$ in equation (3) as $\rho(x_j, y_j, z_j) = \rho(\vec{c}_j) + \epsilon_j$, without invoking the concept of the internal intermediate partition, one can simply argue that it is necessary to evaluate ρ at a known partition generated point rather than at a less explicitly known $(x_j, y_j, z_j) \in S$. The requirement that m be Jordan-like can be described as a common feature that m must share with the volume of the region J . The feature being, that for any infinitesimal partition the configuration composed of all of the infinitely small subrectangles contained within J must have its volume infinitely close the volume of J . This same feature holds for the mass and other such measures since if $\rho \equiv 1$ on J , then $v(J) = m(J)$. I leave to the reader the actual construction of the second derivation.}

3.2 Moments and Center of Mass

It is often difficult to decide whether it is more significant to derive a specific mathematical model completely from fundamental physical observations and the defining properties of the mathematical structure or to replace portions of the derivation with well-known theorems gleaned from the abstract structure itself. It appears that within modern theoretical physics, abstract mathematical results are introduced as soon as practical even though they may be couched in a quasi-physical language. This

procedure follows the routine assumption that the rigorous logic displayed when a proposition is proved abstractly is equivalent to the logic required for a much longer and more complex derivation that utilizes but the fundamental correspondence between the defining properties of the structure and the physical terms that describe the physical scenario. The following theorem is useful in order to illustrate the economy achieved by such an early introduction of well-known mathematical results.

Theorem 3.2.1. A Weighted Mean Value Theorem. *Let $J \subset E \subset R \subset \mathbb{R}^n$, J be Jordan-measurable, and E be compact and connected. Let continuous $f: E \rightarrow \mathbb{R}$, integrable $g: J \rightarrow \mathbb{R}$ and $g(\vec{v}) \geq 0$ for each $\vec{v} \in J$. Then there exists some $\vec{v}_0 \in E$ such that $\int_J f(\vec{v})g(\vec{v}) dX = f(\vec{v}_0) \int_J g(\vec{v}) dX$.*

Even though the elemental derivation process is being stressed throughout this physics manual, it is worthwhile to once again mentioned that the Self-Evident theorems of the Infinitesimal Modeling manual are always available. The Method of Constants, as well as the Maximum-Minimum Method, is actually exemplified within the elemental process derivations. Immediately following equation (1) of this section is the fact that there does exist some $(x_1, y_1, z_1) \in S$ such that $m(S) = \rho(x_1, y_1, z_1)v(S)$. This is the explicit requirement for application of Proposition 6.9 in Appendix 6 assuming the requisite additivity properties for the measure m . If the measure generating function is composed of the product of two or more nonconstant functions, then the Extended Self-Evident Method of Constants Proposition 6.10 may be appropriate.

◇ Notwithstanding our discussion in the above paragraph, let's consider the a moment generating function $M_{(\cdot)}$ defined as follows: Let nonnegative continuous $\rho(x, y, z): J \rightarrow \mathbb{R}$, where $J \subset R \subset \mathbb{R}^3$ is compact and Jordan-measurable. A moment function defined on a partition subrectangle $S \subset J$ is $M_{yz}(S) = x \rho(x, y, z) v(S)$, where $x = f(x, y, z)$ is continuous on $\mathbb{R}^3$. In the same manner, define $M_{xz}(S) = y \rho(x, y, z) v(S)$, $M_{xy}(S) = z \rho(x, y, z) v(S)$. Although it may appear to be sufficient to consider the point (x, y, z) as an arbitrary member of S , certain special selections are necessary. For example, if ρ is constant, then (x, y, z) might be selected as the center of S . It is well-known that if we consider a small enough homogeneous rectangular solid S , then the gravitational field of the Earth in relation to S is effectively a parallel vector field and the acceleration of gravity is a constant g . The center of the rectangle is the point of rotational stability within such a field — the so-called center of gravity. This is demonstrated in the customary manner by considering the expressions $g M_{xy}(S)$, $g M_{yz}(S)$, $g M_{xz}(S)$. Thus from the Newtonian gravitational point of view, the moment function can be viewed as a measure of the rotational effect with respect to the coordinate planes within such a gravitational field. Notice that the necessary observations that lead to this conclusion are relative to the actual small size of the objects.

For the general case of continuous ρ on S , there exist $x_m \rho(x_m, y_m, z_m)$ and $x_M \rho(x_M, y_M, z_M)$, where $(x_m, y_m, z_m), (x_M, y_M, z_M) \in S$ such that

$$x_m \rho(x_m, y_m, z_m) v(S) \leq x \rho(x, y, z) v(S) \leq x_M \rho(x_M, y_M, z_M) v(S). \quad (1)$$

The usual assumption is now imposed upon our problem. Suppose that the actual moment effect $M_{yz}(S)$ lies somewhere between these two extremes. Thus there would exist some $(x'_1, y'_1, z'_1) \in S$ such that

$$M_{yz}(S) = x'_1 \rho(x'_1, y'_1, z'_1) v(S). \quad (2)$$

Equation (2) is now extended to a finite system $\{S_j \mid 1 \leq j \leq n\}$ of partition inner subrectangles using the apparent experiential result that the total moment effect of the system is the sum of the

individual effects. Hence,

$$M_{yz}(\cup S_j) = \sum_{j=1}^n x'_j \rho(x'_j, y'_j, z'_j) v(S_j). \quad (3)$$

By $*$ -transfer, we assume that within the NSP-world the above behavior holds for infinitesimal rectangles. Since this is a modeling technique it is not necessary to assume that such infinitesimal rectangles exist in some type of objective reality. *However, it is possible to describe such behavior within a substratum NSP-world by considering the simplistic behavior of a hyperfinite set of NSP-world infinitesimal rectangles as a superstructure of objects that controls the behavior of a corresponding system of “small” natural world rectangular objects.* As in the least section, assuming that Q is a fine partition of $*R$ this leads to the conclusion that

$$*M_{yz}(\cup\{S_j | S_j \in \text{inn}(Q)\}) = \sum_{j=1}^{\Gamma} x'_j * \rho(x'_j, y'_j, z'_j) * v(S_j). \quad (4)$$

Letting (x_j, y_j, z_j) denote the corner of S_j nearest or equal to $(0, 0, 0)$ then $\{(x_j, y_j, z_j) \mid 1 \leq j \leq \Gamma\}$ is an internal intermediate partition. Uniform continuity of $x\rho$ on J yields that for each j there exists some $\epsilon_j \in \mu(0)$ such that $x'_j * \rho(x'_j, y'_j, z'_j) = x_j * \rho(x_j, y_j, z_j) + \epsilon_j$. Applying the elemental derivation process one obtains

$$\text{st}(*M_{yz}(\cup\{S_j | S_j \in \text{inn}(Q)\})) = \int_J x \rho(x, y, z) dX. \quad (5)$$

The additional requirement that the measure of the moment is Jordan-like yields $M_{yz}(J) = \int_J x \rho(x, y, z) dX$.

Now $R \subset \mathbb{R}^3$ is compact and connected and $f(x, y, z) = x$ is continuous on R . Thus, by the Weighted Mean Value Theorem there exists some real $\bar{x}$ such that

$$M_{yz}(J) = \int_J x \rho(x, y, z) dX = \bar{x} \int_J \rho(x, y, z) dX = \bar{x} m(J). \quad (6)$$

Repeating the above argument there exist real $\bar{y}$ and $\bar{z}$ such $M_{xz}(J) = \int_J y \rho(x, y, z) dX = \bar{y} \int_J \rho(x, y, z) dX = \bar{y} m(J)$ and $M_{xy}(J) = \int_J z \rho(x, y, z) dX = \bar{z} \int_J \rho(x, y, z) dX = \bar{z} m(J)$. Consequently, as far as the moment effects are concerned the object J can be consider as represented by the single point $(\bar{x}, \bar{y}, \bar{z})$ with the mass number $m(J)$ attached to it. $\diamond$

{*Second derivation.* From this point on in this Elementary Physics manual, the second quasi-classical derivation will not be given unless it is substantially different from our previous examples.}

Notice that the above derivation did not start with the concept of the point masses and then derive the integral expression for the center of mass. Rather, we derived the concept by means of infinitesimal analysis. Additionally, the statement that the center of mass is equivalent to the center of gravity appears in the above discussion to depend upon the parallel gravitational field concept. In the next section, we show that the idea of the less substantiated point masses, *if viewed from the NSP-world*, does lead to the same center of mass conclusion.

3.3 Point Masses

In Tipler [1982], the concept of the point mass is used to develop the center of gravity and center of mass for such objects, assuming that this technique has been justified. On page 229 of volume 1, Tipler states: “If the center-of-mass coordinates of a continuous body are to be calculated,

the sum $\sum m_i x_i$ must be replaced by the integral $\int x \, dm$, where dm is an element of mass.” In the text by Young, Riley, McConnell, Rogge [1974, p. 281], when moments of inertia are discussed, once again the student is instructed that such a measure is given by an integral over dm . No further explanation is given as to why this particular technique is justified. This vague modeling technique can be justified within the NSP-world once equations such as (6) of Section 3.2 have been derived.

◇ Assume the hypotheses used to derive (6) of Section 3.2 and let Q once again be a fine partition of ${}^*R$. Let $S_j \in \text{inn}(Q)$. Then there exists some $(x''_j, y''_j, z''_j) \in S_j$ such that ${}^*m(S_j) = {}^*\rho(x''_j, y''_j, z''_j) {}^*v(S_j)$. Now ${}^*\rho(x''_j, y''_j, z''_j) \approx {}^*\rho(x'_j, y'_j, z'_j)$, where (x'_j, y'_j, z'_j) is as described in the derivation of Section 3.2. Hence, $x'_j {}^*\rho(x'_j, y'_j, z'_j) \approx x'_j {}^*\rho(x''_j, y''_j, z''_j)$, for x'_j is limited. The elemental derivation process yields

$$M_{yz}(J) = \int_J x \rho(x, y, z) \, dX = \text{st} \left(\sum_{j=1}^r x'_j {}^*\rho(x''_j, y''_j, z''_j) {}^*v(S_j) \right). \quad (1)$$

Consequently, letting $dm_j = {}^*m(S_j) = {}^*\rho(x''_j, y''_j, z''_j) {}^*v(S_j)$ denote that mass of the infinitesimal subrectangle (not a point!) and repeating the argument for the other two moments one can describe the moment effects within the standard world as follows: The effect is indistinguishable from the effect of a hyperfinite sum of mass numbers attached to the points (x'_j, y'_j, z'_j) . Thus the points (x'_j, y'_j, z'_j) within the NSP-world can be viewed as point masses. Obviously, the point (x'_j, y'_j, z'_j) is not unique since it may be replaced by any $(x, y, z) \in S_j$. ◇

The reader might be inclined to attempt to * -transfer the above derivation to the standard world and arrive at the conclusion that there exists a set of point masses that would yield the moment effects expressed by the integrals. This would be an error, however, since the standard part operator and as well as $\approx$ are external concepts. The technique of * -transfer, at this stage, only allows the hyperfinite sum of point masses to be transferred into a statement about finitely many point masses the sum of the moments of which would approximate the moment effect within an given positive ϵ for all the partitions of R with mesh less than some positive δ . What this implies is that rather than accepting an ad hoc modeling technique that utilizes unrealistic standard world point masses to derive the integral expressions for the moment effects, it may be more conducive to student comprehension to employ the NSP-world point masses since their use can be more rigorously justified. *However, in certain cases once infinitesimal analysis has established equation (6) of Section 3.2, and the like, then standard means can be applied to investigate an effective center of mass for a finite collection of objects.* This we do next.

◇ Suppose that $\{J_1, \dots, J_p\}$ is a nonempty finite set of pairwise disjoint, compact, and Jordan-measurable subsets of $R \subset \mathbb{R}^3$. Further, let nonnegative continuous $\rho_i: J_i \rightarrow \mathbb{R}$, for each i such that $1 \leq i \leq p$. The function $f(x, y, z) = x$ is obviously continuous on compact, connected R . The piecewise well-defined function $h(x, y, z) = \rho_i(x, y, z); (x, y, z) \in J_i, 1 \leq i \leq p$ is continuous on $J = J_1 \cup \dots \cup J_p$. [Since $\mathbb{R}^3$ is a normal topological space it follows that if $\vec{v} \in J_i$, then $\mu(\vec{v}) \cap J_j = \emptyset, i \neq j$.] Now the set J is compact; hence, closed and bounded. Consequently, h is a nonnegative bounded function defined on J . The function $f(x, y, z) = x$ is continuous on compact, connected R and integrable on J . Consider the moment effect generating function $m_{yz} = x h(x, y, z)$. Then from the Weighted Mean Value Theorem there exists $\bar{x}_j$ such that the moment effect

$$M_{yz}(J) = \int_J m_{yz} \, dX = \bar{x}_j m(J). \quad (2)$$

However,

$$\int_J m_{yz} dX = \sum_{i=1}^p \int_{J_i} x \rho_i(x, y, z) dX = \sum_{i=1}^p \bar{x}_i m(J_i). \quad (3)$$

Thus

$$M_{yz}(J) = \sum_{i=1}^p \bar{x}_i m(J_i). \quad (4)$$

Repeating the above derivation yields similar equations as (4) for the other two moments. However, I repeat, once again, that this approach is only relative to a nonempty finite set of disjoint, compact, and Jordan-measurable subsets of R . $\diamond$

3.4 Standard Rules and the Elemental Derivation Process

Although the elemental derivation process is very appealing to the intuition, what happens when this process is viewed as a mathematically stated theorem? If you were to analyze the standard hypotheses needed to model this process, then what would be obtained is the rule IR5 as it appears in the Infinitesimal Modeling manual. This rule coupled with the hypotheses stated in a theorem such as Proposition 6.7 in the Infinitesimal Modeling manual leads to a formal theorem that can be established not by an infinitesimal sum theorem but by the elemental derivation method. We develop such a theorem next — a theorem that allows us to eliminate the actual elemental derivation process. The necessary notation for what follows is defined in this manual.

We will not state what comes next as a formal theorem but state it somewhat informally. We point out that it is but a restatement in a slightly expanded form of Proposition 6.7 in the Infinitesimal Modeling manual. Let $\mathcal{P}$ be any set of partitions of the rectangle $R \subset \mathbb{R}^n$ and compact Jordan-measurable $J \subset R$. Suppose that $\mathcal{P}$ contains a fine partition. Of course, if $\mathcal{P}$ is the set of all simple partitions of R , then such a fine partition exists. Next let B be a real valued function(al) defined on $\{I_{\mathcal{P}}, J\}$ and, at least, additive on $\{I_{\mathcal{P}}, J\}$. Let continuous $f: J \rightarrow \mathbb{R}$. Suppose that for any $P \in \mathcal{P}$ and any $S \in \text{inn}(P)$, it follows that $(f_m)v(S) \leq B(S) \leq (f_M)v(S)$, where (f_m) [resp. (f_M)] is the minimal [resp. maximal] value of f on S and $v(S)$ is the Jordan-measure of S (i.e. its simple volume). Then if B has the ordinary Jordan-like property, it follows (from the elemental derivation process) that $B(J) = \int_J f(\vec{x}) dX$.

What the last paragraph signifies is that in all cases where elementary physical measures are concerned one needs only argue for the acceptance of the stated hypotheses. Once such hypotheses are accepted as reasonable, then the conclusion follows from both the infinite sum theorem or the elemental derivation process.

The assumption that B has the ordinary Jordan-like property is not difficult to accept. A standard criterion appears in the Infinitesimal Modeling manual and that property is apparently necessary in order for B to be obtained by an integral. This comes from the fact that the integral itself when viewed as a functional satisfies this property. It intuitively signifies that an approximation for the value of $B(J)$ that is better than any machine error can be obtained by considering the value $B(C)$, where $C \subset J$ is a configuration composed of subrectangles taken from a partition with “small enough” mesh. Now compare this with the requirements of IR3 (1) in the Infinitesimal Modeling manual and Theorem 6.2.4, where the ordinary Jordan-like property is not assumed. In this case, the maximal–minimal assumption is weakened for boundary rectangles. This weakening is relative to the value of the functional as extended to the boundary rectangles. In an elementary exposition, it may be more reasonable to “build” configurations such as C and accept the ordinary Jordan-like

property for such configurations, then to alter the intuitive acceptance of such maximal-minimal statements as (1) on page 22 of this manual.

Thus far, we have needed to include the strong requirements that the function f be continuous on J and that J be, at least, compact and Jordan-measurable. Can either or both of these requirements be relaxed and an acceptable derivation method developed for integral models? An answer to this question will depend upon what one considers as “acceptable” and the areas of application. We will attempt to answer this question in later sections of this manual.

To be continued by properly trained members of the physics community.

REFERENCES

- Gillispie, C.C.**, [1960], *The Edge of Objectivity*, Princeton University Press, Princeton, NJ.
- Herrmann, R. A.**, [1989], Fractals and ultrasmooth microeffects, *J. Math. Physics*, **30** (4): 805-808.
- Newton, I.**, [1934], *Mathematical Principles of Natural Philosophy*, (Translated by Cajori), University of Cal. Press, Berkeley.
- Tipler, P. A.**, [1982] *Physics*, Worth Publishers, Inc., New York.
- Young, Riley, McConnell**, [1974] *Essentials of Mechanics*, The Iowa State University Press, Ames, Iowa.

Additional Special Symbols

(Alphabetically listed by first symbol letter.)

Symbol.....	Name, if any	Page no.
$\sim_1$	Infinitely Close of Order One.....	170
$\sim_2$	Infinitely Close of Order two.....	172
$\vec{I}$	Impulse Vector	177
$m(S)$	Mass of S	182
$M_{(\cdot)}$	Moment Function	184
dm	Mass Of Infinitesimal $S =$ Element Of Mass	186

- Index for pages 1-153.
- Location is within ± 1 .
- Abel 6.
- acceptable set of partitions 49.
- additive 39.
- additive, simply 37.
- algebraists, information for 15, 17.
- algorithm 78.
- almost parallel 91.
- Apostal [1957] 35.
- applications, simple 23.
- Archimedean 10.
- Archimedes 6.

- Barwise [1977] 21.
- basic:
 - element 47.
 - elementary integral 48.
 - hyperfinite subsets 30.
 - laboratory experimentation 46.
 - region 39.
- bound formula 29.
- boundary subrectangles 62.
- boundedness concept 46.
- bounding method 23.
 - redefinition 24.

- caloric 98.
- Cauchy 6.
 - definition for the integral 35.
 - Principle 88.
- Cavalieri's 36.
- Cesari [1956] 71.
- compactness 32.
- compactness theorem 21.
- comprehend second order rates of change 98.
- conduction, heat 98.
- conductivity, thermal 99.
- consecutive points 84.
- constant values, extension fo 50.
- constant, use of 16.
- constants:
 - elemental method of 64.
 - extended standard 23.
 - external 23.
 - internal 20.
 - internal 23.
 - method of 50.
 - self-evident method of 51.
 - unstarred 23.
 - use of 28.
- continuity 32.
- continuous distribution of matter 49.
- continuous, uniformly 32.
- convergence, sequential 19.
- convex in the direction y 99.
- curvature, radius of 87.
- curve 31.
 - tangent to 71.
- curves, hyperpolygonal 73.
- Cutland [1986] 37.

- d'Alembert 6.
- d'Alembert - Euler 8.
- D-world 14.
- Darboux 35.
 - integral 35.
- de l'Hospital 6, 26.
- De Lillo [1982] 42.
- definition:
 - for Euclidean Spaces 11.
 - for infinite 11.
 - for infinitely close 11.
 - of hyperfinite 31.
 - of infinitesimals 10.
 - of limited 11.
 - of monad 12.
- deformable body rule 93.
- deleted monad 77.
- δ -fine partition 75.
- derivative, n-dimensional 90.
- differentiable 90.
- differential 89.
- differential equation:

- method of maximum and minimum 95.
- models 49.
- dot notation, Newton's 81.
- dynamical (loci) methods 78.
- electron-positron lattice 95.
- element:
 - basic 47.
 - hypertrapezoid 60.
 - infinitesimal 64.
 - surface 73.
- elemental method of constants 64, 76.
- elementary:
 - geometric 46.
 - prototype 47.
 - integral 48.
- elements 48.
 - geometric 59, 73.
 - m-dimensional 47.
 - method of 46, 76.
- energy expended 67.
- entity 17.
 - extended 23.
 - internal 23.
- ϵ -infinitesimal microscope 71.
- equivalent maps 88.
- Euclidean Spaces, definition, 11.
- Euclidean n-spaces 14.
- Eudoxus [370 BC] 27.
- Euler 6.
- evident 44.
- Example:
 - 3.3.1. 19.
 - 3.3.2. 19.
 - 3.3.3. 19.
 - 3.3.4. 20.
 - 3.4.1. 22.
 - 3.4.2. 22.
 - 3.4.3. An argument 23.
 - 4.2.1. 28.
 - 4.2.2. 28.
 - 4.2.3. 28.
 - 4.2.4. 28.
 - 4.3.1. 30.
 - 4.4.1.A. 33.
 - 4.4.1.B. 33.
 - 4.4.2. (Fractals) 33.
 - *-transfer 28.
- extended:
 - entity 23.
 - relations 17.
 - standard 17.
 - standard constants 23.
 - of constant values 50.
 - the finite 50.
- extensions of Jordan measures 39.
- exterior subrectangles 62.
- external constants 23.
- fine 35.
 - partitions 33.
- finite 30.
 - extension of 50.
 - summation 31.
- first-order:
 - ϵ -resolving power. 71.
 - ideals 69.
 - language 19, 20.
- flow lines, postulated properties of 44.
- flowing heat 98.
- fluid motion:
 - lines of 44.
 - tubes of 44.
- fluxions 78.
- force 76.
- formula, bounded 29.
- foundations of infinitesimal modeling 78.
- fractals and infinite length 33.
- free variable 28.
- function, limit of 77.
- Galilean physics 80.
- gauge integral 75.
- Gauss 1827; Art 20 46.
- geometric elements 59, 73.
 - for the curve 59.

- elementary 46.
 - surface element 73.
- geometry 76.
- H-integral 35.
- heat conduction 98.
- heat:
 - flowing 98
 - quantity of 98.
 - specific 99.
 - uniformly conducting 98.
- Henstock [1961] 75.
- Herrmann 1980 46.
 - [1985] 35.
 - [1986] 21.
 - [1989] 33.
- higher order increments, omission of 81.
- Hurd and Loeb [1985] 65.
- hyperfinite:
 - definition 31.
 - partition 35.
 - subsets, basic 30.
 - sum 31.
- hyperline segments 73.
- hyperparallelepiped 74.
- hyperpolygonal curves 73.
- hyperreals, names for 10.
- hyperrectangle, truncated 63.
- hypersurface 73.
- hypertangent planes 73.
- hypertrapezoid element 60.
- hypertrapezoids 73.
- ideal 6, 11.
 - maximal 15.
- idealized behavior 58, 76.
- ideals, order 68.
- identity, preserving 17.
- inconsistencies 6.
- increment:
 - local 90.
 - n th order 70.
- increments, higher order 81.
- individual 17.
 - internal 23.
- indivisible 18, 36.
- induction 20.
- inertia, moment of 49.
- infinite:
 - characterized 11.
 - definition 11.
 - length, fractals 39.
 - magnification operator 71.
 - magnification, of a infinitely small portion of the fluid 44.
 - numbers, different from extended real numbers 26.
 - Sum Theorems 37.
- infinitely close 49.
 - definition 11.
 - of order n 69, 71.
- infinitely small 6.
- infinitesimal:
 - approach, pure 58.
 - elements, physical 64.
 - max. and min. rule 48.
 - microscope 71
 - modeling, foundations 78.
 - parallelepiped 74.
 - reasoning 49, 51.
- infinitesimal:
 - rectangles 73.
- infinitesimalizing procedure 49.
- infinitesimals:
 - and local map 88.
 - definition 10.
 - nonnegative 69.
- inner subrectangles 62.
- integral:
 - Cauchy's definition 35.
 - elementary 48.
 - gauge 75.
 - line 66.
 - M 75.

- McShane 75.
- Riemann-complete 75.
- S 75.
- surface 73.
- intermediate partition 36.
- internal 22.
 - constant names 20.
 - constants 23.
 - definition principle 30.
 - entity 23.
 - individual 23.
 - objects and N-world modeling 30.
- Jarnik, Kurzweil, Schwabik [1983] 75.
- Jordan measures, extended to 39.
- Jordan-measurable 42, 47, 62.
- k -plane 85.
- $(k + 1)$ -consecutive 84.
- Keisler 8.
- Kepler 6.
- laboratory experimentation, basic 46.
- lattice structure 95.
- Leibniz 6, 8.
- Leibniz Principle 21.
 - restated 23.
- Leibniz [1701] 18.
- length by fine partitions 33.
 - by limit of a sequence 33.
- limit of a function 77.
- limit theorems 19.
- limited numbers definition 11.
- line Integral 66.
- linear, locally 88.
- lines of fluid motion 44.
- local:
 - increment 90.
 - map 48.
 - locally linear 88.
 - locus 76.
 - Luxemburg [1962] 15.
 - m-dimensional elements 47.
 - M-integral 75.
 - Machover and Hirschfeld [1969] 15.
 - magnification:
 - operator 71
 - infinite 44.
 - map:
 - equivalent 88.
 - local 88.
 - matter, continuous distribution of 49.
 - Mawhin [1985] 75.
 - max. and min. rule 48.
 - self-evident 49.
 - maximal ideal 15.
 - max. and min., differential equation method of 95.
 - Maxwell [1890] 44.
 - McShane [1973] 75.
 - McShane integral 75.
 - mean value 39.
 - metamathematically, the necessity of arguing 23.
 - method of constants 50.
 - elemental 76.
 - method of elements 76.
 - microconstruction 68.
 - microeffects 68.
 - microguage 68.
 - micropartition 75.
 - regular 75.
 - microscope, infinitesimal 71.
 - modeling:
 - differential equation 49.
 - infinitesimal 78.
 - internal objects 30.
 - N-world 30.
 - moment 78.

- moment of inertia 49.
- monad:
 - definition 12.
 - deleted 77.
- monadic:
 - restrictions of physical processes 94.
 - second law of motion 92.
 - second law of motion, alternative 101.
- Morley's example 52.
 - [1942] 52.
- motion 76.
- motion of points 76.
- moving objects 76.
- moving points, tubes of 44.

- n th difference 70.
- n th order ideal 69.
- n th order increment 70.
- n -dimensional derivative 90.
- N-world 14.
- N-world modeling and internal objects 30.
- names, internal constant 20.
- natural numbers, properties of 20.
- natural philosophy 76.
- non-Archimedean 31.
- non-finite 17.
- nonnegative infinitesimals 69.
- normal vector 73.
- NSP-world 14.
 - behavior, rules for pure 42.

- objective reality, both the infinitesimal and infinite 26.
- objects, internal constant names 20.
- observation, physical 76.
- order ideal, n th 69.
- order ideals 68.
- order pair 16.
- osculating:
 - k -circle 87.
 - k -plane 84.

- parallel, almost 91.
- parallelepipeds, 47.
 - infinitesimal 74.
- partition 35.
 - δ -fine 75.
 - fine 33, 35.
 - hyperfinite 35.
- partitions, acceptable set of 49.
- Peano 20.
- phenomenological approach 46.
- philosophy, natural 76.
- physical:
 - behavior, simplistic 46.
 - infinitesimal elements 64.
 - observation 76.
 - processes, monadic restrictions 94.
- Planck 6.
 - and meaning of physical series 34.
- plane, tangent 73.
- point definable quantities 50.
- point-motion 76.
- power set operator 17.
- predicates in set builder notation 28.
- preserves infinitesimals 88.
- properties of standard part operator 14.
- prototype, elementary 47.
- pure:
 - infinitesimal approach 58.
 - NSP-world behavior 42.

- quantifiers used in set builder notation 28.
- quantity of heat 98.
- quotient ring 15.

- radius of curvature 87.
- real numbers, positive 10.
- rectangle 35.
 - infinitesimal 73.
- region, basic 39.
- regular micropartition 75.

- relation, extended 17.
- Riemann:
 - integral 35.
 - Stieltjes Integral 65.
 - styled sum 58.
 - sums 35.
- Riemann-complete integral 75.
- ring 11.
- ring, quotient 15.
- Robinson 8.
 - [1961] 10, 26.
 - [1966] 15.
- Robinson and Zakon [1969] 15.
- rules 37.
 - for pure NSP-world behavior 42.
 - of correspondence 76.
- S-integral 75.
- second law of motion:
 - monadic 92.
 - monadic alternative 101.
- second order rates of change, comprehending 98.
- segments, hyperline 73.
- self-evident 49.
 - max. and min. 49.
 - method of constants 51.
- sequential convergence 19.
- sets, infinite 26.
- simple:
 - behavior 58.
 - fine partition 35.
- simplistic 76.
 - physical behavior 46.
- simply additive 37.
- sketching 46.
- small, infinitely 6.
- sovereign principle 18.
- specific heat 99.
- Spiegel 8.
- Spivak [1965] 35.
- * operator, properties of 17.
- *-transfer, how two use 22.
- *-transfer, proving things by 22.
- *-transfer 21.
- st operator 14.
- standard:
 - (unstarred) constants 23.
 - constants, extended 23.
 - constants, set of 20.
 - part operator, properties of 14.
 - part operator, using 25.
- steady state 98.
- Stroyan and Luxemburg [1976] 21.
- Struik [1961] 84.
- subrectangle 37.
 - boundary 62.
- subrectangle:
 - exterior 62.
 - inner 62.
- sum, hyperfinite 31.
- summation, finite 31.
- sums, Riemann styled 58.
- supernear 38.
- supernearness 38.
- superstructure 17.
- surface:
 - element 73.
 - integral 73.
 - σ 73.
 - tangent points 75.
- Swartz and Thomson [1988] 75.
- symbol for a monad 12.
- symbol for infinite natural numbers 19.
- symbol for infinitely close 11.
- symbol for limited 11.
- symbol hyperreal 10.
- symbol natural numbers 10.
- symbol positive real numbers 10.
- symbol real numbers 10.
- symbols for infinitesimals 10.
- Synge and Griffith [1959: 173] 49.
- tangent:
 - plane 73.
 - points to the surface 71.

- to a curve 71.
- vector to a curve 82.
- temperature 99.
- The Integral 36.
- thermal conductivity 99.
- truncated hyperrectangular solid 63.
- tubes:
 - of fluid motion 44.
 - of moving points 44.
- types 15.

- ultraproduct 21.
- uniform:
 - rod 49.
 - conducts heat 98.
- uniformly continuous 32.
- unit normal vector to the curve 84.
- unstarred constants 23.

- variable, free 28.
- vector:
 - normal 73.
 - tangent to a curve 82.
 - unit normal 84.
- VR1 44.
- VR2 44.

- Weierstrass 6.
- What if...? 25.
- work 67.

- Zeno 17.